



## Full wwPDB EM Validation Report ⓘ

Feb 13, 2025 – 10:17 AM EST

PDB ID : 9MU9  
EMDB ID : EMD-48626  
Title : Structure of a native *Drosophila melanogaster* Nucleosome Elongation Complex (Pol II EC-nucleosome). Composite map  
Authors : Venette-Smith, N.L.; Vishwakarma, R.K.; Dollinger, R.; Schultz, J.; Venkatakrishnan, V.; Babitzke, P.; Anand, G.; Gilmour, D.S.; Armache, J.-P.; Murakami, K.  
Deposited on : 2025-01-13  
Resolution : 7.80 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 7.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	10	100%
2	T	174	5% 99%
3	N	163	5% 100%
4	c	106	6% 99%
4	g	106	100%
5	d	97	100%
5	h	97	100%
6	a	100	100%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
6	e	100	 92%8%
7	b	80	 100%
7	f	80	 100%
8	A	1470	 100%
9	B	1164	 100%
10	C	269	 5%100%
11	F	84	 99%.
12	H	147	 7%99%.
13	I	116	 100%
14	J	67	 100%
15	K	116	 100%
16	L	46	 100%
17	E	210	 100%

## 2 Entry composition [i](#)

There are 18 unique types of molecules in this entry. The entry contains 42423 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	10	Total	C	N	O	P	0	0
			220	100	50	60	10		

- Molecule 2 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	T	174	Total	C	N	O	P	0	0
			3522	1695	567	1087	173		

- Molecule 3 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	N	163	Total	C	N	O	P	0	0
			3376	1598	673	942	163		

- Molecule 4 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	g	106	Total	C	N	O	S	0	0
			809	511	157	140	1		
4	c	105	Total	C	N	O	S	0	0
			803	506	156	140	1		

- Molecule 5 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	h	97	Total	C	N	O	S	0	0
			753	474	135	142	2		
5	d	97	Total	C	N	O	S	0	0
			753	474	135	142	2		

- Molecule 6 is a protein called Histone H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	e	92	Total	C	N	O	S	0	0
			749	471	142	133	3		
6	a	100	Total	C	N	O	S	0	0
			817	514	158	142	3		

- Molecule 7 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	f	80	Total	C	N	O	S	0	0
			630	395	123	111	1		
7	b	80	Total	C	N	O	S	0	0
			638	401	125	111	1		

- Molecule 8 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	A	1470	Total	C	N	O	S	0	0
			11565	7261	2052	2163	89		

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	B	1164	Total	C	N	O	S	0	0
			9257	5851	1633	1709	64		

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	C	269	Total	C	N	O	S	0	0
			2156	1354	362	432	8		

- Molecule 11 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	F	84	Total	C	N	O	S	0	0
			674	430	115	123	6		

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	H	147	Total	C	N	O	S	0	0
			1186	748	200	228	10		

- Molecule 13 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	I	116	Total	C	N	O	S	0	0
			947	588	164	183	12		

- Molecule 14 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	J	67	Total	C	N	O	S	0	0
			530	342	89	93	6		

- Molecule 15 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	K	116	Total	C	N	O	S	0	0
			935	603	153	176	3		

- Molecule 16 is a protein called DNA-directed RNA polymerase II subunit RPB12.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	L	46	Total	C	N	O	S	0	0
			390	239	78	66	7		

- Molecule 17 is a protein called DNA-directed RNA polymerase II subunit RPB5.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	E	210	Total	C	N	O	S	0	0
			1705	1081	290	322	12		

- Molecule 18 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
18	A	2	Total	Zn	0
			2	2	
18	B	1	Total	Zn	0
			1	1	
18	C	1	Total	Zn	0
			1	1	
18	I	2	Total	Zn	0
			2	2	
18	J	1	Total	Zn	0
			1	1	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
18	L	1	1	1	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA

Chain R:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: Template DNA

Chain T:  99%



- Molecule 3: Non-template DNA

Chain N:  100%




- Molecule 4: Histone H2A

Chain g:  100%



- Molecule 4: Histone H2A

Chain c:  99%



- Molecule 5: Histone H2B



Chain h:  100%

There are no outlier residues recorded for this chain.


- Molecule 5: Histone H2B

Chain d:  100%

There are no outlier residues recorded for this chain.


- Molecule 6: Histone H3

Chain e:  92% 8%



- Molecule 6: Histone H3

Chain a:  100%



- Molecule 7: Histone H4

Chain f:  100%

There are no outlier residues recorded for this chain.


- Molecule 7: Histone H4

Chain b:  100%




- Molecule 8: DNA-directed RNA polymerase II subunit RPB1

Chain A:  100%




- Molecule 9: DNA-directed RNA polymerase II subunit RPB2


Chain B:  100%



- Molecule 10: DNA-directed RNA polymerase II subunit RPB3




- Molecule 11: DNA-directed RNA polymerases I, II, and III subunit RPABC2



- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC3



- Molecule 13: DNA-directed RNA polymerase II subunit RPB9




- Molecule 14: DNA-directed RNA polymerases I, II, and III subunit RPABC5



There are no outlier residues recorded for this chain.

- Molecule 15: DNA-directed RNA polymerase II subunit RPB11



- Molecule 16: DNA-directed RNA polymerase II subunit RPB12



There are no outlier residues recorded for this chain.

- Molecule 17: DNA-directed RNA polymerase II subunit RPB5

Chain E:  100%

There are no outlier residues recorded for this chain.

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	36552	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; 'Patch CTF Estimation' in cryoSPARC	Depositor
Microscope	TFS TALOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50.65	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.572	Depositor
Minimum map value	-0.131	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.024	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	415.368, 415.368, 415.368	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1538, 1.1538, 1.1538	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	R	0.23	0/249	0.80	0/386
2	T	0.56	0/3927	1.06	1/6056 (0.0%)
3	N	0.54	0/3805	0.88	0/5871
4	c	0.25	0/813	0.53	0/1097
4	g	0.25	0/819	0.53	0/1104
5	d	0.25	0/764	0.50	0/1027
5	h	0.25	0/764	0.47	0/1027
6	a	0.26	0/829	0.55	0/1112
6	e	0.26	0/757	0.60	0/1013
7	b	0.26	0/645	0.56	0/862
7	f	0.25	0/637	0.58	0/854
8	A	0.27	1/11779 (0.0%)	0.55	3/15904 (0.0%)
9	B	0.26	0/9444	0.55	1/12756 (0.0%)
10	C	0.25	0/2201	0.50	0/2989
11	F	0.25	0/685	0.58	1/923 (0.1%)
12	H	0.26	0/1211	0.61	1/1627 (0.1%)
13	I	0.26	0/968	0.56	0/1311
14	J	0.27	0/539	0.54	0/726
15	K	0.27	0/954	0.51	0/1286
16	L	0.26	0/397	0.64	0/527
17	E	0.26	0/1735	0.55	0/2339
All	All	0.33	1/43922 (0.0%)	0.66	7/60797 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	150	CYS	C-N	5.32	1.46	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	B	260	ILE	C-N-CA	6.37	137.62	121.70
12	H	41	LEU	CA-CB-CG	5.97	129.04	115.30
2	T	-2	DC	O4'-C1'-N1	5.62	111.93	108.00
11	F	93	LEU	CA-CB-CG	5.61	128.20	115.30
8	A	1328	LEU	CA-CB-CG	5.47	127.89	115.30
8	A	540	MET	CA-CB-CG	5.35	122.40	113.30
8	A	461	MET	CA-CB-CG	5.21	122.15	113.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	c	103/106 (97%)	102 (99%)	1 (1%)	0	100	100
4	g	104/106 (98%)	102 (98%)	2 (2%)	0	100	100
5	d	95/97 (98%)	94 (99%)	1 (1%)	0	100	100
5	h	95/97 (98%)	94 (99%)	1 (1%)	0	100	100
6	a	98/100 (98%)	98 (100%)	0	0	100	100
6	e	90/100 (90%)	88 (98%)	2 (2%)	0	100	100
7	b	78/80 (98%)	77 (99%)	1 (1%)	0	100	100
7	f	78/80 (98%)	75 (96%)	3 (4%)	0	100	100
8	A	1468/1470 (100%)	1419 (97%)	48 (3%)	1 (0%)	48	83
9	B	1162/1164 (100%)	1120 (96%)	42 (4%)	0	100	100
10	C	267/269 (99%)	254 (95%)	13 (5%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
11	F	82/84 (98%)	79 (96%)	3 (4%)	0	100	100
12	H	145/147 (99%)	135 (93%)	10 (7%)	0	100	100
13	I	114/116 (98%)	112 (98%)	2 (2%)	0	100	100
14	J	65/67 (97%)	63 (97%)	2 (3%)	0	100	100
15	K	114/116 (98%)	110 (96%)	4 (4%)	0	100	100
16	L	44/46 (96%)	42 (96%)	2 (4%)	0	100	100
17	E	208/210 (99%)	200 (96%)	8 (4%)	0	100	100
All	All	4410/4455 (99%)	4264 (97%)	145 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	A	1122	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	c	81/82 (99%)	81 (100%)	0	100	100
4	g	81/82 (99%)	81 (100%)	0	100	100
5	d	80/84 (95%)	80 (100%)	0	100	100
5	h	80/84 (95%)	80 (100%)	0	100	100
6	a	85/87 (98%)	85 (100%)	0	100	100
6	e	79/87 (91%)	79 (100%)	0	100	100
7	b	65/65 (100%)	65 (100%)	0	100	100
7	f	63/65 (97%)	63 (100%)	0	100	100
8	A	1269/1290 (98%)	1269 (100%)	0	100	100
9	B	1003/1015 (99%)	1003 (100%)	0	100	100
10	C	244/244 (100%)	244 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	F	74/75 (99%)	74 (100%)	0	100	100
12	H	127/127 (100%)	127 (100%)	0	100	100
13	I	106/108 (98%)	106 (100%)	0	100	100
14	J	56/57 (98%)	56 (100%)	0	100	100
15	K	102/102 (100%)	102 (100%)	0	100	100
16	L	42/42 (100%)	42 (100%)	0	100	100
17	E	190/192 (99%)	190 (100%)	0	100	100
All	All	3827/3888 (98%)	3827 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	9/10 (90%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	N	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	N	-87:DA	O3'	-75:DA	P	23.10


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48626. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 145



Y Index: 173




Z Index: 190


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 844  $\text{nm}^3$ ; this corresponds to an approximate mass of 762 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.128 Å<sup>-1</sup>

## 8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.




## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-48626 and PDB model 9MU9. Per-residue inclusion information can be found in section [3](#) on page [8](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).













































## 9.4 Atom inclusion ⓘ



At the recommended contour level, 97% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9310	 0.2020
A	 0.9120	 0.1960
B	 0.9490	 0.1980
C	 0.8550	 0.2050
E	 0.9430	 0.1990
F	 0.9250	 0.1890
H	 0.8280	 0.2100
I	 0.9390	 0.1940
J	 0.9630	 0.2010
K	 0.8540	 0.1960
L	 0.9570	 0.2180
N	 0.9420	 0.2540
R	 1.0000	 0.2790
T	 0.9350	 0.2510
a	 0.9810	 0.1500
b	 0.9890	 0.1440
c	 0.9280	 0.1600
d	 0.9880	 0.1740
e	 0.9630	 0.1510
f	 0.9900	 0.1570
g	 0.9860	 0.1670
h	 0.9990	 0.1720

