



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 05:55 PM UTC

PDB ID : 5N5Z / pdb_00005n5z
EMDB ID : EMD-3591
Title : Cryo-EM structure of RNA polymerase I in complex with Rrn3 and Core Factor (Orientation II)
Authors : Engel, C.; Gubbey, T.; Neyer, S.; Sainsbury, S.; Oberthuer, C.; Baejen, C.; Bernecky, C.; Cramer, P.
Deposited on : 2017-02-14
Resolution : 7.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 7.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	384 (7.20 - 8.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1664	<p>7% (red), 60% (green), 24% (yellow), 12% (grey). Dot at 12%.</p>
2	B	1203	<p>9% (red), 62% (green), 31% (yellow), 2% (grey). Dots at 2% and 3%.</p>
3	C	335	<p>5% (red), 65% (green), 23% (yellow), 9% (grey). Dots at 2% and 3%.</p>
4	D	137	<p>31% (green), 9% (yellow), 58% (grey). Dots at 1% and 2%.</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	 73% 21%
6	F	155	 48% 14% 35%
7	G	326	 41% 16% 41%
8	H	146	 63% 25% 10%
9	I	125	 20% 52% 26% 21%
10	J	70	 60% 34%
11	K	142	 45% 23% 29%
12	L	70	 6% 30% 24% 9% 37%
13	M	415	 16% 9% 74%
14	N	233	 7% 41% 19% 38%
15	O	627	 36% 33% 6% 25%
16	P	894	 37% 20% 31% 12% 35%
17	Q	514	 53% 27% 43% 9% 20%
18	R	507	 23% 17% 33% 9% 40%

2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 48004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase I subunit RPA190.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1467	Total	C	N	O	S	0	0
			11598	7327	2017	2193	61		

- Molecule 2 is a protein called DNA-directed RNA polymerase I subunit RPA135.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1172	Total	C	N	O	S	0	0
			9312	5891	1633	1738	50		

- Molecule 3 is a protein called DNA-directed RNA polymerases I and III subunit RPAC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	305	Total	C	N	O	S	0	0
			2423	1539	416	460	8		

- Molecule 4 is a protein called DNA-directed RNA polymerase I subunit RPA14.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
4	D	58	Total	C	N	O	0	0
			459	289	78	92		

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	212	Total	C	N	O	S	0	0
			1735	1102	306	316	11		

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	100	Total	C	N	O	S	0	0
			823	522	144	154	3		

- Molecule 7 is a protein called DNA-directed RNA polymerase I subunit RPA43.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	193	1526	985	262	274	5	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	131	1052	664	176	208	4	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase I subunit RPA12.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	99	755	472	125	149	9	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	69	569	362	101	100	6	0	0

- Molecule 11 is a protein called DNA-directed RNA polymerases I and III subunit RPAC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	K	101	793	496	130	162	5	0	0

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	44	352	217	70	61	4	0	0

- Molecule 13 is a protein called DNA-directed RNA polymerase I subunit RPA49.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
13	M	108	856	543	142	171	0	0

- Molecule 14 is a protein called DNA-directed RNA polymerase I subunit RPA34.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	145	Total	C	N	O	S	0	0
			1151	735	188	224	4		

- Molecule 15 is a protein called RNA polymerase I-specific transcription initiation factor RRN3.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	473	Total	C	N	O	S	0	0
			3907	2533	642	711	21		

- Molecule 16 is a protein called RNA polymerase I-specific transcription initiation factor RRN6.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	583	Total	C	N	O	S	0	0
			4729	3010	803	905	11		

- Molecule 17 is a protein called RNA polymerase I-specific transcription initiation factor RRN7.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	410	Total	C	N	O	S	0	0
			3421	2219	579	603	20		

- Molecule 18 is a protein called RNA polymerase I-specific transcription initiation factor RRN11.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	303	Total	C	N	O	S	0	0
			2535	1634	456	434	11		

- Molecule 19 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
19	A	2	Total	Zn	0
			2	2	
19	B	1	Total	Zn	0
			1	1	
19	I	2	Total	Zn	0
			2	2	
19	J	1	Total	Zn	0
			1	1	

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms		AltConf
19	L	1	Total 1	Zn 1	0
19	Q	1	Total 1	Zn 1	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase I subunit RPA190






• Molecule 3: DNA-directed RNA polymerases I and III subunit RPAC1







- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 9: DNA-directed RNA polymerase I subunit RPA12





- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5




- Molecule 11: DNA-directed RNA polymerases I and III subunit RPAC2




- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4




• Molecule 13: DNA-directed RNA polymerase I subunit RPA49




• Molecule 14: DNA-directed RNA polymerase I subunit RPA34



• Molecule 15: RNA polymerase I-specific transcription initiation factor RRN3



• Molecule 16: RNA polymerase I-specific transcription initiation factor RRN6



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	8317	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.071	Depositor
Minimum map value	-0.035	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.018	Depositor
Map size (Å)	324.0, 324.0, 324.0	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.35, 1.35, 1.35	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.64	1/11809 (0.0%)	0.96	36/15943 (0.2%)
2	B	0.62	0/9518	0.98	35/12863 (0.3%)
3	C	0.53	0/2475	0.94	8/3354 (0.2%)
4	D	0.52	0/465	0.81	0/630
5	E	0.49	0/1771	0.88	3/2383 (0.1%)
6	F	0.58	0/838	0.86	0/1129
7	G	0.50	0/1563	0.84	3/2124 (0.1%)
8	H	0.52	0/1070	0.88	0/1449
9	I	0.53	0/765	0.87	0/1030
10	J	0.60	0/578	0.98	6/775 (0.8%)
11	K	0.54	0/804	0.93	6/1083 (0.6%)
12	L	0.50	0/354	0.91	2/468 (0.4%)
13	M	0.49	0/872	0.87	0/1170
14	N	0.49	0/1172	0.88	1/1580 (0.1%)
15	O	0.45	1/3996 (0.0%)	0.82	2/5401 (0.0%)
16	P	0.38	0/4822	0.74	6/6525 (0.1%)
17	Q	0.36	1/3501 (0.0%)	0.67	2/4724 (0.0%)
18	R	0.35	0/2592	0.77	5/3486 (0.1%)
All	All	0.54	3/48965 (0.0%)	0.89	115/66117 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	15
2	B	0	4
3	C	0	1
6	F	0	1
12	L	0	2

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
13	M	0	3
14	N	0	1
15	O	0	4
16	P	0	26
17	Q	0	8
18	R	0	10
All	All	0	75

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	356	PHE	C-N	8.48	1.44	1.33
17	Q	15	CYS	CA-C	7.48	1.62	1.52
15	O	212	THR	CA-CB	5.45	1.62	1.53

The worst 5 of 115 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
18	R	205	VAL	N-CA-C	16.04	125.61	110.53
18	R	208	TYR	N-CA-C	13.16	125.35	111.14
2	B	405	GLY	N-CA-C	-12.23	97.21	113.24
1	A	1071	ASP	N-CA-C	-9.86	97.51	110.53
3	C	223	SER	N-CA-C	-9.70	101.61	112.57

There are no chirality outliers.

5 of 75 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	405	LYS	Peptide
1	A	406	LEU	Peptide
1	A	407	GLN	Peptide
1	A	410	LYS	Peptide
1	A	411	VAL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11598	0	11664	1202	0
2	B	9312	0	9200	1013	0
3	C	2423	0	2412	138	0
4	D	459	0	462	7	0
5	E	1735	0	1764	34	0
6	F	823	0	839	91	0
7	G	1526	0	1534	158	0
8	H	1052	0	1021	56	0
9	I	755	0	728	106	0
10	J	569	0	584	55	0
11	K	793	0	790	54	0
12	L	352	0	371	59	0
13	M	856	0	852	60	0
14	N	1151	0	1168	166	0
15	O	3907	0	3902	417	0
16	P	4729	0	4674	595	0
17	Q	3421	0	3459	784	0
18	R	2535	0	2598	621	0
19	A	2	0	0	1	0
19	B	1	0	0	0	0
19	I	2	0	0	0	0
19	J	1	0	0	0	0
19	L	1	0	0	0	0
19	Q	1	0	0	0	0
All	All	48004	0	48022	4131	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 4131 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
18:R:320:CYS:SG	18:R:374:LEU:HG	1.11	1.67
1:A:995:TYR:CD2	2:B:708:ASP:HA	1.30	1.64
17:Q:20:TRP:CZ3	17:Q:22:ILE:CG2	1.76	1.64
17:Q:186:CYS:HA	18:R:208:TYR:CE1	1.31	1.64
17:Q:326:TYR:CE1	17:Q:452:PHE:CE2	1.87	1.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1449/1664 (87%)	1375 (95%)	60 (4%)	14 (1%)	12	48
2	B	1164/1203 (97%)	1095 (94%)	51 (4%)	18 (2%)	8	40
3	C	303/335 (90%)	288 (95%)	12 (4%)	3 (1%)	12	48
4	D	54/137 (39%)	50 (93%)	2 (4%)	2 (4%)	2	20
5	E	210/215 (98%)	197 (94%)	11 (5%)	2 (1%)	12	48
6	F	98/155 (63%)	95 (97%)	2 (2%)	1 (1%)	12	48
7	G	187/326 (57%)	173 (92%)	12 (6%)	2 (1%)	11	46
8	H	127/146 (87%)	121 (95%)	6 (5%)	0	100	100
9	I	91/125 (73%)	80 (88%)	8 (9%)	3 (3%)	3	21
10	J	67/70 (96%)	63 (94%)	4 (6%)	0	100	100
11	K	99/142 (70%)	92 (93%)	7 (7%)	0	100	100
12	L	42/70 (60%)	36 (86%)	4 (10%)	2 (5%)	2	16
13	M	106/415 (26%)	96 (91%)	8 (8%)	2 (2%)	6	32
14	N	139/233 (60%)	123 (88%)	13 (9%)	3 (2%)	5	29
15	O	467/627 (74%)	426 (91%)	35 (8%)	6 (1%)	9	42
16	P	569/894 (64%)	473 (83%)	76 (13%)	20 (4%)	3	20
17	Q	396/514 (77%)	345 (87%)	41 (10%)	10 (2%)	4	26
18	R	291/507 (57%)	245 (84%)	40 (14%)	6 (2%)	5	30
All	All	5859/7778 (75%)	5373 (92%)	392 (7%)	94 (2%)	10	38

5 of 94 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	408	LYS
1	A	479	ALA
1	A	1651	THR
2	B	111	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	895	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1297/1465 (88%)	1205 (93%)	92 (7%)	13	35
2	B	1025/1053 (97%)	962 (94%)	63 (6%)	17	38
3	C	269/296 (91%)	254 (94%)	15 (6%)	19	40
4	D	55/116 (47%)	50 (91%)	5 (9%)	9	27
5	E	194/197 (98%)	179 (92%)	15 (8%)	12	32
6	F	90/137 (66%)	85 (94%)	5 (6%)	19	40
7	G	171/291 (59%)	158 (92%)	13 (8%)	12	33
8	H	115/128 (90%)	107 (93%)	8 (7%)	14	35
9	I	89/110 (81%)	84 (94%)	5 (6%)	19	40
10	J	64/65 (98%)	56 (88%)	8 (12%)	4	16
11	K	91/130 (70%)	82 (90%)	9 (10%)	7	24
12	L	39/57 (68%)	34 (87%)	5 (13%)	4	15
13	M	98/371 (26%)	85 (87%)	13 (13%)	4	14
14	N	135/220 (61%)	128 (95%)	7 (5%)	21	42
15	O	439/576 (76%)	383 (87%)	56 (13%)	4	15
16	P	539/828 (65%)	386 (72%)	153 (28%)	0	2
17	Q	383/476 (80%)	320 (84%)	63 (16%)	2	10
18	R	286/474 (60%)	223 (78%)	63 (22%)	1	6
All	All	5379/6990 (77%)	4781 (89%)	598 (11%)	8	19

5 of 598 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
16	P	748	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
18	R	266	SER
17	Q	101	LYS
16	P	736	ILE
17	Q	379	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 74 such sidechains are listed below:

Mol	Chain	Res	Type
16	P	226	HIS
18	R	223	ASN
16	P	368	HIS
17	Q	351	ASN
2	B	893	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2
7	G	1
9	I	1
17	Q	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	G	24:VAL	C	25:THR	N	4.53
1	I	45:LEU	C	46:LYS	N	3.17
1	Q	369:TRP	C	370:SER	N	3.11
1	A	438:ILE	C	439:ASP	N	2.97
1	A	991:LYS	C	992:PRO	N	2.87


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3591. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 105



Y Index: 102




Z Index: 122


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.018. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 591 nm³; this corresponds to an approximate mass of 534 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.130\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-3591 and PDB model 5N5Z. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.018 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)

This section was not generated.







































9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 66% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.018) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6620	 0.1160
A	 0.7270	 0.1310
B	 0.7230	 0.1220
C	 0.7900	 0.1330
D	 0.7980	 0.1690
E	 0.8020	 0.1410
F	 0.7480	 0.1420
G	 0.7940	 0.1310
H	 0.8430	 0.1530
I	 0.6210	 0.1250
J	 0.7300	 0.1190
K	 0.7930	 0.1530
L	 0.7440	 0.0950
M	 0.7180	 0.1130
N	 0.7280	 0.1420
O	 0.7620	 0.1370
P	 0.3710	 0.0470
Q	 0.3000	 0.0550
R	 0.4930	 0.1040

