

# wwPDB EM Validation Summary Report (i)

#### Nov 3, 2025 – 12:46 pm GMT

PDB ID : 7NAW / pdb 00007naw

EMDB ID : EMD-12250

Title: Bacterial 30S ribosomal subunit assembly complex state B (Consensus Refine-

ment)

Authors : Schedlbauer, A.; Iturrioz, I.; Ochoa-Lizarralde, B.; Diercks, T.; Kaminishi, T.;

Capuni, R.; Astigarraga, E.; Gil-Carton, D.; Fucini, P.; Connell, S.

Deposited on : 2021-01-25

Resolution : 4.05 Å(reported)

Based on initial model : 4YBB

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev129

Mogul : 1.8.4, CSD as541be (2020) MolProbity : 4-5-2 with Phenix2.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023) EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 4.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive	EM structures	Similar EM resolution
Metric	$(\#  ext{Entries})$	$(\# \mathbf{Entries})$	$(\#  ext{Entries},  ext{ resolution range}(\mathring{ ext{A}}))$
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	6569 ( 3.55 - 4.55 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length		Quality of chain								
1	A	1542		41%		36%	13%	• 7%				
2	В	241	17%		78%		15%	7%				
3	С	233	•		85%		6%	5 9%				







## 2 Entry composition (i)

There are 25 unique types of molecules in this entry. The entry contains 52652 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 16S rRNA.

Mol	Chain	Residues		A	AltConf	Trace			
1	A	1427	Total 30625	C 13663	N 5617	O 9918	P 1427	0	0

• Molecule 2 is a protein called Small ribosomal subunit protein uS2.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
2	В	224	Total 1753	C 1109	N 315	O 321	S 8	0	0

• Molecule 3 is a protein called Small ribosomal subunit protein uS3.

Mol	Chain	Residues		Ato	AltConf	Trace			
3	С	211	Total 1653	C 1046	N 310	O 293	S 4	0	0

• Molecule 4 is a protein called Small ribosomal subunit protein uS4.

Mol	Chain	Residues		Ato	AltConf	Trace			
4	D	205	Total 1643	C 1026	N 315	O 298	S 4	0	0

• Molecule 5 is a protein called Small ribosomal subunit protein uS5.

Mol	Chain	Residues		At	oms			AltConf	Trace
5	Е	155	Total 1144	C 711	N 216	O 211	S 6	0	0

• Molecule 6 is a protein called Small ribosomal subunit protein bS6, fully modified isoform.

Mol	Chain	Residues		At	oms	AltConf	Trace		
6	F	106	Total 862	C 545	N 156	O 154	S 7	0	0



• Molecule 7 is a protein called Small ribosomal subunit protein uS7.

Mol	Chain	Residues		At	oms		AltConf	Trace	
7	G	151	Total 1181	C 735	N 227	O 215	S 4	0	0

• Molecule 8 is a protein called Small ribosomal subunit protein uS8.

Mol	Chain	Residues		At	oms	AltConf	Trace		
0	П	129	Total	С	N	О	S	0	0
0	П	129	979	616	173	184	6	0	0

• Molecule 9 is a protein called Small ribosomal subunit protein uS9.

Mol	Chain	Residues		At	oms			AltConf	Trace
9	I	127	Total 1022	C 634	N 206	O 179	S 3	0	0

• Molecule 10 is a protein called Small ribosomal subunit protein uS10.

Mol	Chain	Residues		At	oms	AltConf	Trace		
10	J	99	Total	С	N	0	S	0	0
			795	498	152	144	1		_

• Molecule 11 is a protein called Small ribosomal subunit protein uS11.

Mol	Chain	Residues		At	oms			AltConf	Trace
11	K	117	Total 877	C 540	N 174	O 160	S 3	0	0

• Molecule 12 is a protein called Small ribosomal subunit protein uS12.

Mol	Chain	Residues		At	oms			AltConf	Trace
12	L	123	Total 957	C 591	N 196	O 165	S 5	0	0

• Molecule 13 is a protein called Small ribosomal subunit protein uS13.

$\mathbf{N}$	Iol	Chain	Residues		${f Atoms}$					Trace
	13	M	114	Total 883	C 546	N 178	O 156	S 3	0	0

• Molecule 14 is a protein called Small ribosomal subunit protein uS14.



Mol	Chain	Residues		Atoms					Trace
14	N	100	Total 805	C 499	N 164	O 139	S 3	0	0

• Molecule 15 is a protein called Small ribosomal subunit protein uS15.

Mol	Chain	Residues		At	oms	AltConf	Trace		
15	0	88	Total	С	N	О	S	0	0
			714	439	144	130	1		

• Molecule 16 is a protein called Small ribosomal subunit protein bS16.

Mol	Chain	Residues		At	oms	AltConf	Trace		
16	D	82	Total	С	N	О	S	0	0
10	1	02	649	406	128	114	1	0	

• Molecule 17 is a protein called Small ribosomal subunit protein uS17.

Mol	Chain	Residues		At	oms	AltConf	Trace		
17	0	90	Total	С	N	О	S	0	0
11	Q	80	648	411	121	113	3	U	U

• Molecule 18 is a protein called Small ribosomal subunit protein bS18.

Mol	Chain	Residues		Ato	oms	AltConf	Trace		
18	R	65	Total 535	C 339	N 100	O 95	S 1	0	0

• Molecule 19 is a protein called Small ribosomal subunit protein uS19.

Mol	Chain	Residues		At	oms	AltConf	Trace		
19	S	82	Total	C	N	0	S	0	0
			658	421	125	110	2		

• Molecule 20 is a protein called Small ribosomal subunit protein bS20.

Mol	Chain	Residues		At	oms	AltConf	Trace		
20	Т	86	Total		N 120	0	S	0	0
			670	414	138	115	3		

• Molecule 21 is a protein called Small ribosomal subunit protein bS21.



Mol	Chain	Residues		Ato	ms	AltConf	Trace		
21	U	52	Total	С	N	О	S	0	0
		Ŭ <b>-</b>	430	269	89	71	1		Ü

• Molecule 22 is a protein called Ribosome maturation factor RimP.

Mol	Chain	Residues		At	oms	AltConf	Trace		
22	v	151	Total	С	N	О	S	0	0
22	Λ	151	1174	742	197	230	5	0	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	151	ALA	-	expression tag	UNP A0A0J3VRH1

• Molecule 23 is a protein called Ribosomal RNA small subunit methyltransferase A.

Mol	Chain	Residues	Atoms				AltConf	Trace	
23	Y	252	Total 1963	C 1252	N 338	O 360	S 13	0	0

• Molecule 24 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
24	A	31	Total Mg 31 31	0

• Molecule 25 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	AltConf
25	В	1	Total Zn 1 1	0




## 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 16S rRNA







 $\bullet$  Molecule 2: Small ribosomal subunit protein uS2














• Molecule 13: Small ribosomal subunit protein uS13

Chain M: 93%

• Molecule 14: Small ribosomal subunit protein uS14


Chain N: 95%

• Molecule 15: Small ribosomal subunit protein uS15

Chain O: 94%


• Molecule 16: Small ribosomal subunit protein bS16





• Molecule 23: Ribosomal RNA small subunit methyltransferase A







# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	23278	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	42	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2250	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.096	Depositor
Minimum map value	-0.022	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.016	Depositor
Map size (Å)	416.64, 416.64, 416.64	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.085, 1.085, 1.085	Depositor



## 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 5MC, MA6, MG, ZN, D2T, 2MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N/L-1	Clasica	Во	nd lengths	Е	Bond angles
Mol	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
1	A	0.84	0/34130	1.28	$154/53237 \ (0.3\%)$
2	В	0.52	0/1784	0.83	0/2403
3	С	0.73	0/1680	1.24	1/2263 (0.0%)
4	D	0.74	0/1665	1.24	0/2227
5	Е	0.74	0/1157	1.31	1/1557 (0.1%)
6	F	0.73	0/881	1.27	2/1189 (0.2%)
7	G	0.82	1/1195 (0.1%)	1.32	2/1602 (0.1%)
8	Н	0.72	0/989	1.21	3/1326 (0.2%)
9	I	0.77	0/1034	1.26	1/1375 (0.1%)
10	J	0.78	0/805	1.37	7/1089 (0.6%)
11	K	0.87	0/893	1.56	9/1205 (0.7%)
12	L	0.83	0/960	1.36	3/1286 (0.2%)
13	M	0.83	0/892	1.30	1/1193 (0.1%)
14	N	0.78	0/817	1.26	0/1088
15	O	0.75	0/722	1.26	2/964 (0.2%)
16	Р	0.79	0/659	1.32	0/884
17	Q	0.73	0/657	1.29	2/881 (0.2%)
18	R	0.79	0/544	1.30	0/731
19	S	0.77	0/675	1.35	3/908 (0.3%)
20	Т	0.73	0/676	1.28	0/895
21	U	0.83	0/436	1.22	0/577
22	X	0.73	0/1192	1.33	3/1619 (0.2%)
23	Y	0.79	1/2006 (0.0%)	1.34	6/2730 (0.2%)
All	All	0.80	$2/56449 \ (0.0\%)$	1.28	200/83229 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.



Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	436
2	В	0	1
3	С	0	1
4	D	0	3
5	Е	0	1
6	F	0	2
7	G	0	1
8	Н	0	1
9	I	0	4
11	K	0	1
12	L	0	2
14	N	0	1
20	Т	0	1
23	Y	0	5
All	All	0	460

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
23	Y	229	LEU	C-O	-6.35	1.15	1.24
7	G	2	PRO	N-CD	5.18	1.55	1.47

The worst 5 of 200 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	A	1242	G	C3'-C2'-O2'	10.52	126.48	110.70
1	A	1208	С	C2'-C3'-O3'	10.33	129.19	113.70
1	A	1303	С	C4'-C3'-O3'	9.69	127.54	113.00
10	J	57	VAL	N-CA-C	9.46	129.01	109.34
10	J	56	HIS	N-CA-C	-9.28	95.97	110.36

There are no chirality outliers.

5 of 460 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	A	Sidechain
1	A	11	G	Sidechain
1	A	15	G	Sidechain
1	A	2	A	Sidechain
1	A	4	U	Sidechain



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	30625	0	15353	166	0
2	В	1753	0	1780	26	0
3	С	1653	0	1727	15	0
4	D	1643	0	1707	1	0
5	Ε	1144	0	1175	16	0
6	F	862	0	864	5	0
7	G	1181	0	1238	2	0
8	Н	979	0	1031	1	0
9	I	1022	0	1070	6	0
10	J	795	0	836	5	0
11	K	877	0	887	4	0
12	L	957	0	1017	7	0
13	M	883	0	941	1	0
14	N	805	0	844	2	0
15	О	714	0	734	3	0
16	Р	649	0	666	1	0
17	Q	648	0	691	0	0
18	R	535	0	552	0	0
19	S	658	0	683	1	0
20	Т	670	0	719	1	0
21	U	430	0	458	0	0
22	X	1174	0	1174	2	0
23	Y	1963	0	1977	6	0
24	A	31	0	0	0	0
25	В	1	0	0	0	0
All	All	52652	0	38124	236	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 236 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:1081:A:C8	5:E:23:LYS:NZ	2.13	1.12
1:A:920:U:H5'	1:A:1080:A:H61	0.94	1.09



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:U:O2	3:C:126:ARG:NE	1.86	1.08
1:A:421:U:O2	3:C:126:ARG:CZ	2.02	1.08
1:A:920:U:H5'	1:A:1080:A:N6	1.70	1.04

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
2	В	222/241 (92%)	218 (98%)	4 (2%)	0	100	100
3	С	209/233 (90%)	201 (96%)	8 (4%)	0	100	100
4	D	203/206 (98%)	198 (98%)	5 (2%)	0	100	100
5	E	153/167 (92%)	143 (94%)	10 (6%)	0	100	100
6	F	104/135 (77%)	104 (100%)	0	0	100	100
7	G	149/179 (83%)	137 (92%)	12 (8%)	0	100	100
8	Н	127/130 (98%)	123 (97%)	4 (3%)	0	100	100
9	I	125/130 (96%)	118 (94%)	7 (6%)	0	100	100
10	J	97/103 (94%)	92 (95%)	5 (5%)	0	100	100
11	K	115/129 (89%)	108 (94%)	7 (6%)	0	100	100
12	L	120/124 (97%)	113 (94%)	7 (6%)	0	100	100
13	M	112/118 (95%)	107 (96%)	5 (4%)	0	100	100
14	N	98/101 (97%)	95 (97%)	3 (3%)	0	100	100
15	О	86/89 (97%)	84 (98%)	2 (2%)	0	100	100
16	Р	80/82 (98%)	76 (95%)	4 (5%)	0	100	100
17	Q	78/84 (93%)	72 (92%)	6 (8%)	0	100	100
18	R	63/75 (84%)	61 (97%)	2 (3%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
19	S	80/92 (87%)	79 (99%)	1 (1%)	0	100	100
20	Τ	84/87 (97%)	83 (99%)	1 (1%)	0	100	100
21	U	50/71 (70%)	50 (100%)	0	0	100	100
22	X	149/151 (99%)	139 (93%)	10 (7%)	0	100	100
23	Y	250/273~(92%)	240 (96%)	10 (4%)	0	100	100
All	All	2754/3000 (92%)	2641 (96%)	113 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	В	186/199~(94%)	186 (100%)	(100%) 0		100	
3	C	$172/190\ (90\%)$	168 (98%)	4 (2%)	45	65	
4	D	172/173~(99%)	172 (100%)	0	100	100	
5	E	$118/126\ (94\%)$	118 (100%)	0	100	100	
6	F	92/116~(79%)	92 (100%)	0	100	100	
7	G	124/147~(84%)	122 (98%)	2 (2%)	58	74	
8	Н	$104/105\ (99\%)$	104 (100%)	0	100	100	
9	I	105/107 (98%)	105 (100%)	0	100	100	
10	J	87/90~(97%)	86 (99%)	1 (1%)	70	80	
11	K	90/99 (91%)	86 (96%)	4 (4%)	24	47	
12	L	$102/103\ (99\%)$	101 (99%)	1 (1%)	73	82	
13	M	92/96 (96%)	91 (99%)	1 (1%)	70	80	
14	N	83/84~(99%)	82 (99%)	1 (1%)	67	79	
15	О	76/77 (99%)	76 (100%)	0	100	100	
16	Р	65/65~(100%)	65 (100%)	0	100	100	
17	Q	74/78~(95%)	74 (100%)	0	100	100	



Mol	Chain	Analysed	Rotameric	Rotameric Outliers		Percentiles		
18	R	56/65~(86%)	55 (98%)	1 (2%)	54	72		
19	S	72/79 (91%)	72 (100%)	0	100	100		
20	Т	65/66 (98%)	65 (100%)	0	100	100		
21	U	43/61 (70%)	43 (100%)	0	100	100		
22	X	130/130 (100%)	129 (99%)	1 (1%)	79	85		
23	Y	216/234 (92%)	213 (99%)	3 (1%)	62	76		
All	All	2324/2490 (93%)	2305 (99%)	19 (1%)	77	85		

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
18	R	33	ILE
23	Y	51	LEU
23	Y	222	ARG
23	Y	42	VAL
11	K	36	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
19	S	53	ASN
19	S	52	HIS
7	G	52	GLN
18	R	54	GLN
6	F	55	HIS

#### 5.3.3 RNA (i)

$\mathbf{Mol}$	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	1425/1542 (92%)	310 (21%)	61 (4%)

5 of 310 RNA backbone outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type
1	A	2	A
1	A	4	U
1	A	9	G
1	A	32	A



Mol	Chain	Res	Type
1	A	39	G

5 of 61 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	812	G
1	A	1332	A
1	A	983	A
1	A	1286	U
1	A	1531	A

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	2MG	A	1207	1	18,26,27	1.10	1 (5%)	16,38,41	1.26	3 (18%)
12	D2T	L	89	12	7,9,10	1.97	3 (42%)	6,11,13	3.49	3 (50%)
1	MA6	A	1518	1	18,26,27	1.06	1 (5%)	19,38,41	2.01	7 (36%)
1	2MG	A	1516	1	18,26,27	0.97	1 (5%)	16,38,41	1.69	5 (31%)
1	5MC	A	967	1	18,22,23	0.98	2 (11%)	26,32,35	1.29	3 (11%)
1	2MG	A	966	1	18,26,27	1.00	1 (5%)	16,38,41	1.37	3 (18%)
1	MA6	A	1519	1	18,26,27	1.17	1 (5%)	19,38,41	2.01	7 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2MG	A	1207	1	-	0/5/27/28	0/3/3/3



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	D2T	L	89	12	-	2/7/12/14	-
1	MA6	A	1518	1	-	4/7/29/30	0/3/3/3
1	2MG	A	1516	1	-	3/5/27/28	0/3/3/3
1	5MC	A	967	1	-	3/7/25/26	0/2/2/2
1	2MG	A	966	1	-	0/5/27/28	0/3/3/3
1	MA6	A	1519	1	-	2/7/29/30	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	Ideal(Å)
12	L	89	D2T	OD2-CG	-2.93	1.21	1.30
12	L	89	D2T	OD1-CG	2.91	1.31	1.22
12	L	89	D2T	O-C	2.78	1.31	1.19
1	A	1207	2MG	C6-N1	-2.63	1.34	1.37
1	A	967	5MC	C6-C5	2.53	1.38	1.34

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
12	L	89	D2T	OD1-CG-CB	-5.43	111.06	122.44
12	L	89	D2T	OD2-CG-CB	5.26	124.52	113.15
1	A	1519	MA6	N1-C6-N6	4.36	121.64	117.06
1	A	1516	2MG	CM2-N2-C2	-3.78	115.51	123.86
1	A	1518	MA6	N1-C6-N6	3.52	120.77	117.06

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	L	89	D2T	CA-CB-SB-CB1
1	A	1516	2MG	C4'-C5'-O5'-P
1	A	1519	MA6	C5-C6-N6-C9
1	A	967	5MC	O4'-C4'-C5'-O5'
1	A	967	5MC	C3'-C4'-C5'-O5'

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1518	MA6	1	0



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	967	5MC	3	0
1	A	966	2MG	2	0
1	A	1519	MA6	3	0

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 32 ligands modelled in this entry, 32 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-12250. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




## 6.2 Central slices (i)


#### 6.2.1 Primary map








Y Index: 192




Z Index: 192


#### 6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192

The images above show central slices of the map in three orthogonal directions.




#### 6.3 Largest variance slices (i)


#### 6.3.1 Primary map








Y Index: 188




Z Index: 175


#### 6.3.2 Raw map



X Index: 191



Y Index: 188




Z Index: 175

The images above show the largest variance slices of the map in three orthogonal directions.




### 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



#### 6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.016. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




### 6.6 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


#### $6.6.1 \quad \mathrm{emd}\_12250\_\mathrm{msk}\_1.\mathrm{map} \ \ \mathbf{\mathring{1}}$









 $6.6.2 \quad \mathrm{emd}\_12250\_\mathrm{msk}\_2.\mathrm{map} \ \ \mathbf{\hat{1}}$ 








## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



#### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $858~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $775~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.247  $\rm \AA^{-1}$ 



## 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.247  $\rm \mathring{A}^{-1}$ 



## 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	4.05	-	-	
Author-provided FSC curve	4.03	5.43	4.10	
Unmasked-calculated*	7.11	10.75	7.50	


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.11 differs from the reported value 4.05 by more than 10 %



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-12250 and PDB model 7NAW. Per-residue inclusion information can be found in section 3 on page 8.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.016 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.016).



### 9.4 Atom inclusion (i)



At the recommended contour level, 98% of all backbone atoms, 94% of all non-hydrogen atoms, are inside the map.



### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.016) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9430	0.2970
A	0.9950	0.3280
В	0.6620	0.1870
С	0.8600	0.2700
D	0.9320	0.3150
Е	0.8920	0.3300
F	0.9020	0.2820
G	0.6260	0.1530
Н	0.9050	0.3360
I	0.9070	0.2190
J	0.9060	0.2310
K	0.9270	0.2520
L	0.8720	0.3040
M	0.8940	0.1640
N	0.9120	0.2440
О	0.9320	0.2920
Р	0.9490	0.3540
Q	0.9350	0.3370
R	0.8950	0.2810
S	0.9420	0.1870
Т	0.9620	0.2890
U	0.8150	0.2620
X	0.9000	0.2080
Y	0.8760	0.2100



