



# wwPDB EM Validation Summary Report i

Jun 29, 2025 – 05:04 am BST

PDB ID : 7NP3 / pdb\_00007np3  
EMDB ID : EMD-12512  
Title : cAMP-free rabbit HCN4 stabilized in LMNG-CHS detergent mixture  
Authors : Giese, H.M.; Chaves-Sanjuan, A.; Saponaro, A.; Clarke, O.; Bolognesi, M.; Mancia, F.; Hendrickson, W.A.; Thiel, G.; Santoro, B.; Moroni, A.  
Deposited on : 2021-02-26  
Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118  
MolProbity : 4-5-2 with Phenix2.0rc1  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *ELECTRON MICROSCOPY*


The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There is only 1 type of molecule in this entry. The entry contains 16132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4, Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	490	Total	C 4033	N 2614	O 679	S 708	32	0
1	B	490	Total	C 4033	N 2614	O 679	S 708	32	0
1	C	490	Total	C 4033	N 2614	O 679	S 708	32	0
1	D	490	Total	C 4033	N 2614	O 679	S 708	32	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP Q9TV66
A	?	-	LEU	deletion	UNP Q9TV66
A	821	ALA	-	insertion	UNP Q9TV66
A	822	GLY	-	insertion	UNP Q9TV66
A	823	GLY	-	insertion	UNP Q9TV66
A	824	PRO	-	insertion	UNP Q9TV66
A	?	-	PRO	deletion	UNP Q9TV66
A	?	-	ALA	deletion	UNP Q9TV66
A	?	-	GLY	deletion	UNP Q9TV66
A	?	-	GLY	deletion	UNP Q9TV66
B	?	-	LEU	deletion	UNP Q9TV66
B	?	-	LEU	deletion	UNP Q9TV66
B	821	ALA	-	insertion	UNP Q9TV66
B	822	GLY	-	insertion	UNP Q9TV66
B	823	GLY	-	insertion	UNP Q9TV66
B	824	PRO	-	insertion	UNP Q9TV66
B	?	-	PRO	deletion	UNP Q9TV66
B	?	-	ALA	deletion	UNP Q9TV66
B	?	-	GLY	deletion	UNP Q9TV66
B	?	-	GLY	deletion	UNP Q9TV66

*Continued on next page...*


*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	LEU	deletion	UNP Q9TV66
C	?	-	LEU	deletion	UNP Q9TV66
C	821	ALA	-	insertion	UNP Q9TV66
C	822	GLY	-	insertion	UNP Q9TV66
C	823	GLY	-	insertion	UNP Q9TV66
C	824	PRO	-	insertion	UNP Q9TV66
C	?	-	PRO	deletion	UNP Q9TV66
C	?	-	ALA	deletion	UNP Q9TV66
C	?	-	GLY	deletion	UNP Q9TV66
C	?	-	GLY	deletion	UNP Q9TV66
D	?	-	LEU	deletion	UNP Q9TV66
D	?	-	LEU	deletion	UNP Q9TV66
D	821	ALA	-	insertion	UNP Q9TV66
D	822	GLY	-	insertion	UNP Q9TV66
D	823	GLY	-	insertion	UNP Q9TV66
D	824	PRO	-	insertion	UNP Q9TV66
D	?	-	PRO	deletion	UNP Q9TV66
D	?	-	ALA	deletion	UNP Q9TV66
D	?	-	GLY	deletion	UNP Q9TV66
D	?	-	GLY	deletion	UNP Q9TV66


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4, Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4













- Molecule 1: Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 4



SER	MET	ASP	ALA	LYS	LEU	PRO
THR	GLN	GLN	ALA	ALA	ALA	ALA
GLU	E242	ALA	ALA	ALA	ALA	ALA
V241	R243	ARG	ALA	ALA	ALA	ALA
I319	V320	PRO	ALA	ALA	ALA	ALA
L387	V321	ALA	ALA	ALA	ALA	ALA
R388	E322	ALA	ALA	ALA	ALA	ALA
M509	E323	ALA	ALA	ALA	ALA	ALA
F610	Q245	ASP	ALA	ALA	ALA	ALA
V605	E246	ALA	ALA	ALA	ALA	ALA
D638	R247	VAL	ALA	ALA	ALA	ALA
Y637	T255	LYS	ALA	ALA	ALA	ALA
W638	V248	VAL	ALA	ALA	ALA	ALA
N324	K249	ASP	ALA	ALA	ALA	ALA
H400	S250	ALA	ALA	ALA	ALA	ALA
I518	A251	GLY	ALA	ALA	ALA	ALA
M608	L329	ALA	ALA	ALA	ALA	ALA
F690	T405	GLY	ALA	ALA	ALA	ALA
M691	Q401	ALA	ALA	ALA	ALA	ALA
GLY	D326	VAL	ALA	ALA	ALA	ALA
ALA	D327	ASP	ALA	ALA	ALA	ALA
GLU	E403	ALA	ALA	ALA	ALA	ALA
ALA	E404	GLY	ALA	ALA	ALA	ALA
VAL	R525	ALA	ALA	ALA	ALA	ALA
GLY	R526	GLY	ALA	ALA	ALA	ALA
ALA	Q527	ALA	ALA	ALA	ALA	ALA
ARG	F406	ALA	ALA	ALA	ALA	ALA
ALA	F407	ALA	ALA	ALA	ALA	ALA
GLY	P331	GLY	ALA	ALA	ALA	ALA
ALA	M332	ALA	ALA	ALA	ALA	ALA
ALA	M333	ALA	ALA	ALA	ALA	ALA
LYS	T409	ALA	ALA	ALA	ALA	ALA
ALA	R533	ALA	ALA	ALA	ALA	ALA
SER	Y338	ALA	ALA	ALA	ALA	ALA
ALA	L339	ALA	ALA	ALA	ALA	ALA
GLU	D261	ALA	ALA	ALA	ALA	ALA
ALA	F262	ALA	ALA	ALA	ALA	ALA
GLY	R263	ALA	ALA	ALA	ALA	ALA
ALA	K335	ALA	ALA	ALA	ALA	ALA
PRO	P258	ALA	ALA	ALA	ALA	ALA
GLY	P259	ALA	ALA	ALA	ALA	ALA
ALA	S260	ALA	ALA	ALA	ALA	ALA
ALA	D261	ALA	ALA	ALA	ALA	ALA
GLY	F262	ALA	ALA	ALA	ALA	ALA
ALA	H257	ALA	ALA	ALA	ALA	ALA
ALA	Y334	ALA	ALA	ALA	ALA	ALA
ALA	D330	ALA	ALA	ALA	ALA	ALA
ALA	P263	ALA	ALA	ALA	ALA	ALA
ALA	V254	ALA	ALA	ALA	ALA	ALA
ALA	E412	ALA	ALA	ALA	ALA	ALA
ALA	A413	ALA	ALA	ALA	ALA	ALA
ALA	S414	ALA	ALA	ALA	ALA	ALA
ALA	L412	ALA	ALA	ALA	ALA	ALA
ALA	V535	ALA	ALA	ALA	ALA	ALA
ALA	E536	ALA	ALA	ALA	ALA	ALA
ALA	Q537	ALA	ALA	ALA	ALA	ALA
ALA	K543	ALA	ALA	ALA	ALA	ALA
ALA	L544	ALA	ALA	ALA	ALA	ALA
ALA	D545	ALA	ALA	ALA	ALA	ALA
ALA	P546	ALA	ALA	ALA	ALA	ALA
ALA	R549	ALA	ALA	ALA	ALA	ALA
ALA	K627	ALA	ALA	ALA	ALA	ALA
ALA	K632	ALA	ALA	ALA	ALA	ALA
ALA	M633	ALA	ALA	ALA	ALA	ALA
ALA	T638	ALA	ALA	ALA	ALA	ALA
ALA	I624	ALA	ALA	ALA	ALA	ALA
ALA	R625	ALA	ALA	ALA	ALA	ALA
ALA	G630	ALA	ALA	ALA	ALA	ALA
ALA	E628	ALA	ALA	ALA	ALA	ALA
ALA	F631	ALA	ALA	ALA	ALA	ALA
ALA	K639	ALA	ALA	ALA	ALA	ALA
ALA	T645	ALA	ALA	ALA	ALA	ALA
ALA	F704	ALA	ALA	ALA	ALA	ALA
ALA	M700	ALA	ALA	ALA	ALA	ALA
ALA	R701	ALA	ALA	ALA	ALA	ALA
ALA	T706	ALA	ALA	ALA	ALA	ALA
ALA	L644	ALA	ALA	ALA	ALA	ALA
ALA	T645	ALA	ALA	ALA	ALA	ALA
ALA	K646	ALA	ALA	ALA	ALA	ALA
ALA	M652	ALA	ALA	ALA	ALA	ALA
ALA	H653	ALA	ALA	ALA	ALA	ALA
ALA	D654	ALA	ALA	ALA	ALA	ALA
ALA	Y655	ALA	ALA	ALA	ALA	ALA
ALA	Y656	ALA	ALA	ALA	ALA	ALA
ALA	E657	ALA	ALA	ALA	ALA	ALA
ALA	L658	ALA	ALA	ALA	ALA	ALA
ALA	K659	ALA	ALA	ALA	ALA	ALA
ALA	S660	ALA	ALA	ALA	ALA	ALA
ALA	Y661	ALA	ALA	ALA	ALA	ALA
ALA	K662	ALA	ALA	ALA	ALA	ALA
ALA	D665	ALA	ALA	ALA	ALA	ALA
ALA	E666	ALA	ALA	ALA	ALA	ALA
ALA	E667	ALA	ALA	ALA	ALA	ALA
ALA	R668	ALA	ALA	ALA	ALA	ALA
ALA	S669	ALA	ALA	ALA	ALA	ALA
ALA	T673	ALA	ALA	ALA	ALA	ALA
ALA	L574	ALA	ALA	ALA	ALA	ALA
ALA	S675	ALA	ALA	ALA	ALA	ALA
ALA	T676	ALA	ALA	ALA	ALA	ALA
ALA	Y678	ALA	ALA	ALA	ALA	ALA
ALA	T682	ALA	ALA	ALA	ALA	ALA
ALA	M587	ALA	ALA	ALA	ALA	ALA
ALA	R588	ALA	ALA	ALA	ALA	ALA
ALA	E681	ALA	ALA	ALA	ALA	ALA
ALA	Y482	ALA	ALA	ALA	ALA	ALA
ALA	I684	ALA	ALA	ALA	ALA	ALA
ALA	V355	ALA	ALA	ALA	ALA	ALA
ALA	I356	ALA	ALA	ALA	ALA	ALA
ALA	I357	ALA	ALA	ALA	ALA	ALA
ALA	P358	ALA	ALA	ALA	ALA	ALA
ALA	V359	ALA	ALA	ALA	ALA	ALA
ALA	F287	ALA	ALA	ALA	ALA	ALA
ALA	F288	ALA	ALA	ALA	ALA	ALA
ALA	T293	ALA	ALA	ALA	ALA	ALA
ALA	Q294	ALA	ALA	ALA	ALA	ALA
ALA	P295	ALA	ALA	ALA	ALA	ALA
ALA	P296	ALA	ALA	ALA	ALA	ALA
ALA	T297	ALA	ALA	ALA	ALA	ALA
ALA	R298	ALA	ALA	ALA	ALA	ALA
ALA	M299	ALA	ALA	ALA	ALA	ALA
ALA	K299	ALA	ALA	ALA	ALA	ALA
ALA	F299	ALA	ALA	ALA	ALA	ALA
ALA	Q299	ALA	ALA	ALA	ALA	ALA
ALA	P300	ALA	ALA	ALA	ALA	ALA
ALA	S301	ALA	ALA	ALA	ALA	ALA
ALA	A373	ALA	ALA	ALA	ALA	ALA
ALA	A374	ALA	ALA	ALA	ALA	ALA
ALA	R675	ALA	ALA	ALA	ALA	ALA
ALA	A676	ALA	ALA	ALA	ALA	ALA
ALA	D677	ALA	ALA	ALA	ALA	ALA
ALA	T678	ALA	ALA	ALA	ALA	ALA
ALA	R679	ALA	ALA	ALA	ALA	ALA



## 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	51758	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	71.85	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	2.574	Depositor
Minimum map value	-1.473	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.068	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	226.84799, 226.84799, 226.84799	wwPDB
Map dimensions	272, 272, 272	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.834, 0.834, 0.834	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.42	0/4131	0.62	0/5583
1	B	0.43	0/4131	0.62	0/5583
1	C	0.43	0/4131	0.62	0/5583
1	D	0.43	0/4131	0.62	0/5583
All	All	0.43	0/16524	0.62	0/22332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4033	0	4046	146	0
1	B	4033	0	4046	141	0
1	C	4033	0	4046	146	0
1	D	4033	0	4046	151	0
All	All	16132	0	16184	536	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 536 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:537:GLN:NE2	1:B:563:LYS:HB2	1.58	1.16
1:C:537:GLN:NE2	1:D:563:LYS:HB2	1.59	1.16
1:B:537:GLN:NE2	1:C:563:LYS:HB2	1.60	1.15
1:A:563:LYS:HB2	1:D:537:GLN:NE2	1.60	1.14
1:A:367:GLU:HG3	1:A:370:LYS:HB2	1.45	0.97

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	488/892 (55%)	450 (92%)	38 (8%)	0	100 100
1	B	488/892 (55%)	449 (92%)	39 (8%)	0	100 100
1	C	488/892 (55%)	450 (92%)	38 (8%)	0	100 100
1	D	488/892 (55%)	451 (92%)	37 (8%)	0	100 100
All	All	1952/3568 (55%)	1800 (92%)	152 (8%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	444/733 (61%)	438 (99%)	6 (1%)	62 78
1	B	444/733 (61%)	438 (99%)	6 (1%)	62 78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	444/733 (61%)	438 (99%)	6 (1%)	62	78
1	D	444/733 (61%)	438 (99%)	6 (1%)	62	78
All	All	1776/2932 (61%)	1752 (99%)	24 (1%)	62	78

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	339	LEU
1	C	582	ILE
1	C	375	LEU
1	D	324	ASN
1	B	324	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	553	HIS
1	D	459	ASN
1	C	324	ASN
1	D	527	GLN
1	D	277	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-12512. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [\(i\)](#)


### 6.2.1 Primary map



X Index: 136




Y Index: 136




Z Index: 136


### 6.2.2 Raw map



X Index: 136



Y Index: 136




Z Index: 136


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [\(i\)](#)


### 6.3.1 Primary map



X Index: 118




Y Index: 154




Z Index: 103


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [\(i\)](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [\(i\)](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_12512\_msk\_1.map [\(i\)](#)



X



Y




Z

## 7 Map analysis (i)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate (i)



The volume at the recommended contour level is  $68 nm^3$ ; this corresponds to an approximate mass of 61 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [\(i\)](#)




\*Reported resolution corresponds to spatial frequency of  $0.303 \text{ \AA}^{-1}$

## 8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [\(i\)](#)



\*Reported resolution corresponds to spatial frequency of 0.303 Å<sup>-1</sup>

## 8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	3.26	3.48	3.28
Unmasked-calculated*	3.60	4.19	3.65

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [\(i\)](#)


This section contains information regarding the fit between EMDB map EMD-12512 and PDB model 7NP3. Per-residue inclusion information can be found in section 3 on page 5.

### 9.1 Map-model overlay [\(i\)](#)




The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

## 9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 64% of all backbone atoms, 55% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5530	0.4190
A	0.5540	0.4180
B	0.5520	0.4170
C	0.5530	0.4200
D	0.5530	0.4210

