

# Full wwPDB X-ray Structure Validation Report (i)

#### Mar 24, 2025 – 12:23 PM EDT

PDB ID		
Title	:	Crystal structure of phosphoglycerate mutase from Trichomonas vaginalis
Authors	:	Seattle Structural Genomics Center for Infectious Disease; Seattle Structural
		Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2025-03-20
Resolution	:	1.95  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
5		2022.3.0, CSD as543be (2022)
Xtriage (Phenix)		
EDS	:	3.0
buster-report	:	1.1.7 (2018)
-		20231227.v01 (using entries in the PDB archive December 27th 2023)
		9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	258	% • 82%	5%	13%
1	В	258	83%	•	12%
1	С	258	5%	6%	16%
1	D	258	79%	6%	16%



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	224	Total	С	Ν	0	$\mathbf{S}$	0	2	0
	А	224	1807	1155	307	330	15	0		
1	В	226	Total	С	Ν	0	S	0	1	0
	D	220	1821	1163	311	333	14			
1	C	218	Total	С	Ν	0	S	0	0	0
	U	210	1731	1106	296	316	13	0		0
1	1 D	218	Total	С	Ν	0	S	0	0	0
		210	1710	1090	291	316	13	0	0	

• Molecule 1 is a protein called Phosphoglycerate mutase.

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
А	-7	MET	-	initiating methionine	UNP A2DUN8
А	-6	ALA	-	expression tag	UNP A2DUN8
А	-5	HIS	-	expression tag	UNP A2DUN8
А	-4	HIS	-	expression tag	UNP A2DUN8
А	-3	HIS	-	expression tag	UNP A2DUN8
А	-2	HIS	-	expression tag	UNP A2DUN8
А	-1	HIS	-	expression tag	UNP A2DUN8
А	0	HIS	-	expression tag	UNP A2DUN8
В	-7	MET	-	initiating methionine	UNP A2DUN8
В	-6	ALA	-	expression tag	UNP A2DUN8
В	-5	HIS	-	expression tag	UNP A2DUN8
В	-4	HIS	-	expression tag	UNP A2DUN8
В	-3	HIS	-	expression tag	UNP A2DUN8
В	-2	HIS	-	expression tag	UNP A2DUN8
В	-1	HIS	-	expression tag	UNP A2DUN8
В	0	HIS	-	expression tag	UNP A2DUN8
С	-7	MET	-	initiating methionine	UNP A2DUN8
С	-6	ALA	-	expression tag	UNP A2DUN8
С	-5	HIS	-	expression tag	UNP A2DUN8
С	-4	HIS	-	expression tag	UNP A2DUN8
С	-3	HIS	-	expression tag	UNP A2DUN8



9NV5	
------	--

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP A2DUN8
С	-1	HIS	-	expression tag	UNP A2DUN8
С	0	HIS	-	expression tag	UNP A2DUN8
D	-7	MET	-	initiating methionine	UNP A2DUN8
D	-6	ALA	-	expression tag	UNP A2DUN8
D	-5	HIS	-	expression tag	UNP A2DUN8
D	-4	HIS	-	expression tag	UNP A2DUN8
D	-3	HIS	-	expression tag	UNP A2DUN8
D	-2	HIS	-	expression tag	UNP A2DUN8
D	-1	HIS	-	expression tag	UNP A2DUN8
D	0	HIS	-	expression tag	UNP A2DUN8

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Δ	1	Total O S	0	0
2	11	T	$5 \ 4 \ 1$	0	0
2	Δ	1	Total O S	0	0
2	Π	1	$5 \ 4 \ 1$	0	
2	В	1	Total O S	0	0
	D	L	$5 \ 4 \ 1$	0	0
2	В	1	Total O S	0	0
	D	1	$5 \ 4 \ 1$	0	0
2	D	1	Total O S	0	0
	D	1	$5 \ 4 \ 1$	0	U



Continued from	previous	page
----------------	----------	------

Mo	bl	Chain	Residues	Atoms	ZeroOcc	AltConf
2		С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2		D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0


• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 6  3  3 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 6  3  3 \end{array}$	0	0


• Molecule 4 is 3,6,9,12,15,18-HEXAOXAICOSANE-1,20-DIOL (three-letter code: P33) (formula:  $C_{14}H_{30}O_8$ ).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	Δ	1	Total	С	0	0	0
4	4 A	1	22	14	8	0	0

• Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total         C         O           11         7         4	0	0
5	А	1	Total         C         O           13         8         5	0	0
5	В	1	Total         C         O           13         8         5	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total         C         O           13         8         5	0	0
5	D	1	Total         C         O           13         8         5	0	0
5	D	1	Total         C         O           10         6         4	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	172	Total O 172 172	0	0
6	В	155	Total O 155 155	0	0
6	С	100	Total O 100 100	0	0
6	D	113	Total O 113 113	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Phosphoglycerate mutase







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	73.78Å 92.70Å 94.00Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $94.09^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	19.96 - 1.95	Depositor
Resolution (A)	19.96 - 1.95	EDS
% Data completeness	99.8 (19.96-1.95)	Depositor
(in resolution range)	99.7 (19.96 - 1.95)	EDS
R <sub>merge</sub>	0.14	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.89 (at 1.94 \text{\AA})$	Xtriage
Refinement program	PHENIX (dev_5634: ???)	Depositor
D D.	0.156 , $0.186$	Depositor
$R, R_{free}$	0.166 , $0.195$	DCC
$R_{free}$ test set	4516 reflections $(4.91%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	26.0	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38, 56.2	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7751	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, SO4, P33, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
	Unain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.83	0/1854	0.77	0/2507
1	В	0.83	0/1865	0.80	0/2520
1	С	0.70	0/1772	0.72	0/2400
1	D	0.71	0/1748	0.71	0/2365
All	All	0.77	0/7239	0.75	0/9792

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	68	ARG	Sidechain

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1807	0	1789	7	0
1	В	1821	0	1809	6	0
1	С	1731	0	1684	7	0
1	D	1710	0	1649	7	0
2	А	10	0	0	0	0
2	В	15	0	0	2	0
2	С	5	0	0	0	0
2	D	5	0	0	0	0
3	А	6	0	8	1	0
3	В	6	0	8	1	0
4	А	22	0	30	1	0
5	А	24	0	31	0	0
5	В	26	0	36	1	0
5	D	23	0	31	0	0
6	А	172	0	0	0	0
6	В	155	0	0	1	0
6	С	100	0	0	0	0
6	D	113	0	0	0	0
All	All	7751	0	7075	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:15:LEU:HD11	1:C:47:ALA:HB1	1.76	0.66
1:D:120:ILE:HB	1:D:127:LEU:HD12	1.86	0.57
1:D:220:LEU:HD23	1:D:235:ILE:HG12	1.89	0.55
1:A:15[A]:LEU:HD23	1:A:219:ALA:HB2	1.88	0.54
1:B:116:GLU:HA	1:B:116:GLU:OE1	2.08	0.53
1:B:198:LYS:HD3	3:B:304:GOL:H31	1.92	0.51
1:B:15:LEU:HD23	1:B:219:ALA:HB2	1.91	0.51
1:B:149:ASP:O	1:B:152:VAL:HG12	2.11	0.50
1:A:107:GLU:H	1:A:107:GLU:CD	2.15	0.50
1:A:107:GLU:O	1:A:111:THR:HG23	2.13	0.49
1:D:39:LYS:HE2	1:D:43:ASP:OD1	2.13	0.48
1:C:198:LYS:HA	1:C:203:MET:HG3	1.97	0.47
2:B:302:SO4:O1	5:B:306:PG4:H12	2.14	0.46
1:B:86:GLN:OE1	2:B:303:SO4:O2	2.32	0.46
1:C:105:ASN:OD1	1:C:108:GLN:HG3	2.16	0.45
1:C:225:ASP:OD1	1:C:229:ASN:HB2	2.17	0.45



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:149:ASP:O	1:C:152:VAL:HG12	2.17	0.45
1:D:149:ASP:O	1:D:152:VAL:HG12	2.17	0.45
1:A:15[A]:LEU:HD11	1:A:51:LEU:HD11	2.00	0.43
1:D:158:SER:OG	1:D:161:MET:HG3	2.18	0.43
4:A:304:P33:H52	6:B:530:HOH:O	2.19	0.42
1:A:86:GLN:HG2	3:A:303:GOL:O3	2.20	0.42
1:A:169:TYR:CE1	1:A:173:GLN:HG3	2.56	0.41
1:C:162:MET:C	1:C:162:MET:SD	2.99	0.41
1:B:151:ARG:NH1	1:D:199:TYR:O	2.55	0.40
1:D:154:PRO:HB2	1:D:157:GLU:HG3	2.02	0.40
1:C:22:SER:HA	1:C:25:ASN:OD1	2.22	0.40
1:A:149:ASP:O	1:A:152:VAL:HG12	2.22	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	222/258~(86%)	217~(98%)	4 (2%)	1 (0%)	25	16
1	В	223/258~(86%)	219~(98%)	4 (2%)	0	100	100
1	С	214/258~(83%)	210 (98%)	4 (2%)	0	100	100
1	D	212/258~(82%)	207~(98%)	5(2%)	0	100	100
All	All	871/1032 (84%)	853~(98%)	17~(2%)	1 (0%)	48	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	95	GLU



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	200/234~(86%)	199~(100%)	1 (0%)	86	86
1	В	202/234~(86%)	200~(99%)	2(1%)	73	72
1	С	187/234 (80%)	183 (98%)	4 (2%)	48	43
1	D	183/234 (78%)	181 (99%)	2(1%)	70	68
All	All	772/936~(82%)	763~(99%)	9 (1%)	67	65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	65	TYR
1	В	4	LYS
1	В	65	TYR
1	С	15	LEU
1	С	16	ARG
1	С	29	SER
1	С	65	TYR
1	D	65	TYR
1	D	81	ASP

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	В	86	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry (i)

#### 16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
10101	туре	Ullaili	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	SO4	А	302	-	4,4,4	0.59	0	$6,\!6,\!6$	1.00	0
5	PG4	D	302	-	12,12,12	0.34	0	11,11,11	0.38	0
3	GOL	А	303	-	$5,\!5,\!5$	0.77	0	$5,\!5,\!5$	1.21	0
2	SO4	D	301	-	4,4,4	0.96	0	$6,\!6,\!6$	0.53	0
2	SO4	В	303	-	4,4,4	0.92	0	$6,\!6,\!6$	0.44	0
5	PG4	D	303	-	9,9,12	0.35	0	8,8,11	0.20	0
2	SO4	С	301	-	4,4,4	0.77	0	$6,\!6,\!6$	0.44	0
2	SO4	В	302	-	4,4,4	0.90	0	$6,\!6,\!6$	0.78	0
2	SO4	В	301	-	4,4,4	0.77	0	$6,\!6,\!6$	0.31	0
5	PG4	А	305	-	10,10,12	0.42	0	$9,\!9,\!11$	0.39	0
5	PG4	В	305	-	12,12,12	0.29	0	$11,\!11,\!11$	0.45	0
5	PG4	В	306	-	12,12,12	0.37	0	11,11,11	0.51	0
4	P33	А	304	-	21,21,21	0.38	0	20,20,20	0.45	0
3	GOL	В	304	-	$5,\!5,\!5$	0.80	0	$5,\!5,\!5$	0.68	0
2	SO4	А	301	-	4,4,4	0.80	0	$6,\!6,\!6$	0.43	0
5	PG4	А	306	-	12,12,12	0.37	0	11,11,11	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PG4	D	302	-	-	6/10/10/10	-
3	GOL	А	303	-	-	0/4/4/4	-
5	PG4	D	303	-	-	3/7/7/10	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PG4	А	305	-	-	3/8/8/10	-
5	PG4	В	305	-	-	6/10/10/10	-
5	PG4	В	306	-	-	7/10/10/10	-
4	P33	А	304	-	-	7/19/19/19	-
3	GOL	В	304	-	-	4/4/4/4	-
5	PG4	А	306	-	-	4/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (40) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	304	GOL	C1-C2-C3-O3
5	D	302	PG4	O2-C3-C4-O3
4	А	304	P33	O16-C17-C18-O19
5	В	305	PG4	O2-C3-C4-O3
5	D	302	PG4	O3-C5-C6-O4
5	В	306	PG4	O2-C3-C4-O3
5	А	306	PG4	O1-C1-C2-O2
5	А	306	PG4	O4-C7-C8-O5
5	В	306	PG4	O4-C7-C8-O5
5	D	303	PG4	O3-C5-C6-O4
3	В	304	GOL	O1-C1-C2-C3
4	А	304	P33	O1-C2-C3-O4
4	А	304	P33	O13-C14-C15-O16
5	В	305	PG4	O4-C7-C8-O5
4	А	304	P33	O7-C8-C9-O10
5	А	305	PG4	O4-C7-C8-O5
5	D	303	PG4	O2-C3-C4-O3
3	В	304	GOL	O2-C2-C3-O3
5	А	305	PG4	C4-C3-O2-C2
5	В	305	PG4	C3-C4-O3-C5
5	D	302	PG4	C4-C3-O2-C2
5	А	305	PG4	C8-C7-O4-C6
5	D	303	PG4	C1-C2-O2-C3
5	D	302	PG4	C6-C5-O3-C4
5	В	305	PG4	C1-C2-O2-C3
3	В	304	GOL	O1-C1-C2-O2



Mol	Chain	Res	Type	Atoms
5	D	302	PG4	O1-C1-C2-O2
4	А	304	P33	C8-C9-O10-C11
5	В	306	PG4	C6-C5-O3-C4
5	D	302	PG4	C1-C2-O2-C3
5	В	306	PG4	C4-C3-O2-C2
5	В	306	PG4	O3-C5-C6-O4
5	В	305	PG4	O3-C5-C6-O4
4	А	304	P33	C2-C3-O4-C5
4	А	304	P33	C14-C15-O16-C17
5	В	306	PG4	C3-C4-O3-C5
5	В	305	PG4	C6-C5-O3-C4
5	В	306	PG4	C8-C7-O4-C6
5	А	306	PG4	C6-C5-O3-C4
5	А	306	PG4	C5-C6-O4-C7

Continued from previous page...


There are no ring outliers.

6 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	303	GOL	1	0
2	В	303	SO4	1	0
2	В	302	SO4	1	0
5	В	306	PG4	1	0
4	А	304	P33	1	0
3	В	304	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSI	RZ>2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	224/258~(86%)	-0.35	3 (1%)	74 79	17, 27, 51, 89	2 (0%)
1	В	226/258~(87%)	-0.32	8 (3%)	47 54	16, 26, 54, 96	1 (0%)
1	С	218/258~(84%)	0.08	12 (5%)	32 38	22, 36, 78, 92	0
1	D	218/258~(84%)	0.04	14 (6%)	27 32	22, 35, 84, 107	0
All	All	886/1032 (85%)	-0.14	37 (4%)	41 48	16, 31, 74, 107	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	121	TRP	5.8
1	D	216	ASN	5.3
1	А	203	MET	4.2
1	С	120	ILE	4.0
1	D	110	CYS	3.9
1	С	24	LEU	3.7
1	D	118	LEU	3.6
1	С	204	THR	3.5
1	В	216	ASN	3.3
1	D	120	ILE	3.2
1	С	238	GLU	3.1
1	D	109	ILE	2.8
1	С	123	LYS	2.8
1	С	23	ASN	2.6
1	С	15	LEU	2.6
1	В	3	ARG	2.6
1	С	110	CYS	2.5
1	В	238	GLU	2.5
1	В	214	LEU	2.5
1	В	203	MET	2.4
1	D	124	ASP	2.4



Mol	Chain	Res Type		RSRZ	
1	D	239	GLU	2.4	
1	D	119	ASN	2.3	
1	D	217	GLY	2.3	
1	D	28	TYR	2.3	
1	В	215	SER	2.2	
1	А	3	ARG	2.2	
1	С	21	LEU	2.2	
1	С	227	LYS	2.2	
1	D	116	GLU	2.1	
1	В	4	LYS	2.1	
1	А	107	GLU	2.1	
1	С	203	MET	2.1	
1	С	113	LEU	2.1	
1	D	111	THR	2.1	
1	В	121	TRP	2.0	
1	D	113	LEU	2.0	

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	SO4	С	301	5/5	0.68	0.14	49,63,69,84	0
5	PG4	А	306	13/13	0.72	0.17	47,61,71,77	0
4	P33	А	304	22/22	0.73	0.15	38,47,56,59	0
5	PG4	D	302	13/13	0.79	0.14	43,51,59,59	0
5	PG4	В	306	13/13	0.82	0.14	47,56,64,66	0
3	GOL	А	303	6/6	0.82	0.14	32,44,48,55	0
5	PG4	D	303	10/13	0.84	0.14	48,58,62,65	0




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
3	GOL	В	304	6/6	0.86	0.15	$39,\!43,\!51,\!53$	0
2	SO4	В	301	5/5	0.87	0.10	68,69,74,75	0
5	PG4	А	305	11/13	0.87	0.12	$32,\!39,\!49,\!55$	0
2	SO4	А	302	5/5	0.89	0.14	36,43,49,58	0
5	PG4	В	305	13/13	0.90	0.10	40,46,54,61	0
2	SO4	D	301	5/5	0.90	0.15	46,49,61,62	0
2	SO4	В	303	5/5	0.90	0.15	$37,\!51,\!55,\!62$	0
2	SO4	А	301	5/5	0.90	0.09	52,62,67,67	0
2	SO4	В	302	5/5	0.94	0.16	43,43,46,48	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




































## 6.5 Other polymers (i)

There are no such residues in this entry.

