



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:24 PM UTC

PDB ID : 6O6V / pdb_00006o6v
Title : Crystal structure of Csm6 in complex with cA4 by soaking cA4 into Csm6
Authors : Jia, N.; Patel, D.J.
Deposited on : 2019-03-07
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)
RNA backbone	3983	1027 (2.66-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	
1	B	440	
2	C	4	
2	D	4	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Csm6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	Total	C	N	O	S	0	0	0
			3450	2220	586	636	8			
1	B	432	Total	C	N	O	S	0	0	0
			3440	2214	583	635	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP B6YWC3
A	0	GLY	-	expression tag	UNP B6YWC3
A	433	HIS	-	expression tag	UNP B6YWC3
A	434	HIS	-	expression tag	UNP B6YWC3
A	435	HIS	-	expression tag	UNP B6YWC3
A	436	HIS	-	expression tag	UNP B6YWC3
A	437	HIS	-	expression tag	UNP B6YWC3
A	438	HIS	-	expression tag	UNP B6YWC3
B	-1	MET	-	initiating methionine	UNP B6YWC3
B	0	GLY	-	expression tag	UNP B6YWC3
B	433	HIS	-	expression tag	UNP B6YWC3
B	434	HIS	-	expression tag	UNP B6YWC3
B	435	HIS	-	expression tag	UNP B6YWC3
B	436	HIS	-	expression tag	UNP B6YWC3
B	437	HIS	-	expression tag	UNP B6YWC3
B	438	HIS	-	expression tag	UNP B6YWC3

- Molecule 2 is a RNA chain called Cyclic RNA cA4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	4	Total	C	N	O	P	0	0	0
			88	40	20	24	4			
2	D	4	Total	C	N	O	P	0	0	0
			88	40	20	24	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total 13	O 13	0	0
3	B	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Csm6



- Molecule 1: Csm6



- Molecule 2: Cyclic RNA cA4



- Molecule 2: Cyclic RNA cA4



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	103.99Å 164.41Å 111.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.53 – 2.35 48.53 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.53-2.35) 99.5 (48.53-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.37Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.227 , 0.271 0.227 , 0.267	Depositor DCC
R_{free} test set	1993 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	58.2	Xtrriage
Anisotropy	0.384	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 26.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7088	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	0/3526	1.47	0/4772
1	B	1.00	0/3515	1.45	0/4757
2	C	1.73	0/99	1.75	2/152 (1.3%)
2	D	1.95	0/99	1.83	3/152 (2.0%)
All	All	1.03	0/7239	1.47	5/9833 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	A	P-O3'-C3'	6.58	130.08	120.20
2	C	1	A	C1'-O4'-C4'	-6.10	103.60	109.70
2	C	1	A	O4'-C1'-N9	5.21	116.02	108.20
2	D	3	A	N9-C1'-C2'	5.04	119.56	112.00
2	D	2	A	O3'-P-O5'	-5.02	96.47	104.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3450	0	3499	26	0
1	B	3440	0	3492	19	0
2	C	88	0	44	0	0
2	D	88	0	44	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	13	0	0	0	0
3	B	9	0	0	0	0
All	All	7088	0	7079	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ASP:OD1	1:A:131:THR:HG23	1.91	0.70
1:B:52:ILE:O	1:B:69:ARG:NH1	2.32	0.62
1:A:361:ARG:NH2	1:A:367:ASP:O	2.36	0.59
1:A:14:TRP:CZ2	1:A:132:HIS:HB3	2.37	0.58
1:A:340:LYS:HG2	2:D:1:A:OP1	2.04	0.56
1:B:50:ASP:HB3	1:B:87:ILE:HG22	1.88	0.56
1:A:145:LEU:O	1:A:149:LEU:HG	2.06	0.56
2:D:1:A:H2'	2:D:4:A:O2'	2.05	0.55
1:A:333:ARG:HD3	1:B:333:ARG:NE	2.22	0.55
1:B:29:PRO:HD2	1:B:77:ILE:HG21	1.90	0.53
1:B:172:ILE:HG12	1:B:172:ILE:O	2.08	0.52
1:B:207:MET:HE3	1:B:212:PHE:HA	1.91	0.52
1:A:134:LEU:HD23	1:A:134:LEU:N	2.25	0.52
1:A:6:VAL:HG22	1:A:129:ASP:HB3	1.90	0.52
1:B:347:LYS:HD2	2:D:2:A:H61	1.76	0.50
1:A:9:GLY:HA2	1:A:134:LEU:CD2	2.43	0.49
1:A:274:VAL:O	1:A:274:VAL:HG13	2.13	0.49
1:B:48:LEU:HD13	1:B:70:VAL:HG21	1.96	0.48
1:A:129:ASP:OD1	1:A:131:THR:CG2	2.62	0.48
1:A:326:PHE:C	1:A:327:ARG:HG3	2.39	0.48
1:A:309:VAL:HG11	1:A:317:LYS:HE2	1.97	0.47
1:A:9:GLY:HA2	1:A:134:LEU:HD21	1.96	0.47
1:A:17:THR:OG1	1:A:18:LYS:N	2.47	0.46
1:A:6:VAL:CG1	1:A:8:TRP:CD1	2.98	0.46
1:B:242:GLY:HA2	1:B:326:PHE:CE1	2.51	0.46
1:A:159:THR:HG22	1:A:185:MET:SD	2.57	0.45
1:A:1:MET:HG2	1:A:119:PRO:HG2	1.98	0.45
1:B:333:ARG:NH1	1:B:337:GLU:OE2	2.50	0.44
1:A:162:ALA:HB3	1:A:182:GLU:HB3	2.00	0.44
1:B:207:MET:HE3	1:B:212:PHE:CA	2.47	0.44
1:B:370:ASN:HD22	1:B:370:ASN:HA	1.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:LEU:N	1:B:33:PRO:CD	2.80	0.44
1:B:129:ASP:OD2	1:B:131:THR:OG1	2.33	0.44
1:A:9:GLY:CA	1:A:134:LEU:CD2	2.96	0.43
1:B:165:SER:HA	1:B:177:ASN:O	2.18	0.43
1:A:31:THR:HG21	1:A:131:THR:HG21	2.00	0.42
1:A:135:ASN:CG	1:B:164:ASN:HD21	2.27	0.42
1:A:242:GLY:HA2	1:A:326:PHE:CE1	2.54	0.42
1:B:216:LYS:HG3	1:B:222:LEU:HD12	2.00	0.42
1:A:11:PHE:HA	1:A:14:TRP:CE2	2.54	0.42
1:A:56:PHE:O	1:A:274:VAL:HG22	2.20	0.42
1:B:325:LEU:HD12	1:B:325:LEU:HA	1.91	0.41
1:A:344:LYS:O	1:A:348:TYR:HD1	2.03	0.41
1:B:11:PHE:HA	1:B:14:TRP:CE2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/440 (98%)	413 (96%)	18 (4%)	0	100	100
1	B	430/440 (98%)	416 (97%)	14 (3%)	0	100	100
All	All	861/880 (98%)	829 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/378 (98%)	358 (96%)	14 (4%)	29	39
1	B	371/378 (98%)	361 (97%)	10 (3%)	39	52
All	All	743/756 (98%)	719 (97%)	24 (3%)	34	46

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	13	ARG
1	A	17	THR
1	A	26	SER
1	A	60	GLU
1	A	61	THR
1	A	83	VAL
1	A	131	THR
1	A	134	LEU
1	A	183	THR
1	A	239	LEU
1	A	356	ARG
1	A	371	ARG
1	A	412	LYS
1	B	30	SER
1	B	61	THR
1	B	132	HIS
1	B	169	VAL
1	B	172	ILE
1	B	173	THR
1	B	177	ASN
1	B	333	ARG
1	B	344	LYS
1	B	402	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	370	ASN
1	B	164	ASN
1	B	370	ASN
1	B	410	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	3/4 (75%)	3 (100%)	0
2	D	3/4 (75%)	3 (100%)	2 (66%)
All	All	6/8 (75%)	6 (100%)	2 (33%)

All (6) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	2	A
2	C	3	A
2	C	4	A
2	D	2	A
2	D	3	A
2	D	4	A

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	2	A
2	D	3	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/440 (98%)	0.59	22 (5%) 33 39	42, 63, 95, 116	0
1	B	432/440 (98%)	0.56	29 (6%) 24 27	44, 67, 101, 119	0
2	C	4/4 (100%)	0.96	0 100 100	100, 108, 117, 133	0
2	D	4/4 (100%)	1.41	0 100 100	111, 116, 164, 175	0
All	All	873/888 (98%)	0.58	51 (5%) 29 33	42, 65, 101, 175	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	48	LEU	5.0
1	A	183	THR	4.2
1	B	114	LEU	3.5
1	B	212	PHE	3.2
1	A	91	ILE	3.2
1	A	172	ILE	3.1
1	B	124	LEU	3.1
1	B	167	PRO	3.1
1	B	1	MET	3.1
1	B	428	ALA	3.0
1	A	368	ALA	3.0
1	A	169	VAL	2.9
1	A	432	SER	2.9
1	A	94	PHE	2.9
1	B	169	VAL	2.9
1	A	11	PHE	2.9
1	B	79	ALA	2.8
1	B	137	VAL	2.8
1	B	432	SER	2.8
1	A	170	PRO	2.7
1	A	225	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	136	TYR	2.6
1	B	370	ASN	2.6
1	A	134	LEU	2.6
1	B	215	LEU	2.5
1	B	170	PRO	2.5
1	B	273	ALA	2.5
1	A	167	PRO	2.4
1	A	52	ILE	2.4
1	B	59	VAL	2.4
1	A	92	GLY	2.3
1	B	180	THR	2.2
1	A	1	MET	2.2
1	A	168	PHE	2.2
1	A	307	LEU	2.2
1	B	52	ILE	2.2
1	B	400	ARG	2.1
1	B	307	LEU	2.1
1	B	55	ASP	2.1
1	B	70	VAL	2.1
1	A	136	TYR	2.1
1	B	107	TYR	2.1
1	A	287	PHE	2.1
1	B	11	PHE	2.1
1	B	213	VAL	2.1
1	A	341	ILE	2.0
1	B	171	LYS	2.0
1	A	369	ALA	2.0
1	B	144	ALA	2.0
1	A	312	PRO	2.0
1	B	333	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.