



wwPDB X-ray Structure Validation Summary Report i

Nov 11, 2025 – 06:18 PM EST

PDB ID : 9OES / pdb_00009oes
Title : HalA with lysine, succinate, chloride, and vanadium(IV)-oxo at pH 4.2
Authors : Kissman, E.N.; Chang, M.C.Y.
Deposited on : 2025-04-29
Resolution : 1.93 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4.5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1306 (1.94-1.94)
Clashscore	180529	1400 (1.94-1.94)
Ramachandran outliers	177936	1387 (1.94-1.94)
Sidechain outliers	177891	1387 (1.94-1.94)
RSRZ outliers	164620	1306 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	F	258	2%	92%	7% •
1	G	258	%	91%	7% •
1	H	258	2%	92%	6% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	VVO	A	304	-	-	X	-
7	FLC	B	504	-	X	X	-
7	FLC	G	303	-	-	X	-

2 Entry composition (i)


There are 9 unique types of molecules in this entry. The entry contains 17579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine halogenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			1993	1254	363	370	6			
1	B	254	Total	C	N	O	S	0	1	0
			1996	1256	363	371	6			
1	C	254	Total	C	N	O	S	0	1	0
			1975	1241	364	364	6			
1	D	255	Total	C	N	O	S	0	1	0
			2008	1262	366	374	6			
1	E	255	Total	C	N	O	S	0	1	0
			2011	1262	369	374	6			
1	F	255	Total	C	N	O	S	0	0	0
			1972	1242	356	368	6			
1	G	253	Total	C	N	O	S	0	3	0
			1991	1250	365	370	6			
1	H	253	Total	C	N	O	S	0	2	0
			1978	1243	361	368	6			

- Molecule 2 is LYSINE (CCD ID: LYS) (formula: C₆H₁₅N₂O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	6	2	2		
2	B	1	Total	C	N	O	0	0
			10	6	2	2		
2	C	1	Total	C	N	O	0	0
			10	6	2	2		
2	D	1	Total	C	N	O	0	0
			10	6	2	2		
2	E	1	Total	C	N	O	0	0
			10	6	2	2		
2	F	1	Total	C	N	O	0	0
			10	6	2	2		
2	G	1	Total	C	N	O	0	0
			10	6	2	2		
2	H	1	Total	C	N	O	0	0
			10	6	2	2		

- Molecule 3 is SUCCINIC ACID (CCD ID: SIN) (formula: C₄H₆O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 4 4	0	0
3	B	1	Total C O 8 4 4	0	0
3	C	1	Total C O 8 4 4	0	0
3	D	1	Total C O 8 4 4	0	0
3	E	1	Total C O 8 4 4	0	0
3	F	1	Total C O 8 4 4	0	0
3	G	1	Total C O 8 4 4	0	0
3	H	1	Total C O 8 4 4	0	0

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 10 6 4	0	0
4	B	1	Total C O 10 6 4	0	0
4	F	1	Total C O 10 6 4	0	0
4	F	1	Total C O 10 6 4	0	0
4	H	1	Total C O 10 6 4	0	0

- Molecule 5 is oxovanadium(2+) (CCD ID: VVO) (formula: OV) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O V 2 1 1	0	0

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	B	1	Total Cl 1 1	0	0
6	C	1	Total Cl 1 1	0	0
6	D	1	Total Cl 1 1	0	0
6	E	1	Total Cl 1 1	0	0
6	F	1	Total Cl 1 1	0	0
6	G	1	Total Cl 1 1	0	0
6	H	1	Total Cl 1 1	0	0

- Molecule 7 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 13 6 7	0	0
7	G	1	Total C O 13 6 7	0	0

- Molecule 8 is VANADIUM ION (CCD ID: V) (formula: V) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total V 1 1	0	0
8	C	1	Total V 1 1	0	0
8	D	1	Total V 1 1	0	0
8	E	1	Total V 1 1	0	0
8	F	1	Total V 1 1	0	0
8	G	1	Total V 1 1	0	0
8	H	1	Total V 1 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	205	Total O 205 205	0	0
9	B	198	Total O 198 198	0	0
9	C	188	Total O 188 188	0	0
9	D	193	Total O 193 193	0	0
9	E	185	Total O 185 185	0	0
9	F	166	Total O 166 166	0	0
9	G	160	Total O 160 160	0	0
9	H	123	Total O 123 123	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lysine halogenase




- Molecule 1: Lysine halogenase




- Molecule 1: Lysine halogenase



- Molecule 1: Lysine halogenase



- Molecule 1: Lysine halogenase




- Molecule 1: Lysine halogenase



- Molecule 1: Lysine halogenase



- Molecule 1: Lysine halogenase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	146.95Å 146.95Å 287.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	95.35 – 1.93 95.35 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.4 (95.35-1.93) 99.3 (95.35-1.93)	Depositor EDS
R_{merge}	0.36	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.23 (at 1.92Å)	Xtriage
Refinement program	PHENIX 1.21rc1_5156	Depositor
R , R_{free}	0.174, 0.216 0.175, 0.216	Depositor DCC
R_{free} test set	8735 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	29.5	Xtriage
Anisotropy	0.582	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33, 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.088 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	17579	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: V, CL, PGE, SIN, VVO, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/2038	0.49	0/2774
1	B	0.31	0/2044	0.50	0/2784
1	C	0.31	0/2020	0.50	0/2755
1	D	0.31	0/2053	0.47	0/2797
1	E	0.30	0/2056	0.46	0/2800
1	F	0.31	0/2016	0.47	0/2749
1	G	0.27	0/2037	0.46	0/2776
1	H	0.26	0/2024	0.46	0/2763
All	All	0.30	0/16288	0.48	0/22198

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1993	0	1906	12	0
1	B	1996	0	1908	7	0
1	C	1975	0	1863	7	0
1	D	2008	0	1917	14	0
1	E	2011	0	1923	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1972	0	1869	8	0
1	G	1991	0	1874	13	0
1	H	1978	0	1853	8	0
2	A	10	0	12	0	0
2	B	10	0	12	0	0
2	C	10	0	12	0	0
2	D	10	0	12	0	0
2	E	10	0	12	0	0
2	F	10	0	12	0	0
2	G	10	0	12	0	0
2	H	10	0	12	0	0
3	A	8	0	4	0	0
3	B	8	0	4	0	0
3	C	8	0	4	1	0
3	D	8	0	4	0	0
3	E	8	0	4	0	0
3	F	8	0	4	0	0
3	G	8	0	4	1	0
3	H	8	0	4	0	0
4	A	10	0	14	0	0
4	B	10	0	14	0	0
4	F	20	0	28	2	0
4	H	10	0	14	1	0
5	A	2	0	0	5	0
6	A	1	0	0	1	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
6	G	1	0	0	0	0
6	H	1	0	0	0	0
7	B	13	0	5	12	0
7	G	13	0	5	4	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
8	E	1	0	0	0	0
8	F	1	0	0	0	0
8	G	1	0	0	0	0
8	H	1	0	0	0	0
9	A	205	0	0	2	8

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	B	198	0	0	11	0
9	C	188	0	0	2	0
9	D	193	0	0	2	0
9	E	185	0	0	5	0
9	F	166	0	0	2	6
9	G	160	0	0	10	2
9	H	123	0	0	2	0
All	All	17579	0	15321	97	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:HIS:NE2	5:A:304:VVO:O1	1.61	1.28
7:B:504:FLC:OB2	9:B:601:HOH:O	1.82	0.96
7:B:504:FLC:HG2	1:D:210:ARG:HH12	1.34	0.93
1:E:133:GLU:OE2	9:E:401:HOH:O	1.89	0.89
7:G:303:FLC:OA2	9:G:401:HOH:O	1.89	0.89

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:589:HOH:O	9:F:542:HOH:O[5_555]	1.95	0.25
9:A:570:HOH:O	9:F:515:HOH:O[5_555]	1.96	0.24
9:A:512:HOH:O	9:G:549:HOH:O[5_555]	2.06	0.14
9:A:604:HOH:O	9:F:564:HOH:O[5_555]	2.07	0.13
9:A:567:HOH:O	9:F:550:HOH:O[5_555]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/258 (98%)	242 (96%)	9 (4%)	1 (0%)	30 22
1	B	253/258 (98%)	246 (97%)	6 (2%)	1 (0%)	30 22
1	C	253/258 (98%)	249 (98%)	4 (2%)	0	100 100
1	D	254/258 (98%)	247 (97%)	7 (3%)	0	100 100
1	E	254/258 (98%)	248 (98%)	6 (2%)	0	100 100
1	F	253/258 (98%)	247 (98%)	6 (2%)	0	100 100
1	G	254/258 (98%)	251 (99%)	3 (1%)	0	100 100
1	H	253/258 (98%)	249 (98%)	4 (2%)	0	100 100
All	All	2026/2064 (98%)	1979 (98%)	45 (2%)	2 (0%)	48 41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	VAL
1	B	3	VAL

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	206/229 (90%)	205 (100%)	1 (0%)	86 86
1	B	207/229 (90%)	204 (99%)	3 (1%)	62 55
1	C	200/229 (87%)	198 (99%)	2 (1%)	73 68
1	D	208/229 (91%)	206 (99%)	2 (1%)	73 68
1	E	209/229 (91%)	204 (98%)	5 (2%)	44 32
1	F	201/229 (88%)	198 (98%)	3 (2%)	60 52
1	G	203/229 (89%)	202 (100%)	1 (0%)	86 86
1	H	201/229 (88%)	199 (99%)	2 (1%)	73 68
All	All	1635/1832 (89%)	1616 (99%)	19 (1%)	67 61

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	121	VAL
1	H	121	VAL
1	H	194	LEU
1	G	255	VAL
1	E	50	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	42	ASN
1	F	190	ASN
1	H	177	ASN
1	H	64	HIS
1	B	64	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 39 ligands modelled in this entry, 15 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	VVO	A	304	1,3	0,1,1	-	-	-	-	
3	SIN	F	303	8	7,7,7	1.27	0	8,8,8	1.63	1 (12%)
4	PGE	H	303	-	9,9,9	0.31	0	8,8,8	0.63	0
2	LYS	C	301	-	8,9,9	0.88	1 (12%)	7,10,10	1.11	1 (14%)
3	SIN	B	503	8	7,7,7	1.31	0	8,8,8	1.54	2 (25%)
4	PGE	A	303	-	9,9,9	0.32	0	8,8,8	0.53	0
4	PGE	F	304	-	9,9,9	0.31	0	8,8,8	0.71	0
7	FLC	G	303	-	12,12,12	1.62	3 (25%)	17,17,17	1.71	4 (23%)
2	LYS	F	302	-	8,9,9	1.01	1 (12%)	7,10,10	1.05	1 (14%)
3	SIN	C	302	8	7,7,7	1.28	0	8,8,8	1.51	2 (25%)
2	LYS	B	502	-	8,9,9	0.89	1 (12%)	7,10,10	1.13	1 (14%)
3	SIN	G	302	8	7,7,7	1.40	0	8,8,8	1.51	1 (12%)
4	PGE	B	501	-	9,9,9	0.33	0	8,8,8	0.67	0
2	LYS	G	301	-	8,9,9	0.83	1 (12%)	7,10,10	0.95	1 (14%)
3	SIN	H	302	8	7,7,7	1.23	0	8,8,8	1.58	2 (25%)
3	SIN	A	302	5	7,7,7	1.27	0	8,8,8	1.62	2 (25%)
2	LYS	H	301	-	8,9,9	0.83	0	7,10,10	0.89	1 (14%)
4	PGE	F	301	-	9,9,9	0.33	0	8,8,8	0.49	0
2	LYS	D	301	-	8,9,9	0.90	1 (12%)	7,10,10	0.94	1 (14%)
3	SIN	D	302	8	7,7,7	1.36	0	8,8,8	1.53	1 (12%)
2	LYS	A	301	-	8,9,9	0.89	1 (12%)	7,10,10	0.89	1 (14%)
7	FLC	B	504	-	12,12,12	1.55	2 (16%)	17,17,17	2.07	5 (29%)
3	SIN	E	302	8	7,7,7	1.36	0	8,8,8	1.38	1 (12%)
2	LYS	E	301	-	8,9,9	0.88	1 (12%)	7,10,10	1.06	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIN	F	303	8	-	3/5/5/5	-
4	PGE	H	303	-	-	5/7/7/7	-
2	LYS	C	301	-	-	1/9/9/9	-
3	SIN	B	503	8	-	3/5/5/5	-
4	PGE	A	303	-	-	4/7/7/7	-
4	PGE	F	304	-	-	6/7/7/7	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	FLC	G	303	-	-	7/16/16/16	-
2	LYS	F	302	-	-	0/9/9/9	-
3	SIN	C	302	8	-	4/5/5/5	-
2	LYS	B	502	-	-	0/9/9/9	-
3	SIN	G	302	8	-	2/5/5/5	-
4	PGE	B	501	-	-	3/7/7/7	-
2	LYS	G	301	-	-	1/9/9/9	-
3	SIN	H	302	8	-	2/5/5/5	-
3	SIN	A	302	5	-	2/5/5/5	-
2	LYS	H	301	-	-	1/9/9/9	-
4	PGE	F	301	-	-	3/7/7/7	-
2	LYS	D	301	-	-	0/9/9/9	-
3	SIN	D	302	8	-	4/5/5/5	-
2	LYS	A	301	-	-	0/9/9/9	-
7	FLC	B	504	-	-	11/16/16/16	-
3	SIN	E	302	8	-	2/5/5/5	-
2	LYS	E	301	-	-	0/9/9/9	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	G	303	FLC	CA-CB	-2.79	1.50	1.54
7	G	303	FLC	OHB-CB	-2.64	1.38	1.43
7	B	504	FLC	OHB-CB	-2.60	1.38	1.43
7	B	504	FLC	CA-CB	-2.49	1.50	1.54
2	B	502	LYS	OXT-C	-2.41	1.23	1.30

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	504	FLC	OB2-CBC-CB	4.54	121.86	113.14
7	B	504	FLC	OHB-CB-CBC	4.36	115.14	108.96
7	G	303	FLC	OB2-CBC-CB	4.25	121.30	113.14
7	G	303	FLC	CB-CA-CAC	-3.09	105.46	113.92
2	C	301	LYS	OXT-C-O	-2.73	117.88	124.08

There are no chirality outliers.

5 of 64 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	B	504	FLC	CA-CB-CBC-OB1
7	B	504	FLC	CA-CB-CBC-OB2
7	B	504	FLC	OHB-CB-CBC-OB1
7	B	504	FLC	OHB-CB-CBC-OB2
7	B	504	FLC	CBC-CB-CG-CGC



There are no ring outliers.



8 monomers are involved in 26 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	304	VVO	5	0
4	H	303	PGE	1	0
4	F	304	PGE	1	0
7	G	303	FLC	4	0
3	C	302	SIN	1	0
3	G	302	SIN	1	0
4	F	301	PGE	1	0
7	B	504	FLC	12	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/258 (98%)	-0.38	4 (1%) 70 75	21, 29, 44, 74	0
1	B	254/258 (98%)	-0.34	5 (1%) 64 70	22, 32, 47, 76	1 (0%)
1	C	254/258 (98%)	-0.32	5 (1%) 64 70	13, 32, 48, 77	1 (0%)
1	D	255/258 (98%)	-0.29	3 (1%) 76 79	13, 31, 49, 76	1 (0%)
1	E	255/258 (98%)	-0.27	4 (1%) 70 75	14, 32, 48, 74	1 (0%)
1	F	255/258 (98%)	-0.29	4 (1%) 70 75	23, 32, 48, 72	0
1	G	253/258 (98%)	-0.20	2 (0%) 82 87	14, 35, 51, 70	3 (1%)
1	H	253/258 (98%)	-0.02	4 (1%) 70 75	15, 38, 55, 76	2 (0%)
All	All	2033/2064 (98%)	-0.26	31 (1%) 71 76	13, 33, 51, 77	9 (0%)

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	255	VAL	4.0
1	H	3	VAL	3.8
1	A	3	VAL	3.6
1	C	255	VAL	3.4
1	A	255	VAL	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PGE	H	303	10/10	0.82	0.14	31,47,50,53	0
4	PGE	F	304	10/10	0.89	0.10	38,47,57,61	0
4	PGE	A	303	10/10	0.89	0.10	39,48,62,66	0
7	FLC	G	303	13/13	0.89	0.15	17,50,58,68	0
7	FLC	B	504	13/13	0.91	0.13	11,40,55,56	0
4	PGE	F	301	10/10	0.91	0.10	42,47,59,65	0
4	PGE	B	501	10/10	0.93	0.07	40,47,56,58	0
3	SIN	G	302	8/8	0.93	0.10	28,39,42,46	0
3	SIN	E	302	8/8	0.93	0.09	28,31,39,40	0
2	LYS	H	301	10/10	0.94	0.08	31,37,41,42	0
3	SIN	F	303	8/8	0.94	0.08	32,35,39,41	0
2	LYS	A	301	10/10	0.95	0.06	24,29,32,34	0
3	SIN	A	302	8/8	0.95	0.07	26,31,36,37	0
3	SIN	H	302	8/8	0.95	0.07	29,35,40,42	0
3	SIN	C	302	8/8	0.95	0.07	27,34,35,36	0
2	LYS	B	502	10/10	0.95	0.06	25,30,35,36	0
3	SIN	D	302	8/8	0.96	0.07	26,34,38,39	0
3	SIN	B	503	8/8	0.96	0.07	27,31,35,40	0
2	LYS	F	302	10/10	0.96	0.07	26,32,35,36	0
2	LYS	D	301	10/10	0.97	0.06	26,29,34,36	0
2	LYS	E	301	10/10	0.97	0.05	30,32,37,43	0
2	LYS	C	301	10/10	0.97	0.05	27,31,34,34	0
2	LYS	G	301	10/10	0.97	0.06	32,35,39,39	0
6	CL	D	304	1/1	0.99	0.04	30,30,30,30	0
6	CL	E	304	1/1	0.99	0.03	29,29,29,29	0
6	CL	F	306	1/1	0.99	0.02	28,28,28,28	0
6	CL	G	305	1/1	0.99	0.05	32,32,32,32	0
6	CL	H	305	1/1	0.99	0.03	33,33,33,33	0
6	CL	B	506	1/1	0.99	0.03	29,29,29,29	0
6	CL	C	304	1/1	0.99	0.04	29,29,29,29	0
8	V	C	303	1/1	0.99	0.04	22,22,22,22	0
8	V	D	303	1/1	0.99	0.06	20,20,20,20	0
8	V	G	304	1/1	0.99	0.06	23,23,23,23	0
8	V	H	304	1/1	0.99	0.03	25,25,25,25	0
5	VVO	A	304	2/2	1.00	0.05	19,19,19,37	0
8	V	E	303	1/1	1.00	0.06	20,20,20,20	0
8	V	F	305	1/1	1.00	0.04	22,22,22,22	0


Continued on next page...


Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	V	B	505	1/1	1.00	0.05	20,20,20,20	0
6	CL	A	305	1/1	1.00	0.02	28,28,28,28	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


































































Electron density around V E 303:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)









6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.