



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2025 – 02:04 PM EDT

PDB ID : 2ONG / pdb_00002ong
Title : Crystal Structure of of limonene synthase with 2-fluorogeranyl diphosphate (FGPP) enzymatically converted to 2-fluorolinalyl diphosphate (FLPP)
Authors : Hyatt, D.C.; Youn, B.; Croteau, R.; Kang, C.
Deposited on : 2007-01-23
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	
1	B	543	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	F3P	A	600	-	-	X	-
3	F3P	B	1600	-	-	X	-
4	BTB	A	605	-	-	X	-
4	BTB	B	1604	-	-	X	-
4	BTB	B	1605	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4S-limonene synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	543	Total	C	N	O	S	0	0	0
			4495	2871	761	843	20			
1	B	543	Total	C	N	O	S	0	0	0
			4491	2870	758	843	20			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	MET	GLU	engineered mutation	UNP Q40322
B	57	MET	GLU	engineered mutation	UNP Q40322

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mn	0	0
			3	3		
2	B	3	Total	Mn	0	0
			3	3		

- Molecule 3 is (3S)-2-fluoro-3,7-dimethylocta-1,6-dien-3-yl trihydrogen diphosphate (CCD ID: F3P) (formula: C₁₀H₁₉FO₇P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	O	P	0	0
			20	10	1	7	2		
3	B	1	Total	C	F	O	P	0	0
			20	10	1	7	2		

- Molecule 4 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	39	Total	O	0	0
			39	39		
5	B	54	Total	O	0	0
			54	54		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4S-limonene synthase



• Molecule 1: 4S-limonene synthase



K550	Q117
D551	R118
S552	M119
F553	G120
F554	E185
G555	L121
K556	L186
S557	L122
F558	K187
T559	E188
G560	D123
C561	H124
A562	F125
V563	N126
D564	Q127
L565	E128
G566	E129
R567	K130
M568	E131
A569	I132
Q570	L133
L571	G134
M572	
Y573	Y137
H574	L138
M575	D139
G576	H140
D577	H141
G578	Y142
H579	Y143
G580	K144
T581	M145
Q582	P146
Q583	F147
P584	P148
I585	K149
I586	E150
H587	R152
Q588	D153
Q589	L154
M590	Y155
T591	
K592	S158
T593	L159
L594	A160
F595	F161
E596	R162
P597	L163
F598	L164
A599	R165
	E166
	H167
	G168
	F169
	E174
	V175
	F176
	D177
	S178
	F179
	K180
	N181
	E182
	E183
	G184
	G185
	L187
	K187
	E188
	S189
	L190
	S191
	D192
	E193
	T194
	R195
	G196
	L197
	L198
	Q199
	L200
	Y201
	E202
	A203
	S204
	F205
	L206
	Y266
	L207
	T208
	E209
	G210
	E211
	T212
	T213
	L214
	E215
	S216
	A217
	R218
	E219
	F220
	A221
	T222
	K223
	F224
	L225
	E226
	E227
	K228
	V229
	N230
	E231
	G232
	G233
	V234
	L235
	G236
	D237
	L238
	T239
	L240
	R241
	L242
	A243
	Y244
	S245
	L246
	D247
	I248
	P249
	L250
	H251
	W252
	R253
	L254
	K255
	R256
	P257
	N258
	A259
	P260
	V261
	W262
	I263
	E264
	W265
	Y266
	R267
	K268
	R269
	P270
	D271
	K272
	K273
	P274
	V275
	V276
	L277
	E278
	L279
	A280
	I281
	L282
	D283
	L284
	K285
	I286
	V287
	Q288
	A289
	Q290
	F291
	Q292
	E293
	E294
	L295
	K296
	E297
	S298
	F299
	L300
	R301
	W301
	W302
	R303
	N304
	G305
	L306
	F307
	V308
	E309
	K310
	L311
	P312
	W313
	A314
	R315
	D316
	R317
	L318
	V319
	E320
	C321
	Y322
	F323
	W324
	W325
	N325
	T326
	I329
	R332
	Q333
	H334
	A335
	S336
	A337
	R338
	I339
	M340
	M341
	G342
	K343
	V344
	N345
	A346
	L347
	I348
	T349
	V350
	I351
	D352
	D353
	I354
	V357
	T360
	L361
	E362
	E363
	L364
	A365
	E366
	Q366
	F367
	T368
	D369
	L370
	I371
	R372
	K373
	W374
	D375
	L376
	N377
	S378
	I379
	D380
	Q381
	L382
	P383
	D384
	Y385
	K386
	Q387
	L388
	C389
	F390
	L391
	A392
	L393
	N394
	N395
	F396
	V397
	D398
	D399
	T400
	S401
	Y402
	D403
	Y404
	M405
	K406
	E407
	K408
	G409
	V410
	M411
	V412
	I413
	P414
	Y415
	L416
	R417
	Q418
	S419
	W420
	V421
	D422
	L423
	A424
	R425
	K426
	Y427
	M428
	V429
	E430
	A431
	R432
	W433
	G436
	L437
	H438
	K439
	P440
	S441
	L442
	E443
	E444
	Y445
	L446
	E447
	N448
	S449
	W450
	Q451
	S452
	L453
	S454
	G455
	P456
	C457
	M458
	L459
	T460
	H461
	I462
	F463
	F464
	R465
	V466
	T467
	D468
	S469
	F470
	T471
	K472
	E473
	T474
	Y475
	D476
	S477
	L478
	Y479
	K480
	Y481
	H482
	D483
	L484
	V485
	R486
	W487
	S488
	S489
	F490
	V491
	L492
	R493
	L494
	A495
	D496
	D497
	L498
	G499
	T500
	S501
	V502
	E503
	E504
	V505
	S506
	R507
	G508
	D509
	Y510
	P511
	K512
	S513
	L514
	Q515
	C516
	Y517
	H518
	S519
	D520
	P521
	N522
	A526
	E527
	A528
	R529
	K530
	H531
	W534
	L535
	T536
	A537
	E538
	V539
	W540
	K541
	M542
	M543
	N544
	A545
	E546
	R547
	V548
	S549

4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	200.48Å 200.48Å 123.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.70 10.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.70) 93.1 (10.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.61Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.208 , 0.241 0.220 , 0.230	Depositor DCC
R_{free} test set	3503 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 97.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.478 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9181	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F3P, BTB, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	39/4607 (0.8%)	2.37	351/6234 (5.6%)
1	B	1.15	42/4603 (0.9%)	2.27	324/6230 (5.2%)
All	All	1.17	81/9210 (0.9%)	2.32	675/12464 (5.4%)

The worst 5 of 81 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	368	THR	C-O	12.06	1.38	1.24
1	B	368	THR	C-O	9.90	1.36	1.24
1	B	413	ILE	CA-CB	8.17	1.59	1.53
1	B	255	LYS	N-CA	-6.65	1.38	1.46
1	A	482	HIS	N-CA	-6.45	1.38	1.46

The worst 5 of 675 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	480	LYS	N-CA-C	-18.45	91.25	111.36
1	A	313	PHE	N-CA-C	-16.93	92.29	113.97
1	A	551	ASP	N-CA-C	-16.47	88.00	111.56
1	A	410	VAL	N-CA-C	15.58	129.93	108.11
1	A	313	PHE	CA-C-N	15.53	142.72	120.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4495	0	4346	592	0
1	B	4491	0	4342	536	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	20	0	18	9	0
3	B	20	0	18	13	0
4	A	28	0	38	21	0
4	B	28	0	38	40	0
5	A	39	0	0	1	0
5	B	54	0	0	2	0
All	All	9181	0	8800	1186	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 66.

The worst 5 of 1186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1604:BTB:N	4:B:1604:BTB:C2	1.68	1.56
4:B:1605:BTB:N	4:B:1605:BTB:C2	1.69	1.51
4:A:605:BTB:N	4:A:605:BTB:C2	1.71	1.49
1:B:579:HIS:CD2	3:B:1600:F3P:H92	1.67	1.28
4:B:1605:BTB:C7	4:B:1605:BTB:H32	1.72	1.19

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	541/543 (100%)	400 (74%)	93 (17%)	48 (9%)	0 0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	541/543 (100%)	409 (76%)	86 (16%)	46 (8%)	0	1
All	All	1082/1086 (100%)	809 (75%)	179 (16%)	94 (9%)	0	0

5 of 94 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	LEU
1	A	78	SER
1	A	83	ASP
1	A	125	PHE
1	A	152	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/492 (100%)	331 (68%)	159 (32%)	0	0
1	B	490/492 (100%)	323 (66%)	167 (34%)	0	0
All	All	980/984 (100%)	654 (67%)	326 (33%)	0	0

5 of 326 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	242	ILE
1	B	459	LEU
1	B	263	ILE
1	B	362	GLU
1	B	510	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	145	ASN
1	B	579	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	395	ASN
1	B	589	GLN
1	B	544	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	F3P	A	600	2	14,19,19	2.54	5 (35%)	20,29,29	1.67	3 (15%)
4	BTB	A	605	-	13,13,13	4.54	6 (46%)	7,16,16	1.18	1 (14%)
4	BTB	B	1604	-	13,13,13	4.55	6 (46%)	7,16,16	1.13	1 (14%)
4	BTB	A	604	-	13,13,13	2.01	4 (30%)	7,16,16	0.57	0
4	BTB	B	1605	-	13,13,13	4.32	6 (46%)	7,16,16	1.08	1 (14%)
3	F3P	B	1600	2	14,19,19	2.49	4 (28%)	20,29,29	2.05	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	F3P	A	600	2	-	4/17/25/25	-
4	BTB	A	605	-	-	12/21/21/21	-
4	BTB	B	1604	-	-	11/21/21/21	-
4	BTB	A	604	-	-	2/21/21/21	-
4	BTB	B	1605	-	-	8/21/21/21	-
3	F3P	B	1600	2	-	3/17/25/25	-

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	605	BTB	C2-N	11.69	1.71	1.48
4	B	1605	BTB	C2-N	10.65	1.69	1.48
4	B	1604	BTB	C2-N	10.43	1.68	1.48
4	B	1604	BTB	C5-N	7.93	1.59	1.48
4	A	605	BTB	C5-N	7.78	1.59	1.48

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1600	F3P	O1-C3-C10	-6.21	86.95	108.11
3	B	1600	F3P	C5-C6-C7	-4.78	111.69	127.64
3	A	600	F3P	C5-C6-C7	-4.57	112.42	127.64
3	A	600	F3P	C10-C3-C2	3.09	116.41	110.69
4	A	605	BTB	C6-C5-N	2.61	121.79	111.59

There are no chirality outliers.

5 of 40 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	600	F3P	C2-C3-C4-C5
3	A	600	F3P	C10-C3-C4-C5
3	B	1600	F3P	C3-C4-C5-C6
4	A	605	BTB	O1-C1-C2-C4
4	A	605	BTB	O1-C1-C2-N


There are no ring outliers.

6 monomers are involved in 83 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	F3P	9	0
4	A	605	BTB	18	0
4	B	1604	BTB	19	0
4	A	604	BTB	3	0
4	B	1605	BTB	21	0
3	B	1600	F3P	13	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	543/543 (100%)	-1.74	0 100 100	18, 46, 94, 100	0
1	B	543/543 (100%)	-1.75	0 100 100	16, 46, 96, 100	0
All	All	1086/1086 (100%)	-1.75	0 100 100	16, 46, 95, 100	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BTB	A	605	14/14	0.98	0.05	98,100,100,100	0
4	BTB	A	604	14/14	0.99	0.07	83,90,96,100	0
4	BTB	B	1604	14/14	0.99	0.05	67,89,100,100	0
4	BTB	B	1605	14/14	0.99	0.04	93,100,100,100	0
2	MN	B	1602	1/1	1.00	0.02	49,49,49,49	0
2	MN	B	1603	1/1	1.00	0.01	44,44,44,44	0
3	F3P	A	600	20/20	1.00	0.05	54,69,76,82	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	F3P	B	1600	20/20	1.00	0.04	44,71,100,100	0
2	MN	A	601	1/1	1.00	0.04	52,52,52,52	0
2	MN	A	602	1/1	1.00	0.03	41,41,41,41	0
2	MN	A	603	1/1	1.00	0.03	43,43,43,43	0
2	MN	B	1601	1/1	1.00	0.04	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around F3P A 600:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.