



wwPDB EM Validation Summary Report i

Jul 24, 2025 – 02:05 PM EDT

PDB ID : 9OP5 / pdb_00009op5
EMDB ID : EMD-70679
Title : Herpes simplex virus type 1 (HSV-1) B-capsid pUL6 portal protein, dodecameric complex
Authors : Crofut, E.H.; Kashyap, S.; Stevens, A.; Jih, J.; Liu, Y.-T.; Zhou, Z.H.
Deposited on : 2025-05-16
Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
MolProbity : 4-5-2 with Phenix2.0rc1
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	I	635	 :	99%		
1	J	635	 :	99%		
1	K	635	 :	99%		
1	L	635	 :	99%		
2	a	676	 :	56%	5%	38%
2	b	676	 :	56%	5%	38%
2	c	676	 :	55%	6%	38%
2	d	676	 :	54%	7%	38%
2	e	676	 :	56%	5%	38%
2	f	676	 :	56%	6%	38%
2	g	676	 :	55%	7%	38%
2	h	676	 :	55%	7%	38%
2	i	676	 :	55%	7%	38%
2	j	676	 :	56%	5%	38%
2	k	676	 :	56%	5%	38%
2	l	676	 :	56%	5%	38%

2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 40476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid scaffolding protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	7	Total	C	N	O	0	0
			56	38	7	11		
1	B	7	Total	C	N	O	0	0
			56	38	7	11		
1	C	7	Total	C	N	O	0	0
			56	38	7	11		
1	D	7	Total	C	N	O	0	0
			56	38	7	11		
1	E	7	Total	C	N	O	0	0
			56	38	7	11		
1	F	7	Total	C	N	O	0	0
			56	38	7	11		
1	G	7	Total	C	N	O	0	0
			56	38	7	11		
1	H	7	Total	C	N	O	0	0
			56	38	7	11		
1	I	7	Total	C	N	O	0	0
			56	38	7	11		
1	J	7	Total	C	N	O	0	0
			56	38	7	11		
1	K	7	Total	C	N	O	0	0
			56	38	7	11		
1	L	7	Total	C	N	O	0	0
			56	38	7	11		

- Molecule 2 is a protein called Capsid portal protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	a	416	Total	C	N	O	S	0
			3317	2110	601	597	9	
2	b	416	Total	C	N	O	S	0
			3317	2110	601	597	9	
2	c	416	Total	C	N	O	S	0
			3317	2110	601	597	9	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	Trace
2	d	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	e	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	f	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	g	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	h	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	i	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	j	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	k	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		
2	l	416	Total	C	N	O	S	0	0
			3317	2110	601	597	9		

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Capsid scaffolding protein

Chain A: :



- Molecule 1: Capsid scaffolding protein

Chain B: :



- Molecule 1: Capsid scaffolding protein

Chain C: 99%

SER PRO VAL LEU PRO PRO MET ASN PRO VAL PRO THR SER GLY GLY GLY ASP GLY ASP SER TYR LEU TRP ILE PRO ALA SER HIS TYR GLN GLN LEU VAL GLY HIS ALA PRO GLN PRO GLN PRO HIS SER PHE PHE PHE PRO ALA ALA ALA GLY

www.english-test.net

- Molecule 1: Capsid scaffolding protein

Chain D:  99%

GLY HIS THR TYR LYS PHE GLN GLU ALA SER GLU LYS GLY ALA GLU PRO VAL ARG GLY TYR LYS ASN GLY ALA PRO SER GLY THR ASP ILE PRO PRO CLN GLY SER ILE ASP ARG CYS PRO CLN GLY VAL ARG GLN ARG GLY GLY VAL ALA LEU

- Molecule 1: Capsid scaffolding protein

Chain E:  99%

- Molecule 1: Capsid scaffolding protein

Chain F: .

99%

- Molecule 1: Capsid scaffolding protein

Chain G: :

99%

- Molecule 1: Capsid scaffolding protein

Chain H: :

99%

- Molecule 1: Capsid scaffolding protein

Chain I: :

For more information, contact the Office of the Vice President for Research and Economic Development at 319-273-2500 or research@uiowa.edu.

1

For more information, contact the Office of the Vice President for Research and Economic Development at 505-274-3000 or research@unm.edu.

- Molecule 1: Capsid scaffolding protein

Chain J: :

99%


- Molecule 1: Capsid scaffolding protein

Chain K: ■

000%





ALA SER ARG GLO GLY
ALA ARG ARG LEU ALA
ALA GLU GLU TLE ALA
ALA LEU SER ALA
ALA GLY ARG THR TRP
ALA GLU ALA LEU PRO
ALA VAL GLU ALA HIS
ALA LEU LEU THR HIS
ALA SER THR ALA LEU
ALA ASN ASN MET MET
ALA ASN ARG ARG SER
ALA GLU ARG ARG LEU
ALA GLN GLN ALA GLY
ALA GLY ILE ILE ALA

GLY H.S.
THR TYR
LEU GLN
ALA SER
GLU GLY
PHE LYS
LYS ME
TRP GLY
ALA GLU
ALA PRO
VAL SER
ALA PRO
ALA ARG
ALA GLY
TYR LYS
ASN GLY
ALA PRO
GLU SER
ILE ASP
LLE PRO
ALA PRO
ALA GLN
GLY GLY
ASP ARG
CYS PRO
ILE VAL
VAL ARG
GLN ARG
GLY GLY
VAL ALA
ALA LEU



- Molecule 1: Capsid scaffolding protein


Chain L: :  99%




- Molecule 2: Capsid portal protein

Chain a: 56% 5% 38%






- Molecule 2: Capsid portal protein

Chain b:  56% 5% 38%


A horizontal progress bar for 'Chain b'. The bar is divided into three segments: a green segment on the left representing 56%, a yellow segment in the middle representing 5%, and a grey segment on the right representing 38%. The total length of the bar is 100%.




- Molecule 2: Capsid portal protein

Chain c: 55% 6% 38%

A horizontal progress bar for 'Chain c'. The bar is divided into three segments: a red segment on the left, a green segment in the middle, and a grey segment on the right. The green segment is labeled '55%' below it. The grey segment is labeled '38%' below it. A small black dot is positioned at the 6% mark on the grey segment.




- Molecule 2: Capsid portal protein




- Molecule 2: Capsid portal protein






- Molecule 2: Capsid portal protein




- Molecule 2: Capsid portal protein





- Molecule 2: Capsid portal protein


Chain h:



- Molecule 2: Capsid portal protein


Chain i:






- Molecule 2: Capsid portal protein


Chain j: 56% 5% 38%



- Molecule 2: Capsid portal protein

Chain k:  56% 5% 38%

A horizontal progress bar for 'Chain k'. The bar is divided into three segments: a green segment on the left, a yellow segment in the middle, and a grey segment on the right. A red dot is positioned on the green segment, indicating the current progress. The text 'Chain k:' is to the left of the bar. Below the bar, the values '56%', '5%', and '38%' are displayed, corresponding to the lengths of the green, yellow, and grey segments respectively.



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C12	Depositor
Number of particles used	799092	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	45	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.012	Depositor
Minimum map value	-1.402	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.044	Depositor
Recommended contour level	0.187	Depositor
Map size (Å)	528.0, 528.0, 528.0	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.08	0/59	0.25	0/80
1	B	0.10	0/59	0.26	0/80
1	C	0.12	0/59	0.26	0/80
1	D	0.08	0/59	0.25	0/80
1	E	0.09	0/59	0.24	0/80
1	F	0.09	0/59	0.26	0/80
1	G	0.10	0/59	0.26	0/80
1	H	0.09	0/59	0.27	0/80
1	I	0.13	0/59	0.36	0/80
1	J	0.09	0/59	0.26	0/80
1	K	0.08	0/59	0.25	0/80
1	L	0.08	0/59	0.28	0/80
2	a	0.11	0/3401	0.28	0/4626
2	b	0.11	0/3401	0.27	0/4626
2	c	0.12	0/3401	0.29	0/4626
2	d	0.11	0/3401	0.28	0/4626
2	e	0.11	0/3401	0.28	0/4626
2	f	0.11	0/3401	0.28	0/4626
2	g	0.12	0/3401	0.29	0/4626
2	h	0.11	0/3401	0.28	0/4626
2	i	0.11	0/3401	0.29	0/4626
2	j	0.11	0/3401	0.29	0/4626
2	k	0.12	0/3401	0.30	0/4626
2	l	0.12	0/3401	0.28	0/4626
All	All	0.11	0/41520	0.28	0/56472

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	56	0	48	1	0
1	B	56	0	48	1	0
1	C	56	0	48	1	0
1	D	56	0	48	2	0
1	E	56	0	48	1	0
1	F	56	0	48	1	0
1	G	56	0	48	3	0
1	H	56	0	48	1	0
1	I	56	0	48	4	0
1	J	56	0	48	1	0
1	K	56	0	48	1	0
1	L	56	0	48	1	0
2	a	3317	0	3264	26	0
2	b	3317	0	3264	25	0
2	c	3317	0	3264	31	0
2	d	3317	0	3264	33	0
2	e	3317	0	3264	23	0
2	f	3317	0	3264	29	0
2	g	3317	0	3264	29	0
2	h	3317	0	3264	32	0
2	i	3317	0	3264	31	0
2	j	3317	0	3264	25	0
2	k	3317	0	3264	30	0
2	l	3317	0	3264	26	0
All	All	40476	0	39744	306	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 306 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:e:78:LEU:HD13	2:e:603:THR:HG21	1.64	0.79
2:a:500:SER:HB2	2:a:503:MET:HE2	1.65	0.77
2:g:503:MET:HE2	2:h:497:ILE:HD12	1.71	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:a:81:THR:HG21	2:a:605:LEU:HD13	1.73	0.70
2:d:499:VAL:HG13	2:d:501:LYS:HE3	1.75	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	5/635 (1%)	5 (100%)	0	0	100 100
1	B	5/635 (1%)	5 (100%)	0	0	100 100
1	C	5/635 (1%)	4 (80%)	1 (20%)	0	100 100
1	D	5/635 (1%)	5 (100%)	0	0	100 100
1	E	5/635 (1%)	5 (100%)	0	0	100 100
1	F	5/635 (1%)	5 (100%)	0	0	100 100
1	G	5/635 (1%)	5 (100%)	0	0	100 100
1	H	5/635 (1%)	5 (100%)	0	0	100 100
1	I	5/635 (1%)	5 (100%)	0	0	100 100
1	J	5/635 (1%)	5 (100%)	0	0	100 100
1	K	5/635 (1%)	5 (100%)	0	0	100 100
1	L	5/635 (1%)	5 (100%)	0	0	100 100
2	a	412/676 (61%)	399 (97%)	13 (3%)	0	100 100
2	b	412/676 (61%)	403 (98%)	9 (2%)	0	100 100
2	c	412/676 (61%)	404 (98%)	8 (2%)	0	100 100
2	d	412/676 (61%)	403 (98%)	9 (2%)	0	100 100
2	e	412/676 (61%)	402 (98%)	10 (2%)	0	100 100
2	f	412/676 (61%)	400 (97%)	12 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	g	412/676 (61%)	400 (97%)	12 (3%)	0	100 100
2	h	412/676 (61%)	402 (98%)	10 (2%)	0	100 100
2	i	412/676 (61%)	401 (97%)	11 (3%)	0	100 100
2	j	412/676 (61%)	400 (97%)	12 (3%)	0	100 100
2	k	412/676 (61%)	400 (97%)	12 (3%)	0	100 100
2	l	412/676 (61%)	402 (98%)	10 (2%)	0	100 100
All	All	5004/15732 (32%)	4875 (97%)	129 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	5/475 (1%)	5 (100%)	0	100 100
1	B	5/475 (1%)	5 (100%)	0	100 100
1	C	5/475 (1%)	5 (100%)	0	100 100
1	D	5/475 (1%)	5 (100%)	0	100 100
1	E	5/475 (1%)	5 (100%)	0	100 100
1	F	5/475 (1%)	5 (100%)	0	100 100
1	G	5/475 (1%)	5 (100%)	0	100 100
1	H	5/475 (1%)	5 (100%)	0	100 100
1	I	5/475 (1%)	5 (100%)	0	100 100
1	J	5/475 (1%)	5 (100%)	0	100 100
1	K	5/475 (1%)	5 (100%)	0	100 100
1	L	5/475 (1%)	5 (100%)	0	100 100
2	a	347/534 (65%)	347 (100%)	0	100 100
2	b	347/534 (65%)	347 (100%)	0	100 100
2	c	347/534 (65%)	347 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	d	347/534 (65%)	347 (100%)	0	100	100
2	e	347/534 (65%)	347 (100%)	0	100	100
2	f	347/534 (65%)	346 (100%)	1 (0%)	91	96
2	g	347/534 (65%)	347 (100%)	0	100	100
2	h	347/534 (65%)	347 (100%)	0	100	100
2	i	347/534 (65%)	347 (100%)	0	100	100
2	j	347/534 (65%)	347 (100%)	0	100	100
2	k	347/534 (65%)	347 (100%)	0	100	100
2	l	347/534 (65%)	347 (100%)	0	100	100
All	All	4224/12108 (35%)	4223 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	f	236	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
2	h	548	ASN
2	k	110	ASN
2	h	622	HIS
2	j	235	ASN
2	k	622	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-70679. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [\(i\)](#)


6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240


6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 261




Y Index: 219




Z Index: 333


6.3.2 Raw map



X Index: 260



Y Index: 260






Z Index: 332




The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)

6.4.1 Primary map




6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [\(i\)](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.187. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [\(i\)](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_70679_msk_1.map [\(i\)](#)



X



Y




Z

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 360 nm^3 ; this corresponds to an approximate mass of 325 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.286 \AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.47	3.83	3.56
Unmasked-calculated*	4.38	7.67	4.43

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.38 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [\(i\)](#)


This section contains information regarding the fit between EMDB map EMD-70679 and PDB model 9OP5. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [\(i\)](#)




The images above show the 3D surface view of the map at the recommended contour level 0.187 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.187).

9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 84% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.187) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7940	0.4740
A	0.6480	0.4120
B	0.6110	0.4040
C	0.7220	0.4190
D	0.6300	0.4170
E	0.6110	0.4010
F	0.6480	0.4180
G	0.6300	0.4050
H	0.6670	0.4080
I	0.6300	0.4050
J	0.6300	0.4190
K	0.6480	0.4100
L	0.6480	0.3840
a	0.7930	0.4750
b	0.7970	0.4740
c	0.8000	0.4740
d	0.7940	0.4760
e	0.7930	0.4770
f	0.7980	0.4740
g	0.7960	0.4750
h	0.7960	0.4770
i	0.7980	0.4740
j	0.7980	0.4740
k	0.7950	0.4730
l	0.7970	0.4730

