



wwPDB EM Validation Summary Report i

Dec 18, 2025 – 03:50 PM EST

PDB ID : 9OT6 / pdb_00009ot6
EMDB ID : EMD-70826
Title : Cryo-EM structure of the PI4KA complex bound to an EFR3 interfering nanobody (F3IN)
Authors : Shaw, A.L.; Suresh, S.; Yip, C.K.; Burke, J.E.
Deposited on : 2025-05-26
Resolution : 3.54 Å(reported)
Based on initial models : 9BAX, .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev129
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 3.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	12891 (3.04 - 4.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 42678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4-kinase alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1724	Total	C	N	O	S	0	0
			13842	8951	2305	2489	97		

Mol	Chain	Residues	Total	C	N	O	S	AltConf	Trace
1	B	1724	Total	C	N	O	S	0	0
			13842	8951	2305	2489	97		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P42356
A	0	SER	-	expression tag	UNP P42356
B	-1	GLY	-	expression tag	UNP P42356
B	0	SER	-	expression tag	UNP P42356

- Molecule 2 is a protein called EFR3 interfering Nanobody (F3IN).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	104	Total	C	N	O	S	0	0
			811	511	141	154	5		

Mol	Chain	Residues	Total	C	N	O	S	AltConf	Trace
2	H	104	811	511	141	154	5	0	0

- Molecule 3 is a protein called Tetrastricopeptide repeat protein 7B.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	592	Total	C	N	O	S	0	0
			4676	2983	811	855	27		

Mol	Chain	Residues	Total	C	N	O	S	AltConf	Trace
3	F	592	4676	2983	811	855	27	0	0

- Molecule 4 is a protein called Hyccin.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	252	Total	C	N	O	S	0	0
			2010	1309	323	366	12		


Mol	Chain	Residues	Total	C	N	O	S	AltConf	Trace
4	G	252	Total	C	N	O	S	0	0
			2010	1309	323	366	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphatidylinositol 4-kinase alpha








- Molecule 2: EFR3 interfering Nanobody (F3IN)





- Molecule 2: EFR3 interfering Nanobody (F3IN)



- Molecule 3: Tetratricopeptide repeat protein 7B







4 Experimental information i

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	282982	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.210	Depositor
Minimum map value	-0.034	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.0371	Depositor
Map size (Å)	431.19998, 431.19998, 431.19998	wwPDB
Map dimensions	560, 560, 560	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.77, 0.77, 0.77	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.10	0/14149	0.26	0/19141
1	B	0.10	0/14149	0.26	0/19141
2	C	0.11	0/826	0.33	0/1116
2	H	0.11	0/826	0.33	0/1116
3	D	0.09	0/4762	0.25	0/6443
3	F	0.09	0/4762	0.25	0/6443
4	E	0.10	0/2058	0.28	0/2793
4	G	0.10	0/2058	0.28	0/2793
All	All	0.10	0/43590	0.26	0/58986

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	13842	0	13974	193	0
1	B	13842	0	13974	197	0
2	C	811	0	771	20	0
2	H	811	0	771	15	0
3	D	4676	0	4734	66	0
3	F	4676	0	4734	68	0
4	E	2010	0	2017	27	0
4	G	2010	0	2017	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	42678	0	42992	598	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 598 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:92:THR:HA	2:H:120:VAL:O	1.73	0.88
2:C:92:THR:HA	2:C:120:VAL:O	1.73	0.88
3:F:378:GLN:HB3	3:F:381:MET:HE1	1.56	0.88
3:D:378:GLN:HB3	3:D:381:MET:HE1	1.56	0.87
3:D:347:ARG:HG3	3:D:349:PRO:HD3	1.66	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1676/2104 (80%)	1605 (96%)	70 (4%)	1 (0%)	48 79
1	B	1676/2104 (80%)	1605 (96%)	70 (4%)	1 (0%)	48 79
2	C	96/165 (58%)	91 (95%)	5 (5%)	0	100 100
2	H	96/165 (58%)	91 (95%)	5 (5%)	0	100 100
3	D	580/843 (69%)	571 (98%)	9 (2%)	0	100 100
3	F	580/843 (69%)	571 (98%)	9 (2%)	0	100 100
4	E	242/308 (79%)	232 (96%)	10 (4%)	0	100 100
4	G	242/308 (79%)	233 (96%)	9 (4%)	0	100 100
All	All	5188/6840 (76%)	4999 (96%)	187 (4%)	2 (0%)	100 100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1980	PHE
1	B	1980	PHE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1527/1842 (83%)	1505 (99%)	22 (1%)	62 80
1	B	1527/1842 (83%)	1506 (99%)	21 (1%)	62 80
2	C	82/129 (64%)	80 (98%)	2 (2%)	44 68
2	H	82/129 (64%)	80 (98%)	2 (2%)	44 68
3	D	494/712 (69%)	484 (98%)	10 (2%)	50 72
3	F	494/712 (69%)	484 (98%)	10 (2%)	50 72
4	E	227/277 (82%)	224 (99%)	3 (1%)	65 81
4	G	227/277 (82%)	224 (99%)	3 (1%)	65 81
All	All	4660/5920 (79%)	4587 (98%)	73 (2%)	58 76

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	E	244	ILE
2	H	49	LEU
3	F	350	GLU
3	F	576	LEU
1	B	200	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 66 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	276	GLN
3	F	763	HIS
2	H	119	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	438	ASN
1	B	91	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-70826. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 280



Y Index: 280




Z Index: 280


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 284



Y Index: 236




Z Index: 305


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [\(i\)](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0371. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [\(i\)](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 438 nm³; this corresponds to an approximate mass of 396 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.282 \AA^{-1}


8 Fourier-Shell correlation [i](#)

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit i


This section contains information regarding the fit between EMDB map EMD-70826 and PDB model 9OT6. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay i




The images above show the 3D surface view of the map at the recommended contour level 0.0371 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0371).

9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 92% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0371) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7980	0.3470
A	0.8570	0.3560
B	0.8620	0.3520
C	0.7180	0.3140
D	0.7060	0.3440
E	0.6120	0.3240
F	0.6940	0.3410
G	0.6550	0.3170
H	0.7420	0.3050

