

wwPDB X-ray Structure Validation Summary Report (i)

Jul 8, 2025 – 06:12 PM EDT

PDB ID : 9PG6 / pdb 00009pg6

Title : Crystal structure of Glyceraldehyde-3-phosphate dehydrogenase from Borde-

tella pertussis (NAD bound)

Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural

Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2025-07-07

Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	344	93%	
1	В	344	90%	7% ••
1	С	344	92%	5% •
1	D	344	90%	8% •



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	E	344	90%	8% •
1	F	344	93%	5% •
1	G	344	93%	5% •
1	Н	344	93%	5% • •



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 21477 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	336	Total	С	N	О	S	0	3	0
1	A	330	2558	1608	447	494	9	U) 	
1	В	336	Total	С	N	О	S	0	4	0
1	Б	330	2573	1618	454	492	9	U	4	
1	С	336	Total	С	N	О	S	0	1	0
1		330	2556	1608	446	492	10	U	1	
1	D	336	Total	С	N	О	S	0	4	0
1	D	330	2573	1618	452	493	10	U		
1	Е	336	Total	С	N	О	S	0	4	0
1	ינו	330	2558	1608	449	492	9	U	4	
1	F	336	Total	С	N	О	S	0	2	0
1	Г	330	2553	1605	447	492	9	U	2	
1	G	336	Total	С	N	О	S	0	7	0
1	G	330	2591	1629	458	495	9	U	1	
1	Н	337	Total	С	N	О	S	0	3	0
1	11	997	2574	1618	452	494	10	0	3	

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP Q7VZB9
A	-6	ALA	-	expression tag	UNP Q7VZB9
A	-5	HIS	-	expression tag	UNP Q7VZB9
A	-4	HIS	-	expression tag	UNP Q7VZB9
A	-3	HIS	-	expression tag	UNP Q7VZB9
A	-2	HIS	-	expression tag	UNP Q7VZB9
A	-1	HIS	-	expression tag	UNP Q7VZB9
A	0	HIS	-	expression tag	UNP Q7VZB9
В	-7	MET	-	expression tag	UNP Q7VZB9
В	-6	ALA	-	expression tag	UNP Q7VZB9
В	-5	HIS	-	expression tag	UNP Q7VZB9
В	-4	HIS	-	expression tag	UNP Q7VZB9
В	-3	HIS	-	expression tag	UNP Q7VZB9



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-2	HIS	-	expression tag	UNP Q7VZB9
В	-1	HIS	_	expression tag	UNP Q7VZB9
В	0	HIS	_	expression tag	UNP Q7VZB9
С	-7	MET	-	expression tag	UNP Q7VZB9
С	-6	ALA	-	expression tag	UNP Q7VZB9
С	-5	HIS	-	expression tag	UNP Q7VZB9
С	-4	HIS	-	expression tag	UNP Q7VZB9
С	-3	HIS	-	expression tag	UNP Q7VZB9
С	-2	HIS	-	expression tag	UNP Q7VZB9
С	-1	HIS	-	expression tag	UNP Q7VZB9
С	0	HIS	-	expression tag	UNP Q7VZB9
D	-7	MET	-	expression tag	UNP Q7VZB9
D	-6	ALA	-	expression tag	UNP Q7VZB9
D	-5	HIS	-	expression tag	UNP Q7VZB9
D	-4	HIS	-	expression tag	UNP Q7VZB9
D	-3	HIS	-	expression tag	UNP Q7VZB9
D	-2	HIS	-	expression tag	UNP Q7VZB9
D	-1	HIS	-	expression tag	UNP Q7VZB9
D	0	HIS	-	expression tag	UNP Q7VZB9
Е	-7	MET	-	expression tag	UNP Q7VZB9
Е	-6	ALA	_	expression tag	UNP Q7VZB9
Е	-5	HIS	-	expression tag	UNP Q7VZB9
Е	-4	HIS	-	expression tag	UNP Q7VZB9
Е	-3	HIS	-	expression tag	UNP Q7VZB9
Е	-2	HIS	-	expression tag	UNP Q7VZB9
Е	-1	HIS	-	expression tag	UNP Q7VZB9
Е	0	HIS	-	expression tag	UNP Q7VZB9
F	-7	MET	-	expression tag	UNP Q7VZB9
F	-6	ALA	-	expression tag	UNP Q7VZB9
F	-5	HIS	-	expression tag	UNP Q7VZB9
F	-4	HIS	-	expression tag	UNP Q7VZB9
F	-3	HIS	-	expression tag	UNP Q7VZB9
F	-2	HIS	-	expression tag	UNP Q7VZB9
F	-1	HIS	-	expression tag	UNP Q7VZB9
F	0	HIS	_	expression tag	UNP Q7VZB9
G	-7	MET	-	expression tag	UNP Q7VZB9
G	-6	ALA	-	expression tag	UNP Q7VZB9
G	-5	HIS	-	expression tag	UNP Q7VZB9
G	-4	HIS	-	expression tag	UNP Q7VZB9
G	-3	HIS	-	expression tag	UNP Q7VZB9
G	-2	HIS	-	expression tag	UNP Q7VZB9
G	-1	HIS	-	expression tag	UNP Q7VZB9



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP Q7VZB9
Н	-7	MET	-	expression tag	UNP Q7VZB9
Н	-6	ALA	-	expression tag	UNP Q7VZB9
Н	-5	HIS	-	expression tag	UNP Q7VZB9
Н	-4	HIS	-	expression tag	UNP Q7VZB9
Н	-3	HIS	-	expression tag	UNP Q7VZB9
Н	-2	HIS	-	expression tag	UNP Q7VZB9
Н	-1	HIS	-	expression tag	UNP Q7VZB9
Н	0	HIS	-	expression tag	UNP Q7VZB9

• Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
	Λ	1	44	21	7	14	2	U	U
2	В	1	Total	С	N	Ο	Р	0	0
	Ъ	1	44	21	7	14	2	U	U
2	С	1	Total	С	N	О	Р	0	0
		1	44	21	7	14	2	U	0
2	D	1	Total	С	N	О	Р	0	0
	ט	1	44	21	7	14	2	U	0
2	Е	1	Total	С	N	О	Р	0	0
	<u> 1</u> 2	1	44	21	7	14	2	U	0
2	F	1	Total	С	N	О	Р	0	0
	I'	1	44	21	7	14	2	U	U



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	C	1	Total	С	N	О	Р	0	0
	G	1	44	21	7	14	2	U	0
9	П	1	Total	С	N	О	Р	0	0
	п	1	44	21	7	14	2	U	0

 \bullet Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0
3	Н	1	Total Mg 1 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	65	Total O 65 65	0	0
4	В	62	Total O 62 62	0	0
4	С	77	Total O 77 77	0	0
4	D	80	Total O 80 80	0	0
4	Е	64	Total O 64 64	0	0
4	F	70	Total O 70 70	0	0
4	G	83	Total O 83 83	0	0
4	Н	83	Total O 83 83	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase



• Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.83Å 130.06Å 140.97Å	Depositor
a, b, c, α , β , γ	90.00° 90.86° 90.00°	Depositor
Resolution (Å)	49.03 - 2.15	Depositor
rtesolution (A)	49.03 - 2.15	EDS
% Data completeness	99.7 (49.03-2.15)	Depositor
(in resolution range)	99.8 (49.03-2.15)	EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.08 (at 2.16Å)	Xtriage
Refinement program	PHENIX (dev_5438: ???)	Depositor
D D.	0.193 , 0.230	Depositor
R, R_{free}	0.199 , 0.233	DCC
R_{free} test set	7791 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.795	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 38.3	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21477	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 55.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1735e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.21	0/2611	0.43	0/3551
1	В	0.20	0/2629	0.41	0/3571
1	С	0.23	0/2603	0.46	0/3536
1	D	0.20	0/2629	0.44	0/3572
1	Ε	0.19	0/2614	0.41	0/3555
1	F	0.20	0/2603	0.43	0/3539
1	G	0.24	0/2656	0.45	0/3608
1	Н	0.24	0/2628	0.44	0/3571
All	All	0.21	0/20973	0.43	0/28503

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Е	0	2
1	Н	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Е	76[A]	ARG	Sidechain
1	Е	76[B]	ARG	Sidechain
1	Н	19	ARG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2558	0	2568	7	0
1	В	2573	0	2601	15	0
1	С	2556	0	2580	8	0
1	D	2573	0	2601	15	0
1	Е	2558	0	2566	18	0
1	F	2553	0	2564	7	0
1	G	2591	0	2624	9	0
1	Н	2574	0	2589	13	0
2	A	44	0	26	1	0
2	В	44	0	26	0	0
2	С	44	0	26	0	0
2	D	44	0	26	0	0
2	Е	44	0	26	0	0
2	F	44	0	26	0	0
2	G	44	0	26	0	0
2	Н	44	0	26	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	D	1	0	0	0	0
3	F	1	0	0	0	0
3	Н	1	0	0	0	0
4	A	65	0	0	0	0
4	В	62	0	0	0	0
4	С	77	0	0	0	0
4	D	80	0	0	0	0
4	Е	64	0	0	0	0
4	F	70	0	0	0	0
4	G	83	0	0	0	0
4	Н	83	0	0	1	0
All	All	21477	0	20901	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ (ext{Å})$	overlap (Å)
1:B:160:LEU:HD22	1:B:178:MET:HE3	1.55	0.87
1:D:160:LEU:HD22	1:D:178:MET:HE3	1.65	0.77
1:H:160:LEU:HD22	1:H:178:MET:HE2	1.73	0.69
1:E:66:ASP:HB3	1:E:76[B]:ARG:HH21	1.57	0.69
1:E:160:LEU:HD22	1:E:178:MET:HE3	1.79	0.64

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	337/344 (98%)	328 (97%)	8 (2%)	1 (0%)	37	34
1	В	338/344 (98%)	328 (97%)	9 (3%)	1 (0%)	37	34
1	С	335/344 (97%)	325 (97%)	9 (3%)	1 (0%)	37	34
1	D	338/344 (98%)	328 (97%)	9 (3%)	1 (0%)	37	34
1	E	338/344 (98%)	330 (98%)	7 (2%)	1 (0%)	37	34
1	F	336/344 (98%)	326 (97%)	9 (3%)	1 (0%)	37	34
1	G	341/344 (99%)	332 (97%)	8 (2%)	1 (0%)	37	34
1	Н	338/344 (98%)	328 (97%)	9 (3%)	1 (0%)	37	34
All	All	2701/2752 (98%)	2625 (97%)	68 (2%)	8 (0%)	37	34

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	242	VAL
1	В	242	VAL
1	С	242	VAL
1	D	242	VAL
1	Ε	242	VAL



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	279/286 (98%)	277 (99%)	2 (1%)	81	86
1	В	281/286 (98%)	279 (99%)	2 (1%)	81	86
1	С	280/286 (98%)	276 (99%)	4 (1%)	62	68
1	D	282/286 (99%)	280 (99%)	2 (1%)	81	86
1	E	278/286 (97%)	276 (99%)	2 (1%)	81	86
1	F	278/286 (97%)	276 (99%)	2 (1%)	81	86
1	G	284/286 (99%)	282 (99%)	2 (1%)	81	86
1	Н	281/286 (98%)	277 (99%)	4 (1%)	62	68
All	All	2243/2288 (98%)	2223 (99%)	20 (1%)	75	81

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	27	LYS
1	Н	90	GLU
1	Н	307	LYS
1	Н	178	MET
1	С	107	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	296	ASN
1	F	241	ASN
1	F	289	ASN
1	G	227	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Bond lengths			В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	D	401	-	42,48,48	1.22	5 (11%)	50,73,73	1.17	4 (8%)
2	NAD	A	401	-	42,48,48	1.37	4 (9%)	50,73,73	1.06	3 (6%)
2	NAD	F	401	-	42,48,48	1.34	4 (9%)	50,73,73	1.14	5 (10%)
2	NAD	Н	401	-	42,48,48	1.25	3 (7%)	50,73,73	1.07	3 (6%)
2	NAD	В	401	-	42,48,48	1.29	4 (9%)	50,73,73	1.06	1 (2%)
2	NAD	С	401	-	42,48,48	1.33	4 (9%)	50,73,73	1.05	3 (6%)
2	NAD	G	401	-	42,48,48	1.34	5 (11%)	50,73,73	1.10	6 (12%)
2	NAD	Е	401	-	42,48,48	1.30	4 (9%)	50,73,73	1.07	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	D	401	-	-	4/26/62/62	0/5/5/5
2	NAD	A	401	-	-	3/26/62/62	0/5/5/5
2	NAD	F	401	-	-	3/26/62/62	0/5/5/5
2	NAD	Н	401	-	-	6/26/62/62	0/5/5/5



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	В	401	-	-	8/26/62/62	0/5/5/5
2	NAD	С	401	-	-	4/26/62/62	0/5/5/5
2	NAD	G	401	-	-	4/26/62/62	0/5/5/5
2	NAD	Е	401	-	-	5/26/62/62	0/5/5/5

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\mathring{A}})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	A	401	NAD	PA-O3	5.95	1.65	1.59
2	F	401	NAD	PA-O3	5.79	1.65	1.59
2	G	401	NAD	PA-O3	5.69	1.65	1.59
2	С	401	NAD	PA-O3	5.67	1.65	1.59
2	Н	401	NAD	PA-O3	5.53	1.65	1.59

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	401	NAD	O3-PA-O1A	-3.31	100.75	110.70
2	Н	401	NAD	O3-PA-O1A	-2.85	102.13	110.70
2	D	401	NAD	O4B-C1B-N9A	-2.71	105.15	108.75
2	D	401	NAD	O2A-PA-O1A	2.64	124.72	112.44
2	F	401	NAD	O3-PA-O1A	-2.59	102.92	110.70

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAD	O4D-C1D-N1N-C2N
2	A	401	NAD	O4D-C1D-N1N-C6N
2	В	401	NAD	C5B-O5B-PA-O1A
2	В	401	NAD	C5D-O5D-PN-O2N
2	В	401	NAD	O4D-C1D-N1N-C2N

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	336/344 (97%)	0.37	9 (2%) 56 61	23, 38, 57, 82	3 (0%)
1	В	336/344 (97%)	0.43	7 (2%) 63 68	17, 39, 58, 78	4 (1%)
1	С	336/344 (97%)	0.30	4 (1%) 76 79	20, 34, 53, 81	1 (0%)
1	D	336/344 (97%)	0.30	4 (1%) 76 79	20, 35, 54, 82	4 (1%)
1	Е	336/344 (97%)	0.55	7 (2%) 63 68	16, 39, 59, 89	4 (1%)
1	F	336/344 (97%)	0.49	7 (2%) 63 68	16, 39, 57, 85	2 (0%)
1	G	336/344 (97%)	0.34	5 (1%) 71 75	18, 34, 53, 69	7 (2%)
1	Н	337/344 (97%)	0.25	5 (1%) 71 75	20, 34, 53, 96	3 (0%)
All	All	2689/2752 (97%)	0.38	48 (1%) 67 71	16, 37, 56, 96	28 (1%)

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	129	ASP	3.9
1	D	171	LEU	3.5
1	A	67	TYR	3.4
1	С	1	MET	3.3
1	В	212	LYS	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	MG	В	402	1/1	0.74	0.19	53,53,53,53	0
3	MG	A	402	1/1	0.89	0.18	57,57,57,57	0
3	MG	D	402	1/1	0.91	0.04	30,30,30,30	0
3	MG	F	402	1/1	0.94	0.05	53,53,53,53	0
2	NAD	A	401	44/44	0.95	0.08	25,32,39,41	0
2	NAD	F	401	44/44	0.95	0.07	23,32,37,43	0
2	NAD	G	401	44/44	0.96	0.06	18,26,31,35	0
2	NAD	Η	401	44/44	0.96	0.06	20,28,31,35	0
2	NAD	С	401	44/44	0.96	0.07	23,29,35,38	0
2	NAD	D	401	44/44	0.96	0.07	20,28,32,34	0
2	NAD	Ε	401	44/44	0.96	0.07	28,35,40,42	0
2	NAD	В	401	44/44	0.96	0.07	24,31,36,42	0
3	MG	Н	402	1/1	0.99	0.03	31,31,31,31	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAD A 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD F 401:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAD G 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD H 401:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAD C 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD D 401:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAD E 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD B 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

