

wwPDB X-ray Structure Validation Summary Report (i)

Apr 29, 2025 - 07:04 AM EDT

PDB ID	:	$3QA8 / pdb_{00003qa8}$
Title	:	Crystal Structure of inhibitor of kappa B kinase beta
Authors	:	Xu, G.; Lo, Y.C.; Li, Q.; Napolitano, G.; Wu, X.; Jiang, X.; Dreano, M.;
		Karin, M.; Wu, H.
Deposited on	:	2011-01-10
Resolution	:	3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#Entries)$	Similar resolution (#Entries, resolution range(Å))		
R _{free}	164625	1563 (3.70-3.50)		
Clashscore	180529	1665 (3.70-3.50)		
Ramachandran outliers	177936	1641 (3.70-3.50)		
Sidechain outliers	177891	1640 (3.70-3.50)		
RSRZ outliers	164620	1562 (3.70-3.50)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length		Quality of cha	in	
1	٨	676	%			
1	A	070	22%	45%	21%	• 8%
1	В	676	22%	43%	23%	• 8%
1	C	070	%			
1	C	676	22%	44%	22%	• 8%
1	л	676	000/	450/	010/	00/
1	D	010	22% %	45%	21%	• 8%
1	Е	676	23%	44%	21%	• 8%



Mol	Chain	Length	Quality of chain					
1	F	676	23%	43%		22%	• 8%	
1	G	676	22%	38%	18%	•	20%	
1	Н	676	20%	39%	18%	·	20%	



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 39026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms					AltConf	Trace
1	Δ	699	Total	С	Ν	0	\mathbf{S}	0	0	0
	A	022	5048	3189	899	929	31	0	0	0
1	В	699	Total	С	Ν	0	S	0	0	0
	D	022	5048	3189	899	929	31	0	0	0
1	C	699	Total	С	Ν	0	S	0	0	0
		022	5048	3189	899	929	31	0	0	0
1	П	699	Total	С	Ν	0	S	0	0	0
	D	022	5048	3189	899	929	31	0	0	0
1	F	699	Total	С	Ν	0	S	0	0	0
1		022	5048	3189	899	929	31	0	0	U
1	Б	699	Total	С	Ν	0	S	0	0	
	_ F	Г 022	5048	3189	899	929	31	0	0	0
1	C	541	Total	С	Ν	0	S	0	0	0
	G	041	4369	2764	779	800	26	0	0	0
1	ц	541	Total	С	Ν	0	S	0	0	0
	п	041	4369	2764	779	800	26	0	0	

• Molecule 1 is a protein called MGC80376 protein.

There are 152 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	0	GLY	-	expression tag	UNP Q6INT1
А	1	GLY	-	expression tag	UNP Q6INT1
А	2	GLY	-	expression tag	UNP Q6INT1
А	3	ARG	-	expression tag	UNP Q6INT1
А	4	SER	-	expression tag	UNP Q6INT1
А	5	PRO	-	expression tag	UNP Q6INT1
А	6	SER	-	expression tag	UNP Q6INT1
А	7	LEU	-	expression tag	UNP Q6INT1
А	8	PRO	-	expression tag	UNP Q6INT1
А	9	THR	-	expression tag	UNP Q6INT1
А	10	GLN	-	expression tag	UNP Q6INT1
A	11	THR	-	expression tag	UNP Q6INT1
А	12	CYS	-	expression tag	UNP Q6INT1



3Q	A8

Chain	Residue	Modelled	Actual	Comment	Reference
А	13	GLY	-	expression tag	UNP Q6INT1
А	14	PRO	-	expression tag	UNP Q6INT1
А	15	TRP	-	expression tag	UNP Q6INT1
А	16	GLU	-	expression tag	UNP Q6INT1
А	177	GLU	SER	engineered mutation	UNP Q6INT1
А	181	GLU	SER	engineered mutation	UNP Q6INT1
В	0	GLY	-	expression tag	UNP Q6INT1
В	1	GLY	-	expression tag	UNP Q6INT1
В	2	GLY	-	expression tag	UNP Q6INT1
В	3	ARG	-	expression tag	UNP Q6INT1
В	4	SER	-	expression tag	UNP Q6INT1
В	5	PRO	-	expression tag	UNP Q6INT1
В	6	SER	-	expression tag	UNP Q6INT1
В	7	LEU	-	expression tag	UNP Q6INT1
В	8	PRO	-	expression tag	UNP Q6INT1
В	9	THR	-	expression tag	UNP Q6INT1
В	10	GLN	-	expression tag	UNP Q6INT1
В	11	THR	-	expression tag	UNP Q6INT1
В	12	CYS	-	expression tag	UNP Q6INT1
В	13	GLY	-	expression tag	UNP Q6INT1
В	14	PRO	-	expression tag	UNP Q6INT1
В	15	TRP	-	expression tag	UNP Q6INT1
В	16	GLU	-	expression tag	UNP Q6INT1
В	177	GLU	SER	engineered mutation	UNP Q6INT1
В	181	GLU	SER	engineered mutation	UNP Q6INT1
С	0	GLY	-	expression tag	UNP Q6INT1
С	1	GLY	-	expression tag	UNP Q6INT1
С	2	GLY	-	expression tag	UNP Q6INT1
С	3	ARG	-	expression tag	UNP Q6INT1
С	4	SER	-	expression tag	UNP Q6INT1
С	5	PRO	-	expression tag	UNP Q6INT1
С	6	SER	-	expression tag	UNP Q6INT1
С	7	LEU	-	expression tag	UNP Q6INT1
С	8	PRO	-	expression tag	UNP Q6INT1
С	9	THR	-	expression tag	UNP Q6INT1
С	10	GLN	-	expression tag	UNP Q6INT1
С	11	THR	-	expression tag	UNP Q6INT1
С	12	CYS	-	expression tag	UNP Q6INT1
С	13	GLY	-	expression tag	UNP Q6INT1
С	14	PRO	-	expression tag	UNP Q6INT1
С	15	TRP	-	expression tag	UNP Q6INT1
C	16	GLU	-	expression tag	UNP Q6INT1



Chain	Residue	Modelled	Actual	Comment	Reference
С	177	GLU	SER	engineered mutation	UNP Q6INT1
С	181	GLU	SER	engineered mutation	UNP Q6INT1
D	0	GLY	-	expression tag	UNP Q6INT1
D	1	GLY	-	expression tag	UNP Q6INT1
D	2	GLY	-	expression tag	UNP Q6INT1
D	3	ARG	-	expression tag	UNP Q6INT1
D	4	SER	_	expression tag	UNP Q6INT1
D	5	PRO	-	expression tag	UNP Q6INT1
D	6	SER	-	expression tag	UNP Q6INT1
D	7	LEU	-	expression tag	UNP Q6INT1
D	8	PRO	-	expression tag	UNP Q6INT1
D	9	THR	-	expression tag	UNP Q6INT1
D	10	GLN	-	expression tag	UNP Q6INT1
D	11	THR	-	expression tag	UNP Q6INT1
D	12	CYS	-	expression tag	UNP Q6INT1
D	13	GLY	-	expression tag	UNP Q6INT1
D	14	PRO	-	expression tag	UNP Q6INT1
D	15	TRP	-	expression tag	UNP Q6INT1
D	16	GLU	-	expression tag	UNP Q6INT1
D	177	GLU	SER	engineered mutation	UNP Q6INT1
D	181	GLU	SER	engineered mutation	UNP Q6INT1
Е	0	GLY	-	expression tag	UNP Q6INT1
Е	1	GLY	-	expression tag	UNP Q6INT1
E	2	GLY	-	expression tag	UNP Q6INT1
Е	3	ARG	-	expression tag	UNP Q6INT1
Е	4	SER	-	expression tag	UNP Q6INT1
Е	5	PRO	-	expression tag	UNP Q6INT1
Е	6	SER	-	expression tag	UNP Q6INT1
E	7	LEU	-	expression tag	UNP Q6INT1
E	8	PRO	-	expression tag	UNP Q6INT1
E	9	THR	-	expression tag	UNP Q6INT1
E	10	GLN	-	expression tag	UNP Q6INT1
E	11	THR	-	expression tag	UNP Q6INT1
E	12	CYS	-	expression tag	UNP Q6INT1
E	13	GLY	-	expression tag	UNP Q6INT1
E	14	PRO	-	expression tag	UNP Q6INT1
E	15	TRP	-	expression tag	UNP Q6INT1
E	16	GLU	-	expression tag	UNP Q6INT1
E	177	GLU	SER	engineered mutation	UNP Q6INT1
Е	181	GLU	SER	engineered mutation	UNP $Q6INT1$
F	0	GLY	-	expression tag	UNP Q6INT1
F	1	GLY	-	expression tag	UNP Q6INT1



3Q	A8
0.00	

Chain	Residue	Modelled	Actual	Comment	Reference
F	2	GLY	-	expression tag	UNP Q6INT1
F	3	ARG	-	expression tag	UNP Q6INT1
F	4	SER	-	expression tag	UNP Q6INT1
F	5	PRO	-	expression tag	UNP Q6INT1
F	6	SER	_	expression tag	UNP Q6INT1
F	7	LEU	-	expression tag	UNP Q6INT1
F	8	PRO	-	expression tag	UNP Q6INT1
F	9	THR	-	expression tag	UNP Q6INT1
F	10	GLN	-	expression tag	UNP Q6INT1
F	11	THR	-	expression tag	UNP Q6INT1
F	12	CYS	-	expression tag	UNP Q6INT1
F	13	GLY	-	expression tag	UNP Q6INT1
F	14	PRO	-	expression tag	UNP Q6INT1
F	15	TRP	-	expression tag	UNP Q6INT1
F	16	GLU	-	expression tag	UNP Q6INT1
F	177	GLU	SER	engineered mutation	UNP Q6INT1
F	181	GLU	SER	engineered mutation	UNP Q6INT1
G	0	GLY	-	expression tag	UNP Q6INT1
G	1	GLY	-	expression tag	UNP Q6INT1
G	2	GLY	-	expression tag	UNP Q6INT1
G	3	ARG	-	expression tag	UNP Q6INT1
G	4	SER	-	expression tag	UNP Q6INT1
G	5	PRO	-	expression tag	UNP Q6INT1
G	6	SER	-	expression tag	UNP Q6INT1
G	7	LEU	-	expression tag	UNP Q6INT1
G	8	PRO	-	expression tag	UNP Q6INT1
G	9	THR	-	expression tag	UNP Q6INT1
G	10	GLN	-	expression tag	UNP Q6INT1
G	11	THR	-	expression tag	UNP Q6INT1
G	12	CYS	-	expression tag	UNP Q6INT1
G	13	GLY	-	expression tag	UNP Q6INT1
G	14	PRO	-	expression tag	UNP Q6INT1
G	15	TRP	-	expression tag	UNP Q6INT1
G	16	GLU	-	expression tag	UNP Q6INT1
G	177	GLU	SER	engineered mutation	UNP Q6INT1
G	181	GLU	SER	engineered mutation	UNP Q6INT1
H	0	GLY	-	expression tag	UNP Q6INT1
H	1	GLY	-	expression tag	UNP Q6INT1
H	2	GLY	-	expression tag	UNP Q6INT1
H	3	ARG	-	expression tag	UNP Q6INT1
H	4	SER	-	expression tag	UNP Q6INT1
H	5	PRO	-	expression tag	UNP Q6INT1



3QA8

Chain	Residue	Modelled	Actual	Comment	Reference
Н	6	SER	-	expression tag	UNP Q6INT1
Н	7	LEU	-	expression tag	UNP Q6INT1
Н	8	PRO	-	expression tag	UNP Q6INT1
Н	9	THR	-	expression tag	UNP Q6INT1
Н	10	GLN	-	expression tag	UNP Q6INT1
Н	11	THR	-	expression tag	UNP Q6INT1
Н	12	CYS	-	expression tag	UNP Q6INT1
Н	13	GLY	-	expression tag	UNP Q6INT1
Н	14	PRO	-	expression tag	UNP Q6INT1
Н	15	TRP	-	expression tag	UNP Q6INT1
Н	16	GLU	-	expression tag	UNP Q6INT1
H	177	GLU	SER	engineered mutation	UNP Q6INT1
Н	181	GLU	SER	engineered mutation	UNP Q6INT1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: MGC80376 protein





B









NS88 NS88 N588 N588 N588 N589 N589 N589 N581 N589 N581 N589 N581 N589 N581 N591 N581 N591 N582 N591 1595 1595 1595 1596 1596 1607 1607 1607 1607 1607 1607 1601 1607 1611 1607 1611 1607 1613 1607 1614 1607 1613 1607 1614 1607 1614 1613 1614 1614 1614 1623 1653 1633 1644 1644 1644 1644 1644 1653 1653 1644 1644 1654 1653 1654</t

W655 W656 L658 L658 K659 A660 A660 A660 A660 A660 A660 Cly Cly Cly SER SER SER

 \bullet Molecule 1: MGC80376 protein



 \bullet Molecule 1: MGC80376 protein



3144 0145 1146













4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	103.17Å 140.34Å 161.17Å	Demositer
a, b, c, α , β , γ	71.28° 79.56° 86.04°	Depositor
$\mathbf{P}_{\mathrm{oscolution}}(\mathbf{\hat{A}})$	15.00 - 3.60	Depositor
Resolution (A)	15.00 - 3.60	EDS
% Data completeness	78.7 (15.00-3.60)	Depositor
(in resolution range)	78.0 (15.00-3.60)	EDS
R_{merge}	0.07	Depositor
R _{sym}	0.04	Depositor
$< I/\sigma(I) > 1$	$1.18 (at 3.48 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
D D.	0.308 , 0.344	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.302 , 0.337	DCC
R_{free} test set	4480 reflections (5.06%)	wwPDB-VP
Wilson B-factor $(Å^2)$	84.5	Xtriage
Anisotropy	0.749	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.19, 107.0	EDS
L-test for $twinning^2$	$ L > = 0.47, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	39026	wwPDB-VP
Average B, all atoms $(Å^2)$	199.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	ond lengths	Bond angles	
	Ullaili	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.96	9/5136~(0.2%)	1.33	50/6931~(0.7%)
1	В	0.98	12/5136~(0.2%)	1.34	53/6931~(0.8%)
1	С	1.02	10/5136~(0.2%)	1.34	59/6931~(0.9%)
1	D	0.96	8/5136~(0.2%)	1.32	50/6931~(0.7%)
1	Е	1.00	9/5136~(0.2%)	1.33	48/6931~(0.7%)
1	F	0.99	10/5136~(0.2%)	1.33	47/6931~(0.7%)
1	G	0.96	7/4448~(0.2%)	1.32	42/6012~(0.7%)
1	Н	0.98	8/4448~(0.2%)	1.33	43/6012~(0.7%)
All	All	0.98	73/39712~(0.2%)	1.33	392/53610~(0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	11
1	В	0	11
1	С	0	10
1	D	0	11
1	Ε	0	10
1	F	0	11
1	G	0	11
1	Н	0	11
All	All	0	86

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	Е	261	THR	CA-C	10.16	1.65	1.52
1	С	261	THR	CA-C	10.05	1.65	1.52
1	G	261	THR	CA-C	9.20	1.64	1.52
1	А	261	THR	CA-C	9.04	1.64	1.52



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	F	119	GLU	N-CA	-8.27	1.38	1.46

The worst 5 of 392 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	F	223	LEU	CA-C-N	-12.93	107.72	120.52
1	F	223	LEU	C-N-CA	-12.93	107.72	120.52
1	А	223	LEU	CA-C-N	-11.84	108.06	120.85
1	А	223	LEU	C-N-CA	-11.84	108.06	120.85
1	F	426	LEU	N-CA-C	-11.84	98.93	113.50

There are no chirality outliers.

5 of 86 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	119	GLU	Peptide
1	А	120	GLY	Peptide
1	А	169	TYR	Peptide
1	А	221	PRO	Peptide
1	А	223	LEU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5048	0	5120	804	3
1	В	5048	0	5120	849	2
1	С	5048	0	5120	870	4
1	D	5048	0	5120	888	0
1	Е	5048	0	5120	849	2
1	F	5048	0	5120	843	0
1	G	4369	0	4430	634	0
1	Н	4369	0	4430	650	1
All	All	39026	0	39580	6134	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 78.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:547:LEU:CD1	1:E:615:THR:HG22	1.18	1.61
1:E:547:LEU:HD13	1:E:615:THR:CG2	1.16	1.54
1:C:496:LYS:CB	1:D:655:TRP:HE1	1.25	1.48
1:A:524:CYS:SG	1:A:643:VAL:HG11	1.55	1.46
1:C:496:LYS:HB2	1:D:655:TRP:NE1	1.23	1.43

The worst 5 of 6134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:501:MET:SD	1:E:364:ALA:O[1_565]	1.61	0.59
1:E:515:ARG:NH1	1:H:394:LYS:NZ[1_465]	1.76	0.44
1:B:617:VAL:CG2	1:C:519:GLN:OE1[1_465]	1.96	0.24
1:A:522:GLU:OE1	1:C:295:PRO:CB[1_565]	2.01	0.19
1:A:522:GLU:OE1	1:C:295:PRO:CG[1_565]	2.04	0.16

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	А	610/676~(90%)	360 (59%)	145 (24%)	105 (17%)	0	2
1	В	610/676~(90%)	362~(59%)	143 (23%)	105 (17%)	0	2
1	C	610/676~(90%)	359~(59%)	147 (24%)	104 (17%)	0	2
1	D	610/676~(90%)	363 (60%)	143 (23%)	104 (17%)	0	2
1	Е	610/676~(90%)	361 (59%)	147 (24%)	102 (17%)	0	2
1	F	610/676~(90%)	362 (59%)	144 (24%)	104 (17%)	0	2
1	G	527/676~(78%)	310 (59%)	126(24%)	91 (17%)	0	2
1	Н	527/676~(78%)	313 (59%)	121 (23%)	93 (18%)	0	1



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	4714/5408 (87%)	2790~(59%)	1116 (24%)	808 (17%)	0 2

5 of 808 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	101	GLY
1	А	106	LYS
1	А	110	GLN
1	А	111	PHE
1	А	166	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	\mathbf{P}	erc	entiles
1	А	563/609~(92%)	498 (88%)	65~(12%)		4	23
1	В	563/609~(92%)	497~(88%)	66 (12%)		4	23
1	С	563/609~(92%)	496 (88%)	67 (12%)		4	22
1	D	563/609~(92%)	498 (88%)	65~(12%)		4	23
1	Ε	563/609~(92%)	500~(89%)	63 (11%)		5	24
1	F	563/609~(92%)	498 (88%)	65~(12%)		4	23
1	G	488/609~(80%)	437 (90%)	51 (10%)		5	26
1	Н	488/609~(80%)	436 (89%)	52 (11%)		5	26
All	All	4354/4872 (89%)	3860 (89%)	494 (11%)		4	24

5 of 494 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	D	416	ASP
1	Н	51	SER
1	Е	314	VAL
1	G	606	ILE
1	Н	390	PHE



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 153 such sidechains are listed below:

Mol	Chain	\mathbf{Res}	Type
1	F	365	GLN
1	Н	230	GLN
1	F	548	GLN
1	G	301	GLN
1	Н	449	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RS	SRZ>	>2	$OWAB(A^2)$	Q<0.9
1	А	622/676~(92%)	-0.43	4 (0%)	85	69	76, 189, 275, 337	0
1	В	622/676~(92%)	-0.40	1 (0%)	92	84	71, 191, 274, 336	0
1	С	622/676~(92%)	-0.16	7 (1%)	77	57	58, 188, 275, 327	0
1	D	622/676~(92%)	-0.53	2(0%)	90	79	85, 194, 278, 327	0
1	Е	622/676~(92%)	-0.28	4 (0%)	85	69	74, 190, 274, 330	0
1	F	622/676~(92%)	-0.52	1 (0%)	92	84	93, 200, 283, 331	0
1	G	541/676~(80%)	-0.51	1 (0%)	92	84	96, 214, 302, 387	0
1	Н	541/676~(80%)	-0.46	1 (0%)	92	84	79, 194, 293, 423	0
All	All	4814/5408 (89%)	-0.41	21 (0%)	89	76	58, 195, 283, 423	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	544	SER	4.0
1	С	508	GLU	4.0
1	С	636	MET	3.7
1	С	648	GLU	3.5
1	С	342	GLN	3.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

