



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 10:42 AM UTC

PDB ID : 7QJK / pdb_00007qjk
Title : Crystal structure of PDE6D in complex with Compound-2
Authors : Yelland, T.; Ismail, S.
Deposited on : 2021-12-16
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	151	
1	BBB	151	
1	CCC	151	
1	DDD	151	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	BBB	145	1178	756	198	218	6	0	0	0
1	AAA	148	1202	772	201	222	7	0	0	0
1	CCC	146	1180	757	194	222	7	0	1	0
1	DDD	141	1134	730	186	212	6	0	1	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
BBB	0	ALA	-	expression tag	UNP O43924
AAA	0	ALA	-	expression tag	UNP O43924
CCC	0	ALA	-	expression tag	UNP O43924
DDD	0	ALA	-	expression tag	UNP O43924

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	BBB	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is N-(phenylmethyl)pyridin-2-amine (CCD ID: 9VO) (formula: $C_{12}H_{12}N_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	BBB	1	Total	C	N	0	0
			14	12	2		
3	AAA	1	Total	C	N	0	0
			14	12	2		
3	AAA	1	Total	C	N	0	0
			14	12	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	CCC	1	Total	C	N	0	0
			14	12	2		
3	DDD	1	Total	C	N	0	0
			14	12	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	BBB	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	BBB	7	Total	O	0	0
			7	7		
5	AAA	5	Total	O	0	0
			5	5		
5	CCC	10	Total	O	0	0
			10	10		
5	DDD	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta




- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta



- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta



- Molecule 1: Retinal rod rhodopsin-sensitive cGMP 3',5'-cyclic phosphodiesterase subunit delta



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	76.55Å 76.55Å 238.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.46 – 3.10 64.46 – 3.10	Depositor EDS
% Data completeness (in resolution range)	88.8 (64.46-3.10) 88.8 (64.46-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 3.13Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.246 , 0.283 0.245 , 0.276	Depositor DCC
R_{free} test set	625 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	69.3	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 66.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4799	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 9VO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.99	0/1226	1.21	0/1650
1	BBB	0.97	0/1202	1.20	0/1620
1	CCC	0.99	0/1204	1.21	0/1625
1	DDD	1.00	0/1157	1.16	0/1563
All	All	0.99	0/4789	1.19	0/6458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1202	0	1197	33	0
1	BBB	1178	0	1161	16	0
1	CCC	1180	0	1144	23	0
1	DDD	1134	0	1098	48	0
2	BBB	4	0	6	0	0
3	AAA	28	0	0	1	0
3	BBB	14	0	0	0	0
3	CCC	14	0	0	0	0
3	DDD	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	BBB	5	0	0	0	0
5	AAA	5	0	0	1	0
5	BBB	7	0	0	0	0
5	CCC	10	0	0	0	0
5	DDD	4	0	0	0	0
All	All	4799	0	4606	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:118:MET:HB2	1:AAA:119:PRO:HD2	1.45	0.96
1:AAA:119:PRO:HG2	1:AAA:122:VAL:HG23	1.55	0.86
1:DDD:75:ARG:HB2	1:DDD:92:PHE:O	1.83	0.79
1:DDD:11:ILE:HD12	1:DDD:69:GLU:HG3	1.65	0.76
1:CCC:62:GLU:HG2	1:CCC:106:GLN:HG3	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	144/151 (95%)	135 (94%)	8 (6%)	1 (1%)	18	49
1	BBB	141/151 (93%)	136 (96%)	5 (4%)	0	100	100
1	CCC	143/151 (95%)	136 (95%)	4 (3%)	3 (2%)	5	25
1	DDD	138/151 (91%)	128 (93%)	6 (4%)	4 (3%)	3	19
All	All	566/604 (94%)	535 (94%)	23 (4%)	8 (1%)	9	34

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	CCC	3	ALA
1	DDD	8	ALA
1	CCC	2	SER
1	AAA	35	THR
1	DDD	6	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	131/135 (97%)	124 (95%)	7 (5%)	20	51
1	BBB	128/135 (95%)	120 (94%)	8 (6%)	16	45
1	CCC	128/135 (95%)	126 (98%)	2 (2%)	55	75
1	DDD	122/135 (90%)	114 (93%)	8 (7%)	15	43
All	All	509/540 (94%)	484 (95%)	25 (5%)	23	53

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	146	ARG
1	DDD	6	GLU
1	DDD	117	MET
1	CCC	118	MET
1	DDD	11	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	BBB	201	-	3,3,3	0.06	0	2,2,2	0.10	0
3	9VO	BBB	202	-	15,15,15	1.67	2 (13%)	18,18,18	1.10	2 (11%)
4	SO4	BBB	203	-	4,4,4	0.35	0	6,6,6	0.07	0
3	9VO	DDD	201	-	15,15,15	1.68	2 (13%)	18,18,18	1.12	2 (11%)
3	9VO	AAA	201	-	15,15,15	1.72	3 (20%)	18,18,18	1.16	2 (11%)
3	9VO	AAA	202	-	15,15,15	1.79	3 (20%)	18,18,18	1.16	3 (16%)
3	9VO	CCC	201	-	15,15,15	1.82	3 (20%)	18,18,18	1.13	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	BBB	201	-	-	0/1/1/1	-
3	9VO	BBB	202	-	-	3/5/5/5	0/2/2/2
3	9VO	DDD	201	-	-	0/5/5/5	0/2/2/2
3	9VO	AAA	201	-	-	2/5/5/5	0/2/2/2
3	9VO	AAA	202	-	-	0/5/5/5	0/2/2/2
3	9VO	CCC	201	-	-	2/5/5/5	0/2/2/2

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	CCC	201	9VO	C8-C9	-5.88	1.38	1.51
3	AAA	202	9VO	C8-C9	-5.63	1.39	1.51
3	DDD	201	9VO	C8-C9	-5.52	1.39	1.51
3	AAA	201	9VO	C8-C9	-5.51	1.39	1.51
3	BBB	202	9VO	C8-C9	-5.21	1.39	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	DDD	201	9VO	C2-N6-C5	2.56	120.87	117.21
3	AAA	201	9VO	N7-C5-N6	2.35	120.53	116.98
3	DDD	201	9VO	C1-C2-N6	-2.32	119.75	123.42
3	AAA	202	9VO	N7-C5-N6	2.25	120.38	116.98
3	AAA	202	9VO	C1-C2-N6	-2.22	119.91	123.42

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BBB	202	9VO	C4-C5-N7-C8
3	BBB	202	9VO	N6-C5-N7-C8
3	AAA	201	9VO	C4-C5-N7-C8
3	AAA	201	9VO	N6-C5-N7-C8
3	CCC	201	9VO	C4-C5-N7-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	202	9VO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	148/151 (98%)	-0.04	0 100 100	54, 81, 109, 133	0
1	BBB	145/151 (96%)	-0.21	1 (0%) 84 68	44, 67, 100, 120	0
1	CCC	146/151 (96%)	-0.03	0 100 100	48, 80, 126, 152	1 (0%)
1	DDD	141/151 (93%)	0.18	1 (0%) 84 68	36, 98, 147, 166	1 (0%)
All	All	580/604 (96%)	-0.03	2 (0%) 90 80	36, 80, 128, 166	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	0	ALA	3.0
1	DDD	67	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	BBB	201	4/4	0.61	0.19	83,88,89,89	0
3	9VO	CCC	201	14/14	0.89	0.15	60,72,82,83	0
3	9VO	AAA	202	14/14	0.91	0.17	81,88,103,105	0
3	9VO	BBB	202	14/14	0.91	0.16	67,72,75,75	0
3	9VO	AAA	201	14/14	0.92	0.14	66,72,81,82	0
3	9VO	DDD	201	14/14	0.93	0.09	74,78,80,81	0
4	SO4	BBB	203	5/5	0.95	0.06	76,79,86,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.