



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 21, 2025 – 01:29 pm BST

PDB ID : 9QRH / pdb\_00009qrh  
Title : Protein Kinase CK2 and bivalent inhibitors  
Authors : Krimm, I.; Gelin, M.; Guichou, J.F.; Grenier, D.  
Deposited on : 2025-04-03  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	 4% 79% 12% 8%
1	B	359	 3% 76% 16% 8%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	0	0
			2736	1756	476	493	11			
1	B	331	Total	C	N	O	S	0	0	0
			2730	1753	475	491	11			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP P68400
A	-20	HIS	-	expression tag	UNP P68400
A	-19	HIS	-	expression tag	UNP P68400
A	-18	HIS	-	expression tag	UNP P68400
A	-17	HIS	-	expression tag	UNP P68400
A	-16	HIS	-	expression tag	UNP P68400
A	-15	HIS	-	expression tag	UNP P68400
A	-14	SER	-	expression tag	UNP P68400
A	-13	SER	-	expression tag	UNP P68400
A	-12	GLY	-	expression tag	UNP P68400
A	-11	VAL	-	expression tag	UNP P68400
A	-10	ASP	-	expression tag	UNP P68400
A	-9	LEU	-	expression tag	UNP P68400
A	-8	GLY	-	expression tag	UNP P68400
A	-7	THR	-	expression tag	UNP P68400
A	-6	GLU	-	expression tag	UNP P68400
A	-5	ASN	-	expression tag	UNP P68400
A	-4	LEU	-	expression tag	UNP P68400
A	-3	TYR	-	expression tag	UNP P68400
A	-2	PHE	-	expression tag	UNP P68400
A	-1	GLN	-	expression tag	UNP P68400
A	0	SER	-	expression tag	UNP P68400
B	-21	MET	-	initiating methionine	UNP P68400
B	-20	HIS	-	expression tag	UNP P68400
B	-19	HIS	-	expression tag	UNP P68400

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	HIS	-	expression tag	UNP P68400
B	-17	HIS	-	expression tag	UNP P68400
B	-16	HIS	-	expression tag	UNP P68400
B	-15	HIS	-	expression tag	UNP P68400
B	-14	SER	-	expression tag	UNP P68400
B	-13	SER	-	expression tag	UNP P68400
B	-12	GLY	-	expression tag	UNP P68400
B	-11	VAL	-	expression tag	UNP P68400
B	-10	ASP	-	expression tag	UNP P68400
B	-9	LEU	-	expression tag	UNP P68400
B	-8	GLY	-	expression tag	UNP P68400
B	-7	THR	-	expression tag	UNP P68400
B	-6	GLU	-	expression tag	UNP P68400
B	-5	ASN	-	expression tag	UNP P68400
B	-4	LEU	-	expression tag	UNP P68400
B	-3	TYR	-	expression tag	UNP P68400
B	-2	PHE	-	expression tag	UNP P68400
B	-1	GLN	-	expression tag	UNP P68400
B	0	SER	-	expression tag	UNP P68400

- Molecule 2 is 5-[[[3 {E}]-3-[4-[2-(3,4-dichlorophenyl)ethylamino]-4-oxidanylidene-butanoyl]iminopropyl]amino]benzo[c][2,6]naphthyridine-8-carboxylic acid (CCD ID: A1I9M) (formula: C<sub>28</sub>H<sub>25</sub>Cl<sub>2</sub>N<sub>5</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0
			39	28	2	5	4	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	
			39	28	2	5	4	
							0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S		
			5	4	1	0	0
3	A	1	Total	O	S		
			5	4	1	0	0
3	A	1	Total	O	S		
			5	4	1	0	0
3	A	1	Total	O	S		
			5	4	1	0	0
3	B	1	Total	O	S		
			5	4	1	0	0
3	B	1	Total	O	S		
			5	4	1	0	0
3	B	1	Total	O	S		
			5	4	1	0	0
3	B	1	Total	O	S		
			5	4	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	15	Total 15	O 15	0	0
4	B	5	Total 5	O 5	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Casein kinase II subunit alpha



#### • Molecule 1: Casein kinase II subunit alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.67Å 127.67Å 124.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.14 – 2.59 45.14 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.14-2.59) 100.0 (45.14-2.59)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.15 (at 2.58Å)	Xtriage
Refinement program	PHENIX (1.19_4092: ???)	Depositor
R, $R_{free}$	0.193 , 0.236 0.193 , 0.232	Depositor DCC
$R_{free}$ test set	1607 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	78.1	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 36.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.002 for -h,-l,-k 0.001 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5609	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.2280e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, A1I9M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2809	0.54	0/3805
1	B	0.37	0/2803	0.56	0/3799
All	All	0.38	0/5612	0.55	0/7604

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2736	0	2650	28	0
1	B	2730	0	2639	28	0
2	A	39	0	0	1	0
2	B	39	0	0	1	0
3	A	20	0	0	1	0
3	B	25	0	0	0	0
4	A	15	0	0	0	0
4	B	5	0	0	0	0
All	All	5609	0	5289	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:157:VAL:HG12	1:B:221:MET:HE1	1.59	0.84
1:A:121:PHE:HA	1:A:124:LEU:HB3	1.75	0.68
1:A:36:GLN:O	1:A:38:ASP:N	2.26	0.67
1:B:135:PHE:HA	1:B:327:VAL:HG21	1.78	0.65
1:A:316:ARG:CZ	1:A:319:MET:HE1	2.28	0.64
1:B:137:MET:HE2	2:B:403:A1I9M:CL17	2.38	0.60
1:B:165:ASP:HB3	1:B:170:LYS:HB3	1.88	0.55
1:A:134:ARG:HG2	1:A:323:TYR:CZ	2.42	0.55
1:A:195:ARG:HD2	1:A:237:ASP:C	2.32	0.54
1:A:279:LYS:NZ	3:A:405:SO4:O4	2.40	0.54
1:A:329:LYS:HD3	1:A:330:ASP:N	2.23	0.53
1:B:123:GLN:O	1:B:127:THR:HG22	2.09	0.52
1:B:105:VAL:HG13	1:B:106:SER:H	1.73	0.52
1:B:267:PRO:HA	1:B:270:ASN:OD1	2.10	0.52
1:B:212:SER:OG	1:B:309:HIS:HB2	2.10	0.52
1:A:88:LEU:HD21	1:A:152:ILE:HD12	1.91	0.52
1:A:316:ARG:NH2	1:A:319:MET:HE1	2.26	0.51
1:B:31:VAL:HG21	1:B:33:TRP:CZ2	2.47	0.50
1:A:329:LYS:HD3	1:A:329:LYS:C	2.35	0.50
1:A:136:TYR:O	1:A:140:ILE:HG13	2.11	0.49
1:B:134:ARG:HG2	1:B:323:TYR:CZ	2.48	0.49
1:B:322:PRO:HA	1:B:325:TYR:CD1	2.48	0.48
1:A:33:TRP:CH2	1:A:109:PRO:HB3	2.48	0.48
1:A:95:ILE:HB	1:A:174:ILE:HG22	1.95	0.48
1:A:36:GLN:C	1:A:38:ASP:N	2.73	0.47
1:B:133:ILE:HD13	1:B:225:MET:HB3	1.97	0.47
1:A:316:ARG:CZ	1:A:319:MET:CE	2.94	0.46
1:B:91:GLY:HA3	1:B:146:TYR:CE2	2.49	0.46
1:A:265:LEU:HD12	1:A:265:LEU:HA	1.84	0.46
1:B:261:TYR:OH	1:B:310:GLN:HG3	2.15	0.46
1:A:243:VAL:HG21	1:A:272:ILE:HG13	1.98	0.46
1:A:267:PRO:HA	1:A:270:ASN:OD1	2.16	0.45
1:B:215:MET:HG3	1:B:313:LEU:O	2.17	0.45
1:B:258:ILE:HG23	1:B:263:ILE:HB	1.97	0.45
1:B:196:TYR:CD2	1:B:231:PRO:HG3	2.52	0.44
1:A:161:ASN:O	1:A:173:LEU:HD12	2.18	0.44
1:B:286:HIS:CE1	1:B:288:GLU:HB2	2.53	0.44
1:B:161:ASN:O	1:B:173:LEU:HD12	2.17	0.44
1:A:128:LEU:HD12	1:A:128:LEU:HA	1.75	0.43
1:B:19:ARG:HD3	1:B:23:TYR:CD2	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ARG:NH2	1:A:25:ASP:OD2	2.32	0.43
1:A:163:MET:HB3	1:A:163:MET:HE3	1.59	0.43
1:B:15:VAL:O	1:B:19:ARG:HG3	2.18	0.43
1:B:265:LEU:HD12	1:B:265:LEU:HA	1.81	0.43
1:B:108:THR:HA	1:B:109:PRO:HD3	1.88	0.43
1:A:225:MET:HE1	2:A:401:A1I9M:CL16	2.55	0.43
1:A:86:GLU:HG2	1:A:89:ARG:HH12	1.83	0.42
1:B:254:LEU:O	1:B:258:ILE:HG12	2.19	0.42
1:A:305:LEU:HD12	1:A:305:LEU:HA	1.90	0.42
1:A:91:GLY:HA3	1:A:146:TYR:CE2	2.55	0.41
1:A:101:VAL:HG23	1:A:110:ALA:HB3	2.02	0.41
1:A:260:LYS:HD3	1:A:261:TYR:CE2	2.54	0.41
1:B:70:LEU:HD13	1:B:78:ILE:HG12	2.02	0.41
1:B:286:HIS:HE1	1:B:288:GLU:HB2	1.86	0.41
1:B:5:VAL:HB	1:B:261:TYR:HA	2.02	0.40
1:B:138:TYR:CE1	1:B:319:MET:HG2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/359 (91%)	306 (93%)	18 (6%)	4 (1%)	11	24
1	B	329/359 (92%)	313 (95%)	14 (4%)	2 (1%)	22	43
All	All	657/718 (92%)	619 (94%)	32 (5%)	6 (1%)	14	31

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	120	ASP

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	120	ASP
1	B	175	ASP
1	A	175	ASP
1	A	272	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/329 (89%)	289 (99%)	3 (1%)	73	88
1	B	290/329 (88%)	282 (97%)	8 (3%)	38	65
All	All	582/658 (88%)	571 (98%)	11 (2%)	52	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	VAL
1	A	102	LYS
1	A	305	LEU
1	B	49	LYS
1	B	65	VAL
1	B	66	VAL
1	B	71	LYS
1	B	73	VAL
1	B	105	VAL
1	B	293	VAL
1	B	330	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	207	GLN
1	B	29	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	36	GLN
1	B	160	HIS
1	B	331	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1I9M	A	401	-	40,42,42	2.14	15 (37%)	49,57,57	1.71	8 (16%)
3	SO4	A	404	-	4,4,4	0.25	0	6,6,6	0.28	0
2	A1I9M	B	403	-	40,42,42	2.23	15 (37%)	49,57,57	1.59	6 (12%)
3	SO4	B	402	-	4,4,4	0.20	0	6,6,6	0.48	0
3	SO4	A	402	-	4,4,4	0.26	0	6,6,6	0.27	0
3	SO4	B	405	-	4,4,4	0.18	0	6,6,6	0.25	0
3	SO4	A	403	-	4,4,4	0.29	0	6,6,6	0.17	0
3	SO4	A	405	-	4,4,4	0.25	0	6,6,6	0.13	0
3	SO4	B	401	-	4,4,4	0.28	0	6,6,6	0.24	0
3	SO4	B	404	-	4,4,4	0.15	0	6,6,6	0.30	0
3	SO4	B	406	-	4,4,4	0.31	0	6,6,6	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1I9M	A	401	-	-	8/20/23/23	0/4/4/4
2	A1I9M	B	403	-	-	7/20/23/23	0/4/4/4

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	A1I9M	C19-N18	5.11	1.45	1.26
2	B	403	A1I9M	C19-N18	5.07	1.45	1.26
2	B	403	A1I9M	C27-C37	4.66	1.59	1.49
2	A	401	A1I9M	C23-N22	4.54	1.42	1.34
2	B	403	A1I9M	C33-C34	4.38	1.42	1.36
2	B	403	A1I9M	C23-N22	4.30	1.42	1.34
2	A	401	A1I9M	C33-C34	4.23	1.42	1.36
2	B	403	A1I9M	C25-N24	3.63	1.43	1.37
2	B	403	A1I9M	C23-N24	3.60	1.37	1.32
2	B	403	A1I9M	C21-C20	3.46	1.56	1.53
2	B	403	A1I9M	C05-N07	3.39	1.41	1.33
2	A	401	A1I9M	C23-N24	3.27	1.37	1.32
2	A	401	A1I9M	C20-C19	3.16	1.52	1.49
2	A	401	A1I9M	C05-N07	2.97	1.40	1.33
2	A	401	A1I9M	C27-C37	2.91	1.55	1.49
2	B	403	A1I9M	C29-C28	2.64	1.42	1.36
2	A	401	A1I9M	C02-N18	2.64	1.44	1.35
2	A	401	A1I9M	C29-C30	2.60	1.46	1.41
2	A	401	A1I9M	O01-C02	2.53	1.25	1.22
2	B	403	A1I9M	C02-N18	2.52	1.44	1.35
2	A	401	A1I9M	C21-C20	2.47	1.55	1.53
2	B	403	A1I9M	C36-N35	2.43	1.36	1.32
2	B	403	A1I9M	C20-C19	2.38	1.51	1.49
2	B	403	A1I9M	O01-C02	2.34	1.25	1.22
2	A	401	A1I9M	C29-C28	2.32	1.41	1.36
2	A	401	A1I9M	C36-N35	2.31	1.36	1.32
2	B	403	A1I9M	O38-C37	2.30	1.29	1.22
2	A	401	A1I9M	O38-C37	2.28	1.29	1.22
2	B	403	A1I9M	C29-C30	2.27	1.45	1.41
2	A	401	A1I9M	C08-N07	-2.13	1.41	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1I9M	C04-C05-N07	5.43	125.56	116.42
2	B	403	A1I9M	C04-C05-N07	5.28	125.30	116.42
2	B	403	A1I9M	C03-C04-C05	5.27	121.79	112.56
2	A	401	A1I9M	C03-C04-C05	4.78	120.94	112.56
2	A	401	A1I9M	O06-C05-N07	-3.93	115.59	123.01
2	B	403	A1I9M	O06-C05-N07	-3.71	116.01	123.01
2	A	401	A1I9M	C04-C03-C02	-2.86	109.43	112.24
2	A	401	A1I9M	O39-C37-C27	2.80	122.12	114.85
2	B	403	A1I9M	O39-C37-C27	2.55	121.46	114.85
2	A	401	A1I9M	C23-N24-C25	2.47	121.57	116.67
2	A	401	A1I9M	C34-C33-C32	-2.42	117.77	119.81
2	B	403	A1I9M	C34-C33-C32	-2.41	117.77	119.81
2	A	401	A1I9M	C31-C36-N35	-2.32	120.95	124.35
2	B	403	A1I9M	C23-N24-C25	2.17	120.96	116.67

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	A1I9M	C19-C20-C21-N22
2	A	401	A1I9M	N07-C08-C09-C10
2	B	403	A1I9M	C19-C20-C21-N22
2	B	403	A1I9M	N07-C08-C09-C10
2	B	403	A1I9M	C02-C03-C04-C05
2	A	401	A1I9M	C08-C09-C10-C11
2	B	403	A1I9M	C08-C09-C10-C11
2	A	401	A1I9M	C08-C09-C10-C15
2	B	403	A1I9M	C08-C09-C10-C15
2	A	401	A1I9M	C26-C27-C37-O38
2	A	401	A1I9M	C28-C27-C37-O38
2	B	403	A1I9M	C26-C27-C37-O38
2	A	401	A1I9M	C26-C27-C37-O39
2	A	401	A1I9M	N18-C19-C20-C21
2	B	403	A1I9M	C03-C04-C05-O06

There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	A1I9M	1	0
2	B	403	A1I9M	1	0
3	A	405	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	330/359 (91%)	0.14	14 (4%) 41 35	56, 74, 114, 164	0
1	B	331/359 (92%)	0.02	9 (2%) 56 50	55, 71, 105, 144	2 (0%)
All	All	661/718 (92%)	0.08	23 (3%) 47 41	55, 73, 111, 164	2 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	ARG	3.2
1	B	120	ASP	3.1
1	B	118	ASN	3.1
1	A	121	PHE	3.0
1	B	105	VAL	3.0
1	A	37	ASP	3.0
1	A	268	ARG	2.7
1	A	75	LYS	2.7
1	B	49	LYS	2.7
1	B	121	PHE	2.7
1	A	127	THR	2.6
1	A	196	TYR	2.4
1	B	125	TYR	2.3
1	A	124	LEU	2.3
1	B	332	ALA	2.3
1	A	74	LYS	2.2
1	B	32	GLU	2.1
1	A	3	GLY	2.1
1	A	50	TYR	2.1
1	A	123	GLN	2.1
1	A	332	ALA	2.1
1	B	122	LYS	2.1
1	A	283	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	404	5/5	0.28	0.17	103,109,117,120	0
3	SO4	B	405	5/5	0.46	0.14	102,104,113,123	0
3	SO4	A	405	5/5	0.63	0.12	101,105,114,119	0
3	SO4	B	406	5/5	0.71	0.10	96,99,109,114	0
3	SO4	A	403	5/5	0.79	0.11	94,94,107,107	0
3	SO4	B	404	5/5	0.80	0.09	102,104,110,111	0
3	SO4	B	401	5/5	0.87	0.21	83,87,97,100	0
3	SO4	B	402	5/5	0.91	0.15	78,79,83,86	0
2	A1I9M	A	401	39/39	0.91	0.11	61,69,88,96	0
3	SO4	A	402	5/5	0.92	0.16	78,80,81,87	0
2	A1I9M	B	403	39/39	0.94	0.10	60,69,80,86	2




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A1I9M A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1I9M B 403:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.