



wwPDB X-ray Structure Validation Summary Report i

Apr 28, 2025 – 09:56 PM EDT

PDB ID : 4R79 / pdb_00004r79
Title : Mos1 transposase paired-end complex with left transposon end
Authors : Richardson, J.M.
Deposited on : 2014-08-27
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1351 (3.10-3.10)
Clashscore	180529	1454 (3.10-3.10)
Ramachandran outliers	177936	1391 (3.10-3.10)
Sidechain outliers	177891	1391 (3.10-3.10)
RSRZ outliers	164620	1351 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain		
3	H	27	19%	85%	
4	A	345	2%	71%	24% ..
4	B	345	2%	70%	26% ..

2 Entry composition [\(i\)](#)

There are 7 unique types of molecules in this entry. The entry contains 8853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called left Inverted repeat NTS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	25	Total	C	N	O	P	0	0	0
			521	248	100	149	24			
1	E	25	Total	C	N	O	P	0	0	0
			521	248	100	149	24			
1	G	25	Total	C	N	O	P	0	0	0
			524	248	100	151	25			

- Molecule 2 is a DNA chain called left Inverted repeat TS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	28	Total	C	N	O	P	0	0	0
			562	270	99	166	27			
2	F	28	Total	C	N	O	P	0	0	0
			562	270	99	166	27			

- Molecule 3 is a DNA chain called left Inverted repeat NTS H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	27	Total	C	N	O	P	0	0	0
			540	260	94	160	26			


- Molecule 4 is a protein called Mariner Mos1 transposase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	336	Total	C	N	O	S	0	0	0
			2803	1778	507	508	10			
4	B	336	Total	C	N	O	S	0	0	0
			2803	1778	507	508	10			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	THR	LYS	variant	UNP Q7JQ07
A	164	ASN	SER	variant	UNP Q7JQ07
A	210	PRO	ARG	variant	UNP Q7JQ07
A	216	ALA	THR	engineered mutation	UNP Q7JQ07
A	344	PHE	LEU	variant	UNP Q7JQ07
B	45	THR	LYS	variant	UNP Q7JQ07
B	164	ASN	SER	variant	UNP Q7JQ07
B	210	PRO	ARG	variant	UNP Q7JQ07
B	216	ALA	THR	engineered mutation	UNP Q7JQ07
B	344	PHE	LEU	variant	UNP Q7JQ07

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	1	Total Mn 1 1	0	0
6	A	1	Total Mn 1 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	C	1	Total O 1 1	0	0
7	D	1	Total O 1 1	0	0
7	F	1	Total O 1 1	0	0
7	A	1	Total O 1 1	0	0
7	B	1	Total O 1 1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: left Inverted repeat NTS




- Molecule 1: left Inverted repeat NTS



- Molecule 1: left Inverted repeat NTS



- Molecule 2: left Inverted repeat TS




- Molecule 2: left Inverted repeat TS




- Molecule 3: left Inverted repeat NTS H



- Molecule 4: Mariner Mos1 transposase



- Molecule 4: Mariner Mos1 transposase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	119.78Å 86.89Å 132.38Å 90.00° 99.03° 90.00°	Depositor
Resolution (Å)	29.87 – 3.10 29.87 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.87-3.10) 99.4 (29.87-3.10)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.20 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R , R_{free}	0.227 , 0.270 0.227 , 0.270	Depositor DCC
R_{free} test set	2444 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	1.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 70.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8853	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.31	0/586	0.67	1/906 (0.1%)
1	E	0.33	0/586	0.67	1/906 (0.1%)
1	G	0.28	0/589	0.57	0/910
2	D	0.36	0/628	0.67	0/965
2	F	0.34	0/628	0.65	0/965
3	H	0.30	0/603	0.62	0/926
4	A	0.38	0/2876	0.80	0/3881
4	B	0.36	0/2876	0.79	3/3881 (0.1%)
All	All	0.35	0/9372	0.74	5/13340 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	5	VAL	CA-C-N	7.17	127.59	119.92
4	B	5	VAL	C-N-CA	7.17	127.59	119.92
1	E	6	DT	C2'-C3'-O3'	-6.24	102.13	111.50
1	C	6	DT	C2'-C3'-O3'	-6.16	102.27	111.50
4	B	212	GLU	N-CA-C	5.20	117.58	110.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	521	0	284	29	0
1	E	521	0	284	23	0
1	G	524	0	283	39	0
2	D	562	0	317	16	0
2	F	562	0	317	14	0
3	H	540	0	306	39	0
4	A	2803	0	2739	83	0
4	B	2803	0	2739	77	0
5	B	5	0	0	0	0
5	D	5	0	0	0	0
6	A	1	0	0	0	0
6	H	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
7	F	1	0	0	1	0
All	All	8853	0	7269	291	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:17:DG:H2"	1:C:18:DG:H5"	1.26	1.13
4:B:227:ASN:ND2	4:B:266:LEU:HD22	1.66	1.10
1:G:7:DG:H5'	4:B:216:ALA:HB2	1.31	1.08
1:E:4:DG:H2"	1:E:5:DG:C5'	1.83	1.07
4:A:262:THR:O	4:A:266:LEU:HD13	1.56	1.04

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	A	332/345 (96%)	311 (94%)	19 (6%)	2 (1%)	22 53
4	B	332/345 (96%)	309 (93%)	21 (6%)	2 (1%)	22 53
All	All	664/690 (96%)	620 (93%)	40 (6%)	4 (1%)	22 53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	B	210	PRO
4	A	210	PRO
4	B	199	GLN
4	A	41	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	A	298/307 (97%)	288 (97%)	10 (3%)	32 62
4	B	298/307 (97%)	294 (99%)	4 (1%)	65 82
All	All	596/614 (97%)	582 (98%)	14 (2%)	45 70

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	214	VAL
4	A	223	LEU
4	B	336	CYS
4	B	260	ARG
4	B	283	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
4	B	248	HIS
4	B	298	GLN
4	A	286	HIS
4	B	122	HIS
4	B	134	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	B	401	-	4,4,4	0.44	0	6,6,6	0.11	0
5	SO4	D	101	-	4,4,4	0.44	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	C	25/25 (100%)	-0.05	0	100	100	44, 65, 82, 88	0
1	E	25/25 (100%)	0.07	0	100	100	35, 70, 87, 92	0
1	G	25/25 (100%)	0.95	2 (8%)	20	11	52, 113, 147, 152	0
2	D	28/28 (100%)	-0.04	0	100	100	37, 62, 82, 89	0
2	F	28/28 (100%)	-0.03	0	100	100	42, 64, 90, 95	0
3	H	27/27 (100%)	1.14	5 (18%)	4	2	53, 109, 145, 148	0
4	A	336/345 (97%)	-0.06	7 (2%)	63	44	33, 67, 109, 134	0
4	B	336/345 (97%)	-0.10	6 (1%)	67	49	27, 70, 111, 145	0
All	All	830/848 (97%)	0.00	20 (2%)	59	41	27, 70, 117, 152	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	54	DT	4.2
4	B	8	LYS	3.5
3	H	55	DG	3.5
4	A	5	VAL	3.3
4	B	155	GLY	2.9

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	D	101	5/5	0.86	0.18	134,134,150,151	0
5	SO4	B	401	5/5	0.89	0.12	108,121,125,128	0
6	MN	H	101	1/1	0.98	0.03	81,81,81,81	0
6	MN	A	401	1/1	0.98	0.07	69,69,69,69	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.