



## wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 12:36 pm GMT

PDB ID : 9RAF / pdb\_00009raf  
EMDB ID : EMD-53876  
Title : Influenza A/H7N9 polymerase in complex with a 70-mer template in stalled elongation with backtracking and stem.  
Authors : Arragain, B.; Cusack, S.  
Deposited on : 2025-05-20  
Resolution : 2.15 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	2551 ( 1.66 - 2.65 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	716	
2	C	788	
3	G	42	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	M	13	 <p>8% 85% 8% 8%</p>
5	B	757	 <p>5% 87% 9% •</p>
6	T	70	 <p>63% 46% 29% 26%</p>

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 31095 atoms, of which 14762 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase acidic protein.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	A	716	11606	3692	5777	989	1107	41	6	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	349	LYS	GLU	engineered mutation	UNP M9TI86
A	490	ILE	ARG	engineered mutation	UNP M9TI86

- Molecule 2 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	C	250	4139	1288	2094	369	371	17	3	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	74	ARG	GLY	engineered mutation	UNP A0A024E1J5
C	760	GLY	-	expression tag	UNP A0A024E1J5
C	761	TRP	-	expression tag	UNP A0A024E1J5
C	762	SER	-	expression tag	UNP A0A024E1J5
C	763	HIS	-	expression tag	UNP A0A024E1J5
C	764	PRO	-	expression tag	UNP A0A024E1J5
C	765	GLN	-	expression tag	UNP A0A024E1J5
C	766	PHE	-	expression tag	UNP A0A024E1J5
C	767	GLU	-	expression tag	UNP A0A024E1J5
C	768	LYS	-	expression tag	UNP A0A024E1J5
C	769	GLY	-	expression tag	UNP A0A024E1J5
C	770	GLY	-	expression tag	UNP A0A024E1J5
C	771	GLY	-	expression tag	UNP A0A024E1J5
C	772	SER	-	expression tag	UNP A0A024E1J5
C	773	GLY	-	expression tag	UNP A0A024E1J5

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	774	GLY	-	expression tag	UNP A0A024E1J5
C	775	GLY	-	expression tag	UNP A0A024E1J5
C	776	SER	-	expression tag	UNP A0A024E1J5
C	777	GLY	-	expression tag	UNP A0A024E1J5
C	778	GLY	-	expression tag	UNP A0A024E1J5
C	779	SER	-	expression tag	UNP A0A024E1J5
C	780	ALA	-	expression tag	UNP A0A024E1J5
C	781	TRP	-	expression tag	UNP A0A024E1J5
C	782	SER	-	expression tag	UNP A0A024E1J5
C	783	HIS	-	expression tag	UNP A0A024E1J5
C	784	PRO	-	expression tag	UNP A0A024E1J5
C	785	GLN	-	expression tag	UNP A0A024E1J5
C	786	PHE	-	expression tag	UNP A0A024E1J5
C	787	GLU	-	expression tag	UNP A0A024E1J5
C	788	LYS	-	expression tag	UNP A0A024E1J5

- Molecule 3 is a protein called RNA Pol II CTD 6 repeats (site 1A/2A).

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			P
3	G	25	351	116	155	25	52	3	0	0

- Molecule 4 is a RNA chain called RNA product.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			P
4	M	13	422	124	143	52	90	13	0	0

- Molecule 5 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
5	B	725	11679	3672	5840	1028	1094	45	8	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	577	GLY	LYS	engineered mutation	UNP S5ME50

- Molecule 6 is a RNA chain called RNA template (70-mer).

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			P
6	T	70	2243	665	753	262	493	70	0	0

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
7	A	1	Total	Mg	0
			1	1	
7	B	2	Total	Mg	0
			2	2	


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		AltConf
8	A	226	Total	O	0
			226	226	
8	C	54	Total	O	0
			54	54	
8	G	16	Total	O	0
			16	16	
8	M	19	Total	O	0
			19	19	
8	B	254	Total	O	0
			254	254	
8	T	83	Total	O	0
			83	83	


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polymerase acidic protein




- Molecule 2: Polymerase basic protein 2







• Molecule 3: RNA Pol II CTD 6 repeats (site 1A/2A)




• Molecule 4: RNA product



• Molecule 5: RNA-directed RNA polymerase catalytic subunit



• Molecule 6: RNA template (70-mer)



165	166	167	168	169	170
-----	-----	-----	-----	-----	-----

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	195294	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	41	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	4.697	Depositor
Minimum map value	-1.832	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.089	Depositor
Recommended contour level	0.6	Depositor
Map size (Å)	306.6, 306.6, 306.6	wwPDB
Map dimensions	420, 420, 420	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.73, 0.73, 0.73	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.12	0/5974	0.23	0/8055
2	C	0.25	1/2093 (0.0%)	0.26	0/2821
3	G	0.12	0/173	0.23	0/235
4	M	0.14	0/311	0.18	0/483
5	B	0.12	0/5979	0.25	0/8055
6	T	0.13	0/1664	0.27	0/2589
All	All	0.15	1/16194 (0.0%)	0.25	0/22238

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	46	ARG	C-O	-9.39	1.13	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5829	5777	5754	33	0
2	C	2045	2094	2083	11	0
3	G	196	155	155	5	0
4	M	279	143	143	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	5839	5840	5805	58	0
6	T	1490	753	753	34	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
8	A	226	0	0	9	0
8	B	254	0	0	17	0
8	C	54	0	0	2	0
8	G	16	0	0	3	0
8	M	19	0	0	0	0
8	T	83	0	0	9	0
All	All	16333	14762	14693	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:T:50:U:O2'	6:T:52:U:OP2	1.90	0.87
6:T:65:U:O2'	6:T:66:U:OP2	1.94	0.86
1:A:488:LYS:NZ	6:T:69:C:OP1	2.12	0.83
6:T:29:C:O2	6:T:51:G:N2	2.11	0.82
1:A:597:GLU:OE1	8:A:901:HOH:O	1.99	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	720/716 (101%)	707 (98%)	13 (2%)	0	100	100
2	C	251/788 (32%)	249 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	G	20/42 (48%)	18 (90%)	2 (10%)	0	100	100
5	B	725/757 (96%)	713 (98%)	12 (2%)	0	100	100
All	All	1716/2303 (74%)	1687 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	651/645 (101%)	645 (99%)	6 (1%)	75	81
2	C	227/686 (33%)	224 (99%)	3 (1%)	65	71
3	G	22/36 (61%)	22 (100%)	0	100	100
5	B	647/667 (97%)	641 (99%)	6 (1%)	75	81
All	All	1547/2034 (76%)	1532 (99%)	15 (1%)	71	78

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	83	ASP
5	B	614	GLU
2	C	217	PHE
5	B	682	ILE
5	B	204	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
5	B	367	GLN
5	B	375	ASN
5	B	690	GLN
5	B	428	GLN
1	A	519	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	M	12/13 (92%)	1 (8%)	0
6	T	69/70 (98%)	19 (27%)	4 (5%)
All	All	81/83 (97%)	20 (24%)	4 (4%)

5 of 20 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	M	20	U
6	T	7	A
6	T	8	A
6	T	11	A
6	T	14	A

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
6	T	13	G
6	T	15	U
6	T	46	U
6	T	65	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SEP	G	21	3	8,9,10	1.50	1 (12%)	8,12,14	1.48	2 (25%)
3	SEP	G	7	3	8,9,10	1.51	1 (12%)	8,12,14	1.47	2 (25%)
3	SEP	G	14	3	8,9,10	1.56	1 (12%)	8,12,14	1.61	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SEP	G	21	3	-	0/5/8/10	-
3	SEP	G	7	3	-	2/5/8/10	-
3	SEP	G	14	3	-	3/5/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	14	SEP	P-O1P	3.41	1.61	1.50
3	G	21	SEP	P-O1P	3.30	1.61	1.50
3	G	7	SEP	P-O1P	3.28	1.61	1.50

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	14	SEP	OG-CB-CA	3.41	111.47	108.14
3	G	21	SEP	OG-CB-CA	2.74	110.81	108.14
3	G	7	SEP	OG-CB-CA	2.71	110.78	108.14
3	G	7	SEP	P-OG-CB	-2.45	111.55	118.30
3	G	21	SEP	P-OG-CB	-2.42	111.62	118.30

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	14	SEP	N-CA-CB-OG
3	G	14	SEP	CA-CB-OG-P
3	G	7	SEP	N-CA-CB-OG
3	G	14	SEP	CB-OG-P-O2P
3	G	7	SEP	CA-CB-OG-P

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	21	SEP	2	0
3	G	14	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-53876. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 210




Y Index: 210




Z Index: 210


### 6.2.2 Raw map



X Index: 210



Y Index: 210




Z Index: 210


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 200




Y Index: 205




Z Index: 215


### 6.3.2 Raw map



X Index: 201



Y Index: 218




Z Index: 215


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 53 nm<sup>3</sup>; this corresponds to an approximate mass of 48 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.465 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.465 \text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.15	-	-
Author-provided FSC curve	2.15	2.52	2.18
Unmasked-calculated*	2.94	3.64	3.01

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.94 differs from the reported value 2.15 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-53876 and PDB model 9RAF. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.6).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7910	 0.6510
A	 0.8200	 0.6770
B	 0.8800	 0.6980
C	 0.8170	 0.6700
G	 0.5780	 0.5810
M	 0.7810	 0.6310
T	 0.3570	 0.3560

