



wwPDB EM Validation Summary Report i

Jan 5, 2026 – 05:18 pm GMT

PDB ID : 9RBQ / pdb_00009rbq
EMDB ID : EMD-53902
Title : Semliki Forest virus trimer 1 in complex with ApoER2 LA5
Authors : Song, X.; Du, B.; Yang, D.; Wang, J.; Huiskonen, J.T.
Deposited on : 2025-05-27
Resolution : 2.70 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev129
Mogul : ?.? (??), CSD ??CSD?? (????)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	10327 (2.20 - 3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 25585 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LDL receptor related protein 8,Ig gamma-2A chain C region, A allele.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	42	312	183	59	64	6	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	200	GLY	-	expression tag	UNP E9PKG2
A	201	SER	-	expression tag	UNP E9PKG2
A	255	ALA	-	linker	UNP E9PKG2
A	256	PRO	-	linker	UNP E9PKG2
A	257	ALA	-	linker	UNP E9PKG2
A	258	ALA	-	linker	UNP E9PKG2
A	259	GLY	-	linker	UNP E9PKG2
A	260	SER	-	linker	UNP E9PKG2
A	261	LEU	-	linker	UNP E9PKG2
A	262	GLU	-	linker	UNP E9PKG2
A	263	VAL	-	linker	UNP E9PKG2
A	264	LEU	-	linker	UNP E9PKG2
A	265	PHE	-	linker	UNP E9PKG2
A	266	GLN	-	linker	UNP E9PKG2
A	267	GLY	-	linker	UNP E9PKG2
A	268	PRO	-	linker	UNP E9PKG2

- Molecule 2 is a protein called Protein E3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	D	54	415	259	69	79	8	0	0
2	E	54	415	259	69	79	8	0	0
2	F	54	415	259	69	79	8	0	0

- Molecule 3 is a protein called Structural polyprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	438	Total	C	N	O	S		
			3329	2109	559	636	25	0	0
3	H	438	Total	C	N	O	S		
			3329	2109	559	636	25	0	0
3	I	438	Total	C	N	O	S		
			3329	2109	559	636	25	0	0

- Molecule 4 is a protein called Envelope glycoprotein E2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	J	421	Total	C	N	O	S		
			3282	2071	583	603	25	0	0
4	K	421	Total	C	N	O	S		
			3282	2071	583	603	25	0	0
4	L	421	Total	C	N	O	S		
			3282	2071	583	603	25	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	162	LYS	GLU	conflict	UNP P0DJZ6
K	162	LYS	GLU	conflict	UNP P0DJZ6
L	162	LYS	GLU	conflict	UNP P0DJZ6

- Molecule 5 is a protein called Capsid protein.


Mol	Chain	Residues	Atoms					AltConf	Trace
5	M	162	Total	C	N	O	S		
			1250	781	227	233	9	0	0
5	N	162	Total	C	N	O	S		
			1250	781	227	233	9	0	0
5	O	162	Total	C	N	O	S		
			1250	781	227	233	9	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
6	B	4	Total	C	N	O	0	0
			50	28	2	20		
6	R	4	Total	C	N	O	0	0
			50	28	2	20		
6	V	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
7	Z	3	Total	C	N	O	0	0
			42	24	3	15		
7	g	3	Total	C	N	O	0	0
			42	24	3	15		
7	n	3	Total	C	N	O	0	0
			42	24	3	15		

- Molecule 8 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
8	c	4	Total	C	N	O	0	0
			56	32	4	20		
8	j	4	Total	C	N	O	0	0
			56	32	4	20		
8	q	4	Total	C	N	O	0	0
			56	32	4	20		


- Molecule 9 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
9	A	1	Total 1 1	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LDL receptor related protein 8, Ig gamma-2A chain C region, A allele



- Molecule 2: Protein E3



- Molecule 2: Protein E3




• Molecule 2: Protein E3







- Molecule 3: Structural polyprotein




- Molecule 3: Structural polyprotein



- Molecule 3: Structural polyprotein




- Molecule 4: Envelope glycoprotein E2




- Molecule 4: Envelope glycoprotein E2

Chain K:  76% 23%




- Molecule 4: Envelope glycoprotein E2

Chain L:  81% 18%




- Molecule 5: Capsid protein


Chain M:  45% 16% 39%




- Molecule 5: Capsid protein

Chain N:  41% 19% 39%





- Molecule 5: Capsid protein



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-D-mannopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

ucopyranose

Chain q:  25% 75%

MAG1
MAG2
MAG3
MAG4

4 Experimental information i

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	2344552	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.066	Depositor
Minimum map value	-0.036	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.005	Depositor
Map size (Å)	276.48, 276.48, 276.48	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.08, 1.08, 1.08	Depositor

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA, MAN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/317	0.25	0/424
2	D	0.09	0/424	0.26	0/580
2	E	0.10	0/424	0.32	0/580
2	F	0.11	0/424	0.31	0/580
3	G	0.13	0/3415	0.30	0/4660
3	H	0.13	0/3415	0.29	0/4660
3	I	0.14	0/3415	0.30	0/4660
4	J	0.13	0/3376	0.32	0/4605
4	K	0.12	0/3376	0.30	0/4605
4	L	0.13	0/3376	0.31	0/4605
5	M	0.10	0/1278	0.30	0/1719
5	N	0.13	0/1278	0.36	0/1719
5	O	0.10	0/1278	0.28	0/1719
All	All	0.13	0/25796	0.30	0/35116

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	312	0	265	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	415	0	389	12	0
2	E	415	0	389	17	0
2	F	415	0	389	11	0
3	G	3329	0	3245	37	0
3	H	3329	0	3247	41	0
3	I	3329	0	3245	45	0
4	J	3282	0	3195	59	0
4	K	3282	0	3197	73	0
4	L	3282	0	3195	60	0
5	M	1250	0	1232	27	0
5	N	1250	0	1232	37	0
5	O	1250	0	1232	35	0
6	B	50	0	43	0	0
6	R	50	0	43	0	0
6	V	50	0	43	0	0
7	Z	42	0	37	1	0
7	g	42	0	37	2	0
7	n	42	0	37	0	0
8	c	56	0	49	0	0
8	j	56	0	49	2	0
8	q	56	0	49	2	0
9	A	1	0	0	0	0
All	All	25585	0	24839	417	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 417 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:327:ASP:OD1	3:I:345:LYS:HG2	1.55	1.06
3:H:327:ASP:OD1	3:H:345:LYS:HG2	1.70	0.92
4:L:201:CYS:HA	4:L:225:CYS:HB2	1.59	0.84
5:O:106:GLY:O	5:O:107:LYS:HG2	1.79	0.83
5:O:173:ILE:HD11	5:O:178:ARG:HD2	1.60	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	40/301 (13%)	38 (95%)	2 (5%)	0	100 100
2	D	52/66 (79%)	51 (98%)	1 (2%)	0	100 100
2	E	52/66 (79%)	47 (90%)	5 (10%)	0	100 100
2	F	52/66 (79%)	47 (90%)	5 (10%)	0	100 100
3	G	436/438 (100%)	430 (99%)	6 (1%)	0	100 100
3	H	436/438 (100%)	429 (98%)	7 (2%)	0	100 100
3	I	436/438 (100%)	430 (99%)	6 (1%)	0	100 100
4	J	419/422 (99%)	404 (96%)	15 (4%)	0	100 100
4	K	419/422 (99%)	405 (97%)	14 (3%)	0	100 100
4	L	419/422 (99%)	400 (96%)	19 (4%)	0	100 100
5	M	160/267 (60%)	153 (96%)	7 (4%)	0	100 100
5	N	160/267 (60%)	150 (94%)	10 (6%)	0	100 100
5	O	160/267 (60%)	155 (97%)	5 (3%)	0	100 100
All	All	3241/3880 (84%)	3139 (97%)	102 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	31/266 (12%)	31 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	46/56 (82%)	46 (100%)	0	100	100
2	E	46/56 (82%)	46 (100%)	0	100	100
2	F	46/56 (82%)	46 (100%)	0	100	100
3	G	372/372 (100%)	372 (100%)	0	100	100
3	H	372/372 (100%)	372 (100%)	0	100	100
3	I	372/372 (100%)	372 (100%)	0	100	100
4	J	360/360 (100%)	360 (100%)	0	100	100
4	K	360/360 (100%)	360 (100%)	0	100	100
4	L	360/360 (100%)	360 (100%)	0	100	100
5	M	132/224 (59%)	132 (100%)	0	100	100
5	N	132/224 (59%)	132 (100%)	0	100	100
5	O	132/224 (59%)	132 (100%)	0	100	100
All	All	2761/3302 (84%)	2761 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
5	M	117	ASN
5	M	194	ASN
5	N	194	ASN
4	J	238	ASN
4	J	224	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

33 monosaccharides are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.























5.6 Ligand geometry [\(i\)](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-53902. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [\(i\)](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 108




Y Index: 154




Z Index: 117


6.3.2 Raw map



X Index: 136



Y Index: 154




Z Index: 108


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [\(i\)](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.005. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [\(i\)](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_53902_msk_1.map [\(i\)](#)



X



Y




Z

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 887 nm^3 ; this corresponds to an approximate mass of 801 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.370 \AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.370 Å⁻¹

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.70	-	-
Author-provided FSC curve	2.66	3.01	2.69
Unmasked-calculated*	2.68	3.06	2.73

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [\(i\)](#)


This section contains information regarding the fit between EMDB map EMD-53902 and PDB model 9RBQ. Per-residue inclusion information can be found in section [3](#) on page [8](#).

9.1 Map-model overlay [\(i\)](#)




The images above show the 3D surface view of the map at the recommended contour level 0.005 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.005).

9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 95% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.005) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8900	0.6020
A	0.6130	0.4890
B	0.5200	0.4260
D	0.4900	0.5120
E	0.6110	0.5470
F	0.4560	0.5170
G	0.9440	0.6430
H	0.9410	0.6310
I	0.9560	0.6530
J	0.9210	0.6180
K	0.9240	0.6260
L	0.9290	0.6260
M	0.8170	0.4670
N	0.8120	0.4770
O	0.8420	0.5160
R	0.5000	0.3870
V	0.6600	0.5630
Z	0.6670	0.4390
c	0.6790	0.4700
g	0.7140	0.4540
j	0.6250	0.4690
n	0.7380	0.4840
q	0.6960	0.5130

