



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 16, 2026 – 03:45 pm GMT

PDB ID : 9ROS / pdb\_00009ros  
Title : Atomic resolution (1.15 Å) XFEL structure of as-isolated copper nitrite reductase from *Bradyrhizobium* sp. at low pH (5.5) determined by serial femtosecond rotation crystallography (SF-ROX) at 100 K  
Authors : Rose, S.L.; Antonyuk, S.V.; Ferroni, F.M.; Sugimoto, H.; Yamashita, K.; Hirata, K.; Ago, H.; Ueno, G.; Murakami, H.; Eady, R.R.; Tosha, T.; Yamamoto, M.; Hasnain, S.S.  
Deposited on : 2025-06-22  
Resolution : 1.15 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1095 (1.16-1.12)
Clashscore	180529	1248 (1.16-1.12)
Ramachandran outliers	177936	1224 (1.16-1.12)
Sidechain outliers	177891	1224 (1.16-1.12)
RSRZ outliers	164620	1095 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6081 atoms, of which 2793 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	345	5491	1755	2741	481	500	14	0	14	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	GLU	-	expression tag	UNP H0SLX7
A	343	ASN	-	expression tag	UNP H0SLX7
A	344	LEU	-	expression tag	UNP H0SLX7
A	345	TYR	-	expression tag	UNP H0SLX7
A	346	PHE	-	expression tag	UNP H0SLX7
A	347	GLN	-	expression tag	UNP H0SLX7

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	0
			2	2		

- Molecule 3 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	H	O	0	1
			21	6	10	5		
3	A	1	Total	C	H	O	0	1
			21	6	10	5		

- Molecule 4 is beta-D-fructofuranose (CCD ID: FRU) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	A	1	Total	C	H	O	0	1
			24	6	12	6		
4	A	1	Total	C	H	O	0	1
			24	6	12	6		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	1
5	A	1	Total O S 5 4 1	0	1
5	A	1	Total O S 5 4 1	0	1
5	A	1	Total O S 5 4 1	0	1
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	1
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	421	Total	O	0	34
			444	444		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Copper-containing nitrite reductase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.32Å 107.32Å 107.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.00 – 1.15 17.00 – 1.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (17.00-1.15) 100.0 (17.00-1.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.65 (at 1.10Å)	Xtrriage
Refinement program	SHELXL	Depositor
R, $R_{free}$	0.150 , 0.174 0.152 , 0.174	Depositor DCC
$R_{free}$ test set	8333 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.49 , 73.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.063 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6081	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, GLC, GOL, FRU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.90	0/2863	1.04	3/3884 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	92	ASP	CA-CB-CG	9.74	122.34	112.60
1	A	16	PRO	O-C-N	-6.63	118.26	121.31
1	A	337	THR	CA-CB-OG1	-5.69	101.06	109.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	ARG	Sidechain
1	A	154	ARG	Sidechain
1	A	299	VAL	Peptide
1	A	301	HIS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	2741	2740	23	2
2	A	2	0	0	0	0
3	A	22	20	20	3	0
4	A	24	24	24	3	0
5	A	40	0	0	2	0
6	A	6	8	8	3	0
7	A	444	0	0	10	4
All	All	3288	2793	2792	29	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:505[A]:GLC:C1	4:A:506[A]:FRU:O2	1.89	1.21
3:A:503[A]:GLC:C1	4:A:504[A]:FRU:O2	1.92	1.16
1:A:261[B]:GLU:HG3	7:A:608:HOH:O	1.75	0.85
1:A:320:LYS:HE2	7:A:609:HOH:O	1.79	0.83
1:A:43[A]:LYS:HE2	7:A:771:HOH:O	1.87	0.74

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:604:HOH:O	7:A:876:HOH:O[2_565]	1.51	0.69
1:A:48[B]:GLU:OE1	7:A:879:HOH:O[5_555]	2.06	0.14
7:A:775:HOH:O	7:A:981:HOH:O[6_456]	2.11	0.09
1:A:279[B]:ARG:HH22	7:A:939[B]:HOH:O[6_456]	1.55	0.05

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/346 (103%)	352 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/282 (104%)	289 (98%)	5 (2%)	56	19

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166[A]	ARG
1	A	166[B]	ARG
1	A	242	GLN
1	A	279[A]	ARG
1	A	279[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	54	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	A	514	-	4,4,4	0.28	0	6,6,6	0.28	0
5	SO4	A	509[A]	-	4,4,4	0.40	0	6,6,6	0.19	0
5	SO4	A	512	-	4,4,4	0.42	0	6,6,6	0.45	0
4	FRU	A	506[A]	-	11,12,12	0.50	0	10,18,18	0.92	0
5	SO4	A	513	-	4,4,4	0.46	0	6,6,6	0.25	0
5	SO4	A	515	-	4,4,4	0.65	0	6,6,6	0.20	0
3	GLC	A	503[A]	-	11,11,12	0.74	0	15,15,17	1.70	3 (20%)
3	GLC	A	505[A]	-	11,11,12	1.21	1 (9%)	15,15,17	2.15	3 (20%)
5	SO4	A	508[A]	-	4,4,4	0.53	0	6,6,6	0.09	0
5	SO4	A	510[A]	-	4,4,4	1.49	1 (25%)	6,6,6	0.50	0
4	FRU	A	504[A]	-	11,12,12	0.61	0	10,18,18	0.75	0
6	GOL	A	511[A]	-	5,5,5	0.22	0	5,5,5	0.25	0
5	SO4	A	507[A]	-	4,4,4	0.39	0	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FRU	A	506[A]	-	-	0/5/24/24	0/1/1/1
3	GLC	A	503[A]	-	-	2/2/19/22	0/1/1/1
3	GLC	A	505[A]	-	-	0/2/19/22	0/1/1/1
4	FRU	A	504[A]	-	-	0/5/24/24	0/1/1/1
6	GOL	A	511[A]	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	510[A]	SO4	O2-S	2.73	1.60	1.46
3	A	505[A]	GLC	O5-C5	2.23	1.48	1.43

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	505[A]	GLC	C1-O5-C5	5.94	120.24	112.19
3	A	505[A]	GLC	C1-C2-C3	4.05	114.64	109.67
3	A	503[A]	GLC	O5-C1-C2	3.45	116.09	110.77
3	A	503[A]	GLC	O5-C5-C4	-2.65	104.39	110.83
3	A	503[A]	GLC	O3-C3-C2	-2.50	105.20	109.99

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	511[A]	GOL	O1-C1-C2-C3
6	A	511[A]	GOL	C1-C2-C3-O3
6	A	511[A]	GOL	O1-C1-C2-O2
6	A	511[A]	GOL	O2-C2-C3-O3
3	A	503[A]	GLC	O5-C5-C6-O6

There are no ring outliers.

7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	509[A]	SO4	1	0
4	A	506[A]	FRU	2	0
5	A	515	SO4	1	0
3	A	503[A]	GLC	1	0
3	A	505[A]	GLC	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	504[A]	FRU	1	0
6	A	511[A]	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/346 (99%)	-0.46	9 (2%) 57 65	5, 14, 28, 73	32 (9%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	LEU	5.8
1	A	346	PHE	5.1
1	A	341	ASN	3.7
1	A	345	TYR	3.4
1	A	347	GLN	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	FRU	A	506[A]	12/12	0.74	0.18	29,34,37,41	24
4	FRU	A	504[A]	12/12	0.85	0.14	33,46,52,55	24

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	A	510[A]	5/5	0.85	0.18	117,118,126,129	5
5	SO4	A	513	5/5	0.88	0.11	70,70,75,75	0
3	GLC	A	503[A]	11/12	0.90	0.10	28,34,43,47	21
5	SO4	A	514	5/5	0.90	0.12	31,35,36,37	5
5	SO4	A	515	5/5	0.90	0.13	25,32,40,40	5
3	GLC	A	505[A]	11/12	0.91	0.12	14,22,25,27	21
5	SO4	A	508[A]	5/5	0.91	0.10	52,54,57,59	5
5	SO4	A	507[A]	5/5	0.93	0.10	23,24,27,28	5
6	GOL	A	511[A]	6/6	0.93	0.07	39,41,46,50	0
5	SO4	A	509[A]	5/5	0.97	0.07	34,37,39,46	5
5	SO4	A	512	5/5	0.97	0.06	24,25,26,27	5
2	CU	A	502	1/1	1.00	0.03	12,12,12,12	0
2	CU	A	501	1/1	1.00	0.01	12,12,12,12	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.