

wwPDB EM Validation Summary Report (i)

Feb 24, 2025 - 03:04 pm GMT

PDB ID : 8S3O EMDB ID : EMD-19699 Title : Structure of native human CD109 Authors : Almeida, V.A.; Andersen, G.R. Deposited on 2024-02-20 : 2.99 Å(reported) Resolution : Based on initial model :

Based on initial model : .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis	:	FAILED
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	FAILED
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive}\ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	А	1447	84%	8%	9%
2	В	2	50% 50	%	
2	С	2	100%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 21074 atoms, of which 10518 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CD109 antigen.

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	А	1324	Total	С	Η	Ν	0	\mathbf{S}	0	0
		1021	20860	6676	10416	1716	2015	37		5

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-20	SER	-	expression tag	UNP Q6YHK3
А	-19	VAL	-	expression tag	UNP Q6YHK3
А	-18	SER	-	expression tag	UNP Q6YHK3
А	-17	GLY	-	expression tag	UNP Q6YHK3
А	-16	TRP	-	expression tag	UNP Q6YHK3
A	-15	SER	-	expression tag	UNP Q6YHK3
А	-14	HIS	-	expression tag	UNP Q6YHK3
А	-13	PRO	-	expression tag	UNP Q6YHK3
А	-12	GLN	-	expression tag	UNP Q6YHK3
А	-11	PHE	-	expression tag	UNP Q6YHK3
А	-10	GLU	-	expression tag	UNP Q6YHK3
А	-9	LYS	-	expression tag	UNP Q6YHK3
А	-8	GLY	-	expression tag	UNP Q6YHK3
А	-7	GLY	-	expression tag	UNP Q6YHK3
А	-6	GLY	-	expression tag	UNP Q6YHK3
А	-5	SER	-	expression tag	UNP Q6YHK3
А	-4	GLY	-	expression tag	UNP Q6YHK3
А	-3	GLY	-	expression tag	UNP Q6YHK3
А	-2	GLY	-	expression tag	UNP Q6YHK3
А	-1	SER	-	expression tag	UNP Q6YHK3
А	0	GLY	-	expression tag	UNP Q6YHK3
А	1	GLY	-	expression tag	UNP Q6YHK3
А	2	SER	-	expression tag	UNP Q6YHK3
А	3	ALA	-	expression tag	UNP Q6YHK3
А	4	TRP	-	expression tag	UNP Q6YHK3
А	5	SER	-	expression tag	UNP Q6YHK3
А	6	HIS	-	expression tag	UNP Q6YHK3
А	7	PRO	-	expression tag	UNP Q6YHK3

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
А	8	GLN	-	expression tag	UNP Q6YHK3
А	9	PHE	-	expression tag	UNP Q6YHK3
А	10	GLU	-	expression tag	UNP Q6YHK3
А	11	LYS	-	expression tag	UNP Q6YHK3
А	12	GLY	-	expression tag	UNP Q6YHK3
А	13	GLU	-	expression tag	UNP Q6YHK3
А	14	ASN	-	expression tag	UNP Q6YHK3
А	15	LEU	-	expression tag	UNP Q6YHK3
А	16	TYR	-	expression tag	UNP Q6YHK3
А	17	PHE	-	expression tag	UNP Q6YHK3
А	18	GLN	-	expression tag	UNP Q6YHK3
А	19	SER	-	expression tag	UNP Q6YHK3
А	20	GLY	-	expression tag	UNP Q6YHK3
А	21	SER	-	expression tag	UNP Q6YHK3
А	1421	HIS	-	expression tag	UNP Q6YHK3
А	1422	HIS	-	expression tag	UNP Q6YHK3
А	1423	HIS	-	expression tag	UNP Q6YHK3
А	1424	HIS	-	expression tag	UNP Q6YHK3
А	1425	HIS	-	expression tag	UNP Q6YHK3
А	1426	HIS	-	expression tag	UNP Q6YHK3

Continued from previous page...


• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace	
2	В	2	Total	-			-	0	0
		2	53	16	25	2	10	0	0
2	С	9	Total	С	Η	Ν	Ο	0	0
2	U	2	53	16	25	2	10	0	

• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf	
3	٨	1	Total	С	Η	Ν	Ο	0
0	A	1	27	8	13	1	5	0
9	٨	1	Total	С	Η	Ν	Ο	0
3	А	1	27	8	13	1	5	0
3	Δ	1	Total	С	Η	Ν	Ο	0
J	А	1	27	8	13	1	5	0
3	٨	1	Total	С	Η	Ν	Ο	0
3	A	1	27	8	13	1	5	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:	84%	8% 9%
SER VAL SER CLP TRP CLP SER PFIG CLU CLU CLU CLU CLV CLU CLV CLV CLV CLV CLV CLV CLV CLV CLV CLV	HIS PHE PHE CUN CUN CUN CUN CUN CUN CUN CUN CUN CUN	L48 157 157 157 157 82 892 1104 1104 1110
q111 1113 1114 1114 1114 1114 1114 1114	ASP LEU SER SER SER E322 E322 E325 E355 T356 T356 T356 T356 T356 T356 T356 T	1394 E400 TTR TTR TTR TTR TTR SER ASN ASN ASN ASN ASN ASN ASN ASN ASN
K411 V415 K417 K417 S425 S425 F432 I434 I434 K444 K444 K444 K444 K444 K444	1515 C530 C530 N557 K565 K565 K565 K565 K566 N603 D604 D604	ASP ASP ASP ASP ASP ASP ASP ASP
ASN ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	SER PRO HISO VAL ARG LISS D698 D698 S728 S728 S728 S728 S728 S728 S728 S774 IT770	7776 1799 1809 1808 1808 1808 1818 1818
L819 1830 V881 2872 1872 2922 6923 6923 6923 6924 6924 6926 7929 7929 7976 8976 8976 8976 8976	11066 V1068 V1068 N1085 L1086 E1106 E1106 E1116 C1120 C1120 C1120 C1120 C1120 C1120	11184 E1201 E1201 R1227 L1235 L1235 C128 GLY
SER SER ARG ARG ARG ARG B1274 E1279 D1282 L1283 L1283 L1283 N1368 N1368 N1384 T1384 R1383 S1408 S1408 S1408	GLU ASP GLY GLY HLS HLS HLS HLS HLS HLS HLS	
• Molecule 2: 2-acetamido-2-deoxy-	-beta-D-glucopyranose-(1-4)-2	2-acetamido-2-deoxy

• Molecule 1: CD109 antigen

y-beta-D-gluc opyranose

Chain B:	50%	50%
8		

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain C:

100%

NAC



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	273564	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	59.9	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1700	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	ol Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.29	0/10660	0.55	0/14476	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	10444	10416	10420	56	0
2	В	28	25	25	1	0
2	С	28	25	25	0	0
3	А	56	52	52	2	0
All	All	10556	10518	10522	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1283:LEU:O	1:A:1393:ARG:NH1	2.28	0.67
1:A:604:ASP:O	1:A:609:ASN:ND2	2.31	0.63
1:A:394:THR:HG23	1:A:415:VAL:HG12	1.82	0.62
1:A:48:LEU:HD11	1:A:56:VAL:HG22	1.82	0.61
1:A:819:LEU:HD12	1:A:1408:SER:HB3	1.85	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	1314/1447~(91%)	1257 (96%)	54 (4%)	3~(0%)	44 77	

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	929	PHE
1	А	228	LYS
1	А	391	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	1190/1289~(92%)	1174 (99%)	16 (1%)	65 85		



5 of 16 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	1288	LYS
1	А	921	CYS
1	А	698	ASP
1	А	872	ARG
1	А	626	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

М	Mol Type	Turne	Chain	Res	Link	Bo	Bond lengths			Bond angles		
		rybe	Chain			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
2		NAG	В	1	2,1	14,14,15	0.34	0	$17,\!19,\!21$	0.43	0	
2	2	NAG	В	2	2	14,14,15	0.22	0	17,19,21	0.81	1 (5%)	
2	2	NAG	С	1	2,1	14,14,15	0.20	0	17,19,21	0.52	0	
2	2	NAG	С	2	2	14,14,15	0.16	0	17,19,21	0.43	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	В	2	2	-	2/6/23/26	0/1/1/1
2	NAG	С	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	В	2	NAG	C2-N2-C7	2.67	126.71	122.90

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	2	NAG	O5-C5-C6-O6
2	С	2	NAG	C4-C5-C6-O6
2	С	1	NAG	O5-C5-C6-O6
2	В	2	NAG	C8-C7-N2-C2
2	В	2	NAG	O7-C7-N2-C2


There are no ring outliers.

1 monomer is involved in 1 short contact:


M	bl	Chain	Res	Type	Clashes	Symm-Clashes
2		В	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	Mol Type Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
3	NAG	А	1503	1	14,14,15	0.20	0	17,19,21	0.41	0
3	NAG	А	1502	1	14,14,15	0.24	0	17,19,21	1.32	2 (11%)
3	NAG	А	1504	1	14,14,15	0.26	0	17,19,21	0.42	0
3	NAG	А	1501	1	14,14,15	0.20	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	А	1503	1	-	1/6/23/26	0/1/1/1
3	NAG	А	1502	1	-	3/6/23/26	0/1/1/1
3	NAG	А	1504	1	-	0/6/23/26	0/1/1/1
3	NAG	А	1501	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	1502	NAG	C2-N2-C7	4.47	129.27	122.90
3	А	1502	NAG	C1-C2-N2	2.03	113.96	110.49

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	1501	NAG	O5-C5-C6-O6
3	А	1502	NAG	C8-C7-N2-C2
3	А	1502	NAG	O7-C7-N2-C2
3	А	1501	NAG	C4-C5-C6-O6
3	А	1502	NAG	C3-C2-N2-C7

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	1502	NAG	1	0
3	А	1501	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

