



## wwPDB EM Validation Summary Report ⓘ

Apr 21, 2026 – 07:41 pm BST

PDB ID : 9SDL / pdb\_00009sdl  
EMDB ID : EMD-54787  
Title : Cryo-EM structure of PfHT1 bound to 2,5-anhydro-D-mannitol  
Authors : Gulati, A.; Suades, A.; Drew, D.  
Deposited on : 2025-08-14  
Resolution : 2.42 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4-5-2 with Phenix2.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49



# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	5729 ( 1.92 - 2.92 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	749	 59% 5% 36%
1	B	749	 59% 5% 36%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15452 atoms, of which 7848 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hexose transporter 1, Green fluorescent protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	478	Total	C	H	N	O	S	0	0
			7703	2526	3912	586	659	20		
1	B	478	Total	C	H	N	O	S	0	0
			7703	2526	3912	586	659	20		

There are 58 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	15	MET	-	initiating methionine	UNP O97467
A	16	GLU	-	expression tag	UNP O97467
A	17	LYS	-	expression tag	UNP O97467
A	18	GLU	-	expression tag	UNP O97467
A	19	ASP	-	expression tag	UNP O97467
A	505	GLU	-	linker	UNP O97467
A	506	ASN	-	linker	UNP O97467
A	507	LEU	-	linker	UNP O97467
A	508	TYR	-	linker	UNP O97467
A	509	PHE	-	linker	UNP O97467
A	510	GLN	-	linker	UNP O97467
A	511	GLY	-	linker	UNP O97467
A	512	GLN	-	linker	UNP O97467
A	513	PHE	-	linker	UNP O97467
A	577	GLY	SER	conflict	UNP P42212
A	584	ALA	SER	conflict	UNP P42212
A	751	ASP	-	expression tag	UNP P42212
A	752	GLU	-	expression tag	UNP P42212
A	753	LEU	-	expression tag	UNP P42212
A	754	TYR	-	expression tag	UNP P42212
A	755	LYS	-	expression tag	UNP P42212
A	756	HIS	-	expression tag	UNP P42212
A	757	HIS	-	expression tag	UNP P42212
A	758	HIS	-	expression tag	UNP P42212
A	759	HIS	-	expression tag	UNP P42212
A	760	HIS	-	expression tag	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	761	HIS	-	expression tag	UNP P42212
A	762	HIS	-	expression tag	UNP P42212
A	763	HIS	-	expression tag	UNP P42212
B	15	MET	-	initiating methionine	UNP O97467
B	16	GLU	-	expression tag	UNP O97467
B	17	LYS	-	expression tag	UNP O97467
B	18	GLU	-	expression tag	UNP O97467
B	19	ASP	-	expression tag	UNP O97467
B	505	GLU	-	linker	UNP O97467
B	506	ASN	-	linker	UNP O97467
B	507	LEU	-	linker	UNP O97467
B	508	TYR	-	linker	UNP O97467
B	509	PHE	-	linker	UNP O97467
B	510	GLN	-	linker	UNP O97467
B	511	GLY	-	linker	UNP O97467
B	512	GLN	-	linker	UNP O97467
B	513	PHE	-	linker	UNP O97467
B	577	GLY	SER	conflict	UNP P42212
B	584	ALA	SER	conflict	UNP P42212
B	751	ASP	-	expression tag	UNP P42212
B	752	GLU	-	expression tag	UNP P42212
B	753	LEU	-	expression tag	UNP P42212
B	754	TYR	-	expression tag	UNP P42212
B	755	LYS	-	expression tag	UNP P42212
B	756	HIS	-	expression tag	UNP P42212
B	757	HIS	-	expression tag	UNP P42212
B	758	HIS	-	expression tag	UNP P42212
B	759	HIS	-	expression tag	UNP P42212
B	760	HIS	-	expression tag	UNP P42212
B	761	HIS	-	expression tag	UNP P42212
B	762	HIS	-	expression tag	UNP P42212
B	763	HIS	-	expression tag	UNP P42212

- Molecule 2 is 2.5-anhydro-D-mannitol (CCD ID: A1IVP) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	H	O	
2	A	1	23	6	12	5	0
2	B	1	Total	C	H	O	0
			23	6	12	5	



HIS	HIS	GLY	ASP	GLN
HIS	HIS	ASP	PHE	CYS
		GLY	LYS	PHE
		PRO	GLU	ALA
		VAL	ASP	ARG
		LEU	GLY	TYR
		LEU	ASN	PRO
		PRO	ILE	ASP
		ASP	LEU	HIS
		ASN	GLY	MET
		HIS	HIS	LYS
		TYR	LYS	GLN
		LEU	LEU	HIS
		SER	GLU	ASP
		THR	TYR	PHE
		GLN	ASN	PHE
		SER	TYR	LYS
		ALA	ASN	SER
		LEU	SER	ALA
		SER	HIS	MET
		LYS	ASN	PRO
		ASP	VAL	GLU
		PRO	TYR	GLY
		ASN	ILE	TYR
		GLU	MET	VAL
		LYS	ALA	GLN
		ARG	ASP	GLU
		ASP	LYS	ARG
		HIS	GLN	THR
		MET	LYS	ILE
		VAL	ASN	PHE
		LEU	GLY	PHE
		LEU	ILE	LYS
		GLU	LYS	ASP
		PHE	VAL	ASP
		VAL	ASN	GLY
		THR	PHE	ASN
		ALA	LYS	TYR
		ALA	ILE	LYS
		GLY	ARG	THR
		ILE	HIS	ARG
		THR	ASN	ALA
		HIS	ILE	GLU
		GLY	GLU	VAL
		MET	ASP	LYS
		ASP	GLY	PHE
		GLU	SER	GLY
		LEU	VAL	GLY
		TYR	GLN	ASP
		LYS	LEU	THR
		ASP	ALA	LEU
		GLU	ASP	VAL
		LEU	HIS	ASN
		TYR	TYR	ARG
		LYS	GLN	ILE
		HIS	GLU	GLU
		HIS	ASN	LEU
		HIS	THR	LYS
		HIS	PRO	GLY
		HIS	ILE	THR

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	300678	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	62.3	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.430	Depositor
Minimum map value	-0.172	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.075	Depositor
Map size (Å)	267.8, 267.8, 267.8	wwPDB
Map dimensions	412, 412, 412	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.65, 0.65, 0.65	Depositor



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1IVP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.12	0/3884	0.24	0/5263
1	B	0.12	0/3884	0.24	0/5263
All	All	0.12	0/7768	0.24	0/10526

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3791	3912	3911	23	0
1	B	3791	3912	3911	20	0
2	A	11	12	0	0	0
2	B	11	12	0	0	0
All	All	7604	7848	7822	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:SER:OG	1:B:168:HIS:ND1	2.19	0.76
1:A:37:SER:OG	1:A:168:HIS:ND1	2.19	0.75
1:A:142:GLY:O	1:A:145:THR:OG1	2.17	0.62
1:B:142:GLY:O	1:B:145:THR:OG1	2.17	0.62
1:A:234:TYR:OH	1:A:272:ASN:ND2	2.25	0.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/749 (64%)	464 (98%)	12 (2%)	0	100	100
1	B	476/749 (64%)	464 (98%)	12 (2%)	0	100	100
All	All	952/1498 (64%)	928 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/661 (64%)	422 (100%)	2 (0%)	81	90
1	B	424/661 (64%)	422 (100%)	2 (0%)	81	90
All	All	848/1322 (64%)	844 (100%)	4 (0%)	78	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	GLU
1	A	433	LEU
1	B	196	GLU
1	B	433	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	125	HIS
1	B	125	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	A1IVP	A	801	-	11,11,11	2.09	4 (36%)	15,15,15	1.75	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1IVP	B	801	-	11,11,11	2.08	4 (36%)	15,15,15	1.75	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1IVP	A	801	-	-	1/4/20/20	0/1/1/1
2	A1IVP	B	801	-	-	1/4/20/20	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	A1IVP	O11-C03	-4.57	1.34	1.45
2	B	801	A1IVP	O11-C03	-4.57	1.34	1.45
2	A	801	A1IVP	C06-C08	2.75	1.60	1.53
2	B	801	A1IVP	C06-C08	2.75	1.60	1.53
2	A	801	A1IVP	O05-C04	-2.62	1.36	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	A1IVP	C02-C03-C04	-3.21	107.34	115.09
2	A	801	A1IVP	C02-C03-C04	-3.21	107.34	115.09
2	B	801	A1IVP	O11-C03-C04	2.71	110.48	105.11
2	A	801	A1IVP	O11-C03-C04	2.71	110.48	105.11
2	A	801	A1IVP	O05-C04-C03	2.33	117.79	111.05

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	A1IVP	O01-C02-C03-O11
2	B	801	A1IVP	O01-C02-C03-O11


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-54787. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.




## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 206




Y Index: 206




Z Index: 206


### 6.2.2 Raw map



X Index: 206



Y Index: 206




Z Index: 206

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 206




Y Index: 184




Z Index: 243


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.075. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_54787\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 38 nm<sup>3</sup>; this corresponds to an approximate mass of 35 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.413  $\text{\AA}^{-1}$



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.413  $\text{\AA}^{-1}$



## 8.2 Resolution estimates

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.42	-	-
Author-provided FSC curve	2.42	2.89	2.49
Unmasked-calculated*	2.99	3.60	3.06

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.99 differs from the reported value 2.42 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-54787 and PDB model 9SDL. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.075 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.075).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.075) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8200	<div></div> 0.5910
A	<div></div> 0.8220	<div></div> 0.5920
B	<div></div> 0.8230	<div></div> 0.5910

