



wwPDB EM Validation Summary Report ⓘ

Jun 29, 2026 – 03:47 pm BST

PDB ID : 9TIM / pdb_00009tim
EMDB ID : EMD-55961
Title : Phage 812 baseplate in the pre-contraction state - upper arm
Authors : Binovsky, J.; Plevka, P.
Deposited on : 2025-12-05
Resolution : 5.10 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 5.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.
























Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	844 (4.60 - 5.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1019	
2	B	173	
2	C	173	
2	D	173	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	173	 95%
2	F	173	 83% 12%
2	G	173	 17% 91% 8%
2	H	173	 46% 85% 13%
2	I	173	 75% 88% 10%
2	J	173	 73% 91% 6%
2	K	173	 69% 88% 10%
2	L	173	 64% 87% 10%
2	M	173	 81% 90% 9%
3	N	1152	 39% 80% 10% 8%
3	O	1152	 36% 78% 11% 8%
3	P	1152	 39% 82% 9% 8%
3	Q	1152	 11% 80% 10% 8%
3	R	1152	 19% 79% 10% 8%
3	S	1152	 19% 80% 9% 8%
4	T	458	 19% 96%
4	U	458	 23% 97%
4	V	458	 17% 98%
5	W	640	 19% 96%
5	X	640	 14% 95% 5%
5	Y	640	 15% 95% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 97106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ORF63.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	628	5070	3240	807	1013	10	0	0

- Molecule 2 is a protein called ORF64.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	163	1280	818	209	252	1	0	0
2	C	162	1270	811	207	251	1	0	0
2	D	163	1278	817	208	252	1	0	0
2	E	172	1350	858	221	270	1	0	0
2	F	152	1199	769	196	233	1	0	0
2	G	172	1350	858	221	270	1	0	0
2	H	172	1350	858	221	270	1	0	0
2	I	172	1350	858	221	270	1	0	0
2	J	172	1350	858	221	270	1	0	0
2	K	172	1350	858	221	270	1	0	0
2	L	172	1350	858	221	270	1	0	0
2	M	172	1350	858	221	270	1	0	0

- Molecule 3 is a protein called ORF65.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	N	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		
3	O	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		
3	P	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		
3	Q	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		
3	R	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		
3	S	1062	Total	C	N	O	S	0	0
			8364	5260	1398	1682	24		

- Molecule 4 is a protein called ORF68.



Mol	Chain	Residues	Atoms					AltConf	Trace
4	T	458	Total	C	N	O	S	0	0
			3548	2224	592	719	13		
4	U	458	Total	C	N	O	S	0	0
			3548	2224	592	719	13		
4	V	458	Total	C	N	O	S	0	0
			3548	2224	592	719	13		

- Molecule 5 is a protein called CBM-cenC domain-containing protein.



Mol	Chain	Residues	Atoms					AltConf	Trace
5	W	640	Total	C	N	O	S	0	0
			5127	3258	840	1016	13		
5	X	640	Total	C	N	O	S	0	0
			5127	3258	840	1016	13		
5	Y	640	Total	C	N	O	S	0	0
			5127	3258	840	1016	13		

TYR	ASP	LYS	PRO	ARG	ASN	THR	ASP	VAL	ARG	TYR	LYS	LEU	ILE	VAL	GLU	GLY	GLY	PHE	LYS	ASP	ASN	LEU	ILE	VAL	PHE	ARG	VAL	LEU	TYR	ASN	ASN	ASN	ASN	GLU	THR	THR	PHE	ILE	TYR	PRO	SER	SER	GLU	ILE	ILE	ASN	GLY	VAL	ALA	GLU	LYS	GLU	PHE	ILE	ALA	LYS	PRO	SER	THR	GLU		
ASP	LYS	PRO	ARG	ASN	THR	ASP	VAL	ARG	TYR	LYS	LEU	ILE	VAL	ARG	PRO	TYR	ASP	ASP	ASN	LYS	THR	ILE	SER	LYS	VAL	ARG	VAL	LEU	ARG	ASN	LYS	VAL	ASN	ASN	PHE	ASN	GLU	THR	PHE	ILE	TYR	PRO	SER	SER	GLU	ILE	ILE	ASN	GLY	VAL	ALA	GLU	LYS	GLU	PHE	ILE	ALA	LYS	PRO	SER	THR	GLU


● Molecule 2: ORF64




● Molecule 2: ORF64





● Molecule 2: ORF64





● Molecule 2: ORF64




● Molecule 2: ORF64





● Molecule 2: ORF64









• Molecule 2: ORF64




• Molecule 2: ORF64




• Molecule 2: ORF64





• Molecule 2: ORF64




- Molecule 3: ORF65






• Molecule 3: ORF65






• Molecule 3: ORF65






• Molecule 4: ORF68




• Molecule 4: ORF68







- Molecule 4: ORF68




- Molecule 5: CBM-cenC domain-containing protein



- Molecule 5: CBM-cenC domain-containing protein



• Molecule 5: CBM-cenC domain-containing protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	19727	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40.8	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.112	Depositor
Minimum map value	-0.074	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.03	Depositor
Map size (\AA)	600.192, 600.192, 600.192	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.6672001, 1.6672001, 1.6672001	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/5179	1.19	7/7040 (0.1%)
2	B	0.67	0/1306	1.14	2/1774 (0.1%)
2	C	0.66	0/1296	1.17	0/1760
2	D	0.65	0/1304	1.12	0/1771
2	E	0.66	0/1378	1.18	3/1872 (0.2%)
2	F	0.68	0/1224	1.27	4/1660 (0.2%)
2	G	0.73	0/1378	1.33	2/1872 (0.1%)
2	H	0.74	0/1378	1.38	6/1872 (0.3%)
2	I	0.77	0/1378	1.39	6/1872 (0.3%)
2	J	0.77	0/1378	1.37	5/1872 (0.3%)
2	K	0.76	0/1378	1.39	3/1872 (0.2%)
2	L	0.78	0/1378	1.36	4/1872 (0.2%)
2	M	0.79	0/1378	1.35	3/1872 (0.2%)
3	N	0.78	0/8526	1.40	42/11565 (0.4%)
3	O	0.79	0/8526	1.42	54/11565 (0.5%)
3	P	0.79	0/8526	1.33	31/11565 (0.3%)
3	Q	0.75	0/8526	1.38	42/11565 (0.4%)
3	R	0.76	0/8526	1.36	36/11565 (0.3%)
3	S	0.76	0/8526	1.37	45/11565 (0.4%)
4	T	0.75	0/3619	1.25	4/4913 (0.1%)
4	U	0.72	0/3619	1.22	3/4913 (0.1%)
4	V	0.72	0/3619	1.19	4/4913 (0.1%)
5	W	0.73	0/5253	1.22	6/7135 (0.1%)
5	X	0.72	0/5253	1.23	6/7135 (0.1%)
5	Y	0.72	0/5253	1.23	11/7135 (0.2%)
All	All	0.75	0/99105	1.32	329/134515 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
2	B	0	1
2	F	0	1
2	G	0	2
2	H	0	1
2	I	0	2
2	J	0	1
2	L	0	4
2	M	0	2
3	N	0	19
3	O	0	17
3	P	0	17
3	Q	0	21
3	R	0	16
3	S	0	16
4	T	0	2
4	U	0	3
4	V	0	5
5	W	0	2
5	X	0	5
5	Y	0	4
All	All	0	147

There are no bond length outliers.

The worst 5 of 329 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	O	171	ASN	CA-CB-CG	10.40	123.00	112.60
3	S	26	ASN	CA-C-N	8.94	129.84	119.47
3	S	26	ASN	C-N-CA	8.94	129.84	119.47
3	R	779	ASP	CA-CB-CG	8.89	121.50	112.60
3	S	28	ASP	CA-CB-CG	8.59	121.19	112.60

There are no chirality outliers.

5 of 147 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	205	ARG	Sidechain
1	A	293	TYR	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	453	TYR	Sidechain
1	A	468	TYR	Sidechain
1	A	599	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5070	0	4932	12	0
2	B	1280	0	1275	2	0
2	C	1270	0	1263	4	0
2	D	1278	0	1274	0	0
2	E	1350	0	1339	2	0
2	F	1199	0	1195	4	0
2	G	1350	0	1339	2	0
2	H	1350	0	1339	2	0
2	I	1350	0	1339	3	0
2	J	1350	0	1339	4	0
2	K	1350	0	1339	5	0
2	L	1350	0	1339	6	0
2	M	1350	0	1339	1	0
3	N	8364	0	8206	28	0
3	O	8364	0	8206	40	0
3	P	8364	0	8206	23	0
3	Q	8364	0	8206	20	0
3	R	8364	0	8206	28	0
3	S	8364	0	8206	27	0
4	T	3548	0	3468	3	0
4	U	3548	0	3468	3	0
4	V	3548	0	3468	1	0
5	W	5127	0	4969	7	0
5	X	5127	0	4969	6	0
5	Y	5127	0	4969	9	0
All	All	97106	0	95198	222	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 222 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:Q:981:LYS:HA	3:Q:981:LYS:HE2	1.73	0.69
3:O:268:LYS:HB2	3:O:403:TYR:CZ	2.35	0.62
1:A:197:ILE:HD11	2:B:2:ALA:HA	1.81	0.61
3:S:147:LEU:HD23	3:S:148:THR:N	2.16	0.61
3:O:582:ASN:HA	3:O:889:THR:HG22	1.83	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	620/1019 (61%)	590 (95%)	28 (4%)	2 (0%)	36	72
2	B	159/173 (92%)	155 (98%)	4 (2%)	0	100	100
2	C	158/173 (91%)	153 (97%)	4 (2%)	1 (1%)	21	58
2	D	159/173 (92%)	152 (96%)	7 (4%)	0	100	100
2	E	170/173 (98%)	162 (95%)	8 (5%)	0	100	100
2	F	146/173 (84%)	138 (94%)	8 (6%)	0	100	100
2	G	170/173 (98%)	157 (92%)	11 (6%)	2 (1%)	10	42
2	H	170/173 (98%)	152 (89%)	16 (9%)	2 (1%)	10	42
2	I	170/173 (98%)	161 (95%)	8 (5%)	1 (1%)	21	58
2	J	170/173 (98%)	161 (95%)	9 (5%)	0	100	100
2	K	170/173 (98%)	162 (95%)	8 (5%)	0	100	100
2	L	170/173 (98%)	159 (94%)	10 (6%)	1 (1%)	21	58
2	M	170/173 (98%)	161 (95%)	7 (4%)	2 (1%)	10	42
3	N	1056/1152 (92%)	927 (88%)	112 (11%)	17 (2%)	7	36
3	O	1056/1152 (92%)	943 (89%)	96 (9%)	17 (2%)	7	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	P	1056/1152 (92%)	962 (91%)	80 (8%)	14 (1%)	9	40
3	Q	1056/1152 (92%)	943 (89%)	88 (8%)	25 (2%)	4	26
3	R	1056/1152 (92%)	941 (89%)	100 (10%)	15 (1%)	9	38
3	S	1056/1152 (92%)	945 (90%)	99 (9%)	12 (1%)	11	45
4	T	456/458 (100%)	429 (94%)	25 (6%)	2 (0%)	30	67
4	U	456/458 (100%)	439 (96%)	17 (4%)	0	100	100
4	V	456/458 (100%)	438 (96%)	18 (4%)	0	100	100
5	W	638/640 (100%)	611 (96%)	27 (4%)	0	100	100
5	X	638/640 (100%)	601 (94%)	36 (6%)	1 (0%)	43	78
5	Y	638/640 (100%)	598 (94%)	38 (6%)	2 (0%)	36	72
All	All	12220/13301 (92%)	11240 (92%)	864 (7%)	116 (1%)	16	50

5 of 116 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	G	93	ALA
2	H	93	ALA
2	I	93	ALA
2	L	93	ALA
3	N	997	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	573/928 (62%)	568 (99%)	5 (1%)	70	77
2	B	143/153 (94%)	141 (99%)	2 (1%)	59	72
2	C	142/153 (93%)	141 (99%)	1 (1%)	76	80
2	D	143/153 (94%)	143 (100%)	0	100	100
2	E	152/153 (99%)	150 (99%)	2 (1%)	61	73
2	F	134/153 (88%)	132 (98%)	2 (2%)	57	71

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	152/153 (99%)	146 (96%)	6 (4%)	28	49
2	H	152/153 (99%)	138 (91%)	14 (9%)	8	27
2	I	152/153 (99%)	143 (94%)	9 (6%)	18	39
2	J	152/153 (99%)	143 (94%)	9 (6%)	18	39
2	K	152/153 (99%)	141 (93%)	11 (7%)	13	34
2	L	152/153 (99%)	142 (93%)	10 (7%)	15	37
2	M	152/153 (99%)	141 (93%)	11 (7%)	13	34
3	N	934/1010 (92%)	881 (94%)	53 (6%)	18	40
3	O	934/1010 (92%)	868 (93%)	66 (7%)	13	35
3	P	934/1010 (92%)	883 (94%)	51 (6%)	19	41
3	Q	934/1010 (92%)	887 (95%)	47 (5%)	22	43
3	R	934/1010 (92%)	863 (92%)	71 (8%)	12	32
3	S	934/1010 (92%)	867 (93%)	67 (7%)	13	34
4	T	405/405 (100%)	401 (99%)	4 (1%)	68	77
4	U	405/405 (100%)	403 (100%)	2 (0%)	81	82
4	V	405/405 (100%)	404 (100%)	1 (0%)	87	86
5	W	577/577 (100%)	572 (99%)	5 (1%)	70	77
5	X	577/577 (100%)	568 (98%)	9 (2%)	55	69
5	Y	577/577 (100%)	570 (99%)	7 (1%)	63	74
All	All	10901/11770 (93%)	10436 (96%)	465 (4%)	27	47

5 of 465 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	P	588	ILE
4	V	192	ASP
3	Q	876	GLU
4	T	94	MET
3	S	443	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 103 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	257	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	S	332	GLN
5	Y	126	ASN
3	Q	484	ASN
3	R	441	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-55961. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 159




Y Index: 177




Z Index: 179


6.3.2 Raw map



X Index: 159



Y Index: 177




Z Index: 179


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.03. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_55961_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 849 nm³; this corresponds to an approximate mass of 767 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.196 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.196\AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.10	-	-
Author-provided FSC curve	5.10	7.35	5.19
Unmasked-calculated*	7.62	9.91	7.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.62 differs from the reported value 5.1 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-55961 and PDB model 9TIM. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).





















































9.4 Atom inclusion [i](#)



At the recommended contour level, 73% of all backbone atoms, 55% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5490	 0.2680
A	 0.5480	 0.2880
B	 0.7220	 0.3860
C	 0.7280	 0.3890
D	 0.7440	 0.3990
E	 0.7150	 0.3630
F	 0.6980	 0.3360
G	 0.6030	 0.2950
H	 0.4420	 0.2130
I	 0.2840	 0.1570
J	 0.2470	 0.1280
K	 0.2960	 0.1550
L	 0.3210	 0.1500
M	 0.2150	 0.1420
N	 0.4540	 0.2200
O	 0.4720	 0.2190
P	 0.4500	 0.2150
Q	 0.6550	 0.2980
R	 0.6000	 0.2840
S	 0.5960	 0.2850
T	 0.5970	 0.3080
U	 0.5640	 0.2950
V	 0.5960	 0.3200
W	 0.5960	 0.2790
X	 0.6330	 0.3120
Y	 0.6250	 0.2940

