



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 8, 2025 – 12:09 PM JST

PDB ID : 9UDE / pdb_00009ude
Title : Crystal structure of MonCI in complex with diepoxidized farnesyl acetate
Authors : Xiao, H.L.; Guan, Y.Z.; Chen, X.
Deposited on : 2025-04-06
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1003 (2.66-2.66)
Clashscore	180529	1063 (2.66-2.66)
Ramachandran outliers	177936	1052 (2.66-2.66)
Sidechain outliers	177891	1052 (2.66-2.66)
RSRZ outliers	164620	1003 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	 3% 84% 9% 8%
1	B	511	 5% 77% 14% 8%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MonCI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	472	3547	2219	645	673	10	0	1	0
1	B	472	3503	2196	638	660	9	0	0	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	initiating methionine	UNP Q846W9
A	-13	ASN	-	expression tag	UNP Q846W9
A	-12	HIS	-	expression tag	UNP Q846W9
A	-11	LYS	-	expression tag	UNP Q846W9
A	-10	VAL	-	expression tag	UNP Q846W9
A	-9	HIS	-	expression tag	UNP Q846W9
A	-8	HIS	-	expression tag	UNP Q846W9
A	-7	HIS	-	expression tag	UNP Q846W9
A	-6	HIS	-	expression tag	UNP Q846W9
A	-5	HIS	-	expression tag	UNP Q846W9
A	-4	HIS	-	expression tag	UNP Q846W9
A	-3	ILE	-	expression tag	UNP Q846W9
A	-2	GLU	-	expression tag	UNP Q846W9
A	-1	GLY	-	expression tag	UNP Q846W9
A	0	ARG	-	expression tag	UNP Q846W9
B	-14	MET	-	initiating methionine	UNP Q846W9
B	-13	ASN	-	expression tag	UNP Q846W9
B	-12	HIS	-	expression tag	UNP Q846W9
B	-11	LYS	-	expression tag	UNP Q846W9
B	-10	VAL	-	expression tag	UNP Q846W9
B	-9	HIS	-	expression tag	UNP Q846W9
B	-8	HIS	-	expression tag	UNP Q846W9
B	-7	HIS	-	expression tag	UNP Q846W9
B	-6	HIS	-	expression tag	UNP Q846W9
B	-5	HIS	-	expression tag	UNP Q846W9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	HIS	-	expression tag	UNP Q846W9
B	-3	ILE	-	expression tag	UNP Q846W9
B	-2	GLU	-	expression tag	UNP Q846W9
B	-1	GLY	-	expression tag	UNP Q846W9
B	0	ARG	-	expression tag	UNP Q846W9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is diepoxidized farnesyl acetate (CCD ID: YK6) (formula: $C_{17}H_{28}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			21	17	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	45	Total O 45 45	0	0
6	B	8	Total O 8 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MonCI



- Molecule 1: MonCI



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.62Å 52.77Å 145.83Å 90.00° 92.37° 90.00°	Depositor
Resolution (Å)	145.70 – 2.65 145.70 – 2.65	Depositor EDS
% Data completeness (in resolution range)	95.7 (145.70-2.65) 95.7 (145.70-2.65)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 2.65Å)	Xtrriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.199 , 0.261 0.203 , 0.262	Depositor DCC
R_{free} test set	1483 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	46.5	Xtrriage
Anisotropy	0.491	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7237	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, YK6, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3620	0.49	0/4939
1	B	0.28	0/3573	0.50	0/4881
All	All	0.28	0/7193	0.50	0/9820

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3547	0	3502	30	0
1	B	3503	0	3443	55	0
2	A	53	0	31	2	0
2	B	53	0	31	1	0
3	A	21	0	0	0	0
4	A	6	0	8	0	0
5	A	1	0	0	0	0
6	A	45	0	0	0	0
6	B	8	0	0	0	0
All	All	7237	0	7015	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:HIS:CE1	1:B:382:LEU:HD23	1.85	1.10
1:B:334:HIS:CE1	1:B:382:LEU:CD2	2.56	0.88
1:A:233:ARG:HH21	1:A:411:LEU:HD23	1.39	0.87
1:B:195:PRO:HG2	1:B:311:LEU:HD11	1.61	0.81
1:B:334:HIS:HE1	1:B:382:LEU:HD23	1.47	0.80
1:B:176:ILE:HD13	1:B:319:LEU:HB2	1.68	0.74
1:A:469:PRO:HG2	1:A:470:GLU:OE2	1.88	0.73
1:B:4:THR:HG22	1:B:5:ARG:H	1.56	0.71
1:B:36:GLU:HG2	1:B:40:LEU:HD21	1.73	0.70
1:B:211:ARG:HG3	1:B:295:VAL:HG12	1.79	0.64
1:B:212:LEU:HD21	1:B:250:ARG:HD2	1.79	0.64
1:B:141:VAL:O	1:B:188:TRP:NE1	2.32	0.62
1:A:407:THR:O	1:A:411:LEU:HD12	2.00	0.61
1:A:4:THR:OG1	1:A:5:ARG:N	2.35	0.60
1:B:189:LEU:HD22	1:B:194:VAL:HG21	1.82	0.59
1:B:309:GLU:HB2	1:B:370:LYS:HG2	1.85	0.59
1:B:230:ALA:H	1:B:240:PHE:HD1	1.52	0.58
1:B:38:ASP:HA	1:B:137:ARG:NH2	2.21	0.55
1:A:233:ARG:HG2	1:A:390:TYR:CE1	2.43	0.54
1:A:208:TYR:CZ	1:A:330:PRO:HG2	2.43	0.53
1:B:334:HIS:HE1	1:B:382:LEU:CD2	2.12	0.53
1:A:468:LEU:HD23	1:A:470:GLU:OE1	2.08	0.53
1:A:23:HIS:HB2	1:A:126:ALA:HB1	1.92	0.52
1:A:37:ARG:HD3	1:A:139:GLU:HG3	1.92	0.52
1:B:202:VAL:HG22	1:B:395:VAL:HG23	1.90	0.52
1:B:208:TYR:CE1	1:B:330:PRO:HG2	2.44	0.52
1:B:381:ILE:HD13	1:B:401:ARG:CZ	2.39	0.52
1:B:32:VAL:O	1:B:132:ILE:HA	2.10	0.51
1:B:95:LEU:HB2	1:B:100:TRP:CZ3	2.46	0.50
1:A:5:ARG:HH21	1:A:5:ARG:HG2	1.77	0.50
1:A:233:ARG:HG2	1:A:390:TYR:CZ	2.46	0.50
1:B:38:ASP:OD2	1:B:49:GLY:N	2.45	0.50
1:B:179:THR:OG1	1:B:183:SER:HB2	2.11	0.50
1:A:208:TYR:OH	1:A:331:ILE:HG13	2.12	0.49
1:B:206:VAL:HB	1:B:301:GLY:HA3	1.94	0.49
1:B:233:ARG:HG3	1:B:390:TYR:CE1	2.47	0.49
1:B:161:ASP:HB3	1:B:164:SER:HG	1.77	0.49
1:B:208:TYR:CZ	1:B:330:PRO:HG2	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:259:ARG:NH1	1:B:392:ASN:O	2.44	0.49
1:B:472:LEU:HD22	1:B:477:LEU:HB2	1.95	0.49
1:B:10:VAL:HG23	1:B:172:ALA:HB2	1.95	0.48
1:A:242:VAL:HG11	1:A:244:TYR:CE2	2.48	0.48
1:B:177:ASP:OD1	1:B:179:THR:OG1	2.26	0.48
1:A:92:LEU:HG	1:A:225:ALA:HB3	1.95	0.48
1:A:338:SER:HB2	1:A:379:PRO:HG3	1.95	0.48
1:B:188:TRP:O	1:B:192:LEU:HD13	2.14	0.48
1:B:407:THR:O	1:B:411:LEU:HD12	2.14	0.48
1:A:34:VAL:HB	1:A:134:VAL:HG12	1.94	0.47
1:B:4:THR:HG22	1:B:5:ARG:HG3	1.96	0.47
1:B:232:ASP:OD1	1:B:233:ARG:N	2.48	0.47
1:B:258:THR:HG21	1:B:390:TYR:CE1	2.49	0.47
1:B:353:SER:C	1:B:355:GLN:H	2.18	0.47
1:A:243:VAL:HG23	1:A:282:LEU:HD22	1.97	0.46
1:A:208:TYR:CE1	1:A:330:PRO:HG2	2.50	0.46
1:B:38:ASP:HA	1:B:137:ARG:HH21	1.79	0.46
1:A:37:ARG:HG3	1:A:162:LEU:HD11	1.98	0.46
1:B:382:LEU:HD11	1:B:434:VAL:HG12	1.97	0.46
1:B:382:LEU:HD12	1:B:441:GLN:HG3	1.98	0.46
1:A:95:LEU:HB2	1:A:100:TRP:CZ3	2.51	0.46
1:B:208:TYR:OH	1:B:331:ILE:HG13	2.16	0.45
1:A:319:LEU:HD11	1:A:368:LEU:HD13	1.98	0.45
1:A:258:THR:HG21	1:A:390:TYR:CE1	2.52	0.45
1:B:259:ARG:HD2	1:B:392:ASN:HB3	1.99	0.45
1:B:212:LEU:HD13	1:B:252:LEU:HD13	1.98	0.45
1:B:385:THR:HG21	1:B:441:GLN:HE22	1.82	0.45
1:B:211:ARG:CG	1:B:295:VAL:HG12	2.47	0.44
1:B:181:ARG:HG2	1:B:304:ARG:NH2	2.33	0.44
1:A:470:GLU:H	1:A:470:GLU:CD	2.20	0.43
1:B:144:THR:HG21	1:B:169:GLN:HE21	1.82	0.43
1:A:55:HIS:NE2	1:A:298:SER:HB2	2.34	0.43
1:B:161:ASP:HB3	1:B:164:SER:OG	2.18	0.43
1:B:189:LEU:HB3	1:B:194:VAL:HG23	2.01	0.43
1:B:112:VAL:HG21	1:B:244:TYR:HB3	1.99	0.43
1:B:57:HIS:HA	2:B:700:FAD:O4	2.19	0.42
1:B:247:GLU:OE1	1:B:250:ARG:NH2	2.53	0.42
1:A:242:VAL:HG11	1:A:244:TYR:CZ	2.55	0.41
1:B:17:ALA:HB1	1:B:321:ILE:HD12	2.01	0.41
1:A:335:GLY:HA3	2:A:501:FAD:H1'2	2.03	0.41
1:B:243:VAL:HG23	1:B:282:LEU:HD22	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:LEU:HG	1:B:225:ALA:HB3	2.02	0.41
1:B:381:ILE:HD13	1:B:401:ARG:NE	2.36	0.41
1:A:446:SER:O	1:A:450:LEU:HG	2.21	0.40
1:A:472:LEU:HD13	1:A:478:ASP:HA	2.03	0.40
1:A:206:VAL:HA	1:A:257:CYS:O	2.22	0.40
1:A:57:HIS:HA	2:A:501:FAD:O4	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	469/511 (92%)	454 (97%)	15 (3%)	0	100	100
1	B	468/511 (92%)	452 (97%)	15 (3%)	1 (0%)	44	61
All	All	937/1022 (92%)	906 (97%)	30 (3%)	1 (0%)	48	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	354	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/401 (90%)	359 (99%)	2 (1%)	84	92
1	B	351/401 (88%)	346 (99%)	5 (1%)	62	79
All	All	712/802 (89%)	705 (99%)	7 (1%)	73	85

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	160	ARG
1	A	304	ARG
1	B	135	ARG
1	B	183	SER
1	B	232	ASP
1	B	233	ARG
1	B	355	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	355	GLN
1	B	169	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	YK6	A	502	-	22,22,22	2.87	5 (22%)	33,33,33	3.63	9 (27%)
2	FAD	B	700	-	53,58,58	0.45	0	68,89,89	0.49	1 (1%)
4	GOL	A	503	-	5,5,5	0.90	0	5,5,5	0.99	0
2	FAD	A	501	-	53,58,58	0.46	0	68,89,89	0.50	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	YK6	A	502	-	-	13/17/33/33	0/2/2/2
2	FAD	B	700	-	-	5/30/50/50	0/6/6/6
4	GOL	A	503	-	-	0/4/4/4	-
2	FAD	A	501	-	-	5/30/50/50	0/6/6/6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	YK6	O21-C10	9.59	1.60	1.45
3	A	502	YK6	O20-C14	8.35	1.60	1.46
3	A	502	YK6	C10-C09	-2.15	1.44	1.47
3	A	502	YK6	O20-C13	-2.11	1.41	1.45
3	A	502	YK6	O21-C09	-2.07	1.41	1.45

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	YK6	O21-C09-C10	13.49	68.52	59.86
3	A	502	YK6	O20-C13-C14	10.91	68.51	59.91
3	A	502	YK6	O21-C10-C09	-5.57	54.84	59.26
3	A	502	YK6	C10-O21-C09	-5.42	56.64	60.88
3	A	502	YK6	O20-C13-C12	-4.16	110.61	117.74
3	A	502	YK6	C14-O20-C13	-3.96	56.76	60.77
3	A	502	YK6	O20-C14-C13	-3.58	54.73	58.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	YK6	O21-C09-C08	-3.01	112.59	117.74
3	A	502	YK6	C08-C09-C10	-2.44	121.74	125.35
2	A	501	FAD	C5A-C6A-N6A	2.32	123.88	120.35
2	B	700	FAD	C5A-C6A-N6A	2.28	123.81	120.35

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FAD	C5'-O5'-P-O3P
2	B	700	FAD	C5'-O5'-P-O3P
3	A	502	YK6	C01-C02-O03-C04
3	A	502	YK6	O16-C02-O03-C04
3	A	502	YK6	C06-C07-C08-C09
3	A	502	YK6	C17-C06-C07-C08
3	A	502	YK6	C05-C06-C07-C08
3	A	502	YK6	C10-C11-C12-C13
2	A	501	FAD	PA-O3P-P-O5'
2	B	700	FAD	PA-O3P-P-O5'
3	A	502	YK6	C07-C08-C09-C10
2	A	501	FAD	C5'-O5'-P-O2P
2	B	700	FAD	C5'-O5'-P-O2P
3	A	502	YK6	C11-C12-C13-C14
3	A	502	YK6	C18-C10-C11-C12
3	A	502	YK6	O21-C10-C11-C12
3	A	502	YK6	C07-C08-C09-O21
3	A	502	YK6	C11-C12-C13-O20
2	A	501	FAD	P-O3P-PA-O2A
2	B	700	FAD	P-O3P-PA-O2A
3	A	502	YK6	C09-C10-C11-C12
2	A	501	FAD	O4B-C4B-C5B-O5B
2	B	700	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.



2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	700	FAD	1	0
2	A	501	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	472/511 (92%)	-0.11	13 (2%) 55 53	20, 39, 67, 97	1 (0%)
1	B	472/511 (92%)	0.53	28 (5%) 29 27	35, 63, 96, 116	0
All	All	944/1022 (92%)	0.21	41 (4%) 40 37	20, 49, 90, 116	1 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	480	ALA	7.7
1	A	480	ALA	5.5
1	B	383	ALA	5.5
1	B	231	ASP	4.6
1	B	382	LEU	4.6
1	B	208	TYR	4.3
1	A	479	ALA	4.1
1	A	355	GLN	3.8
1	B	363	ALA	3.7
1	B	479	ALA	3.6
1	B	151	GLY	3.3
1	A	362	ARG	3.2
1	B	361	ALA	3.1
1	B	362	ARG	3.0
1	B	230	ALA	3.0
1	A	361	ALA	2.9
1	B	4	THR	2.9
1	B	354	VAL	2.9
1	B	150	SER	2.8
1	B	451	MET	2.8
1	A	363	ALA	2.7
1	B	158	VAL	2.7
1	B	168	GLU	2.6
1	B	166	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	355	GLN	2.5
1	B	147	GLY	2.5
1	B	143	LEU	2.4
1	A	164	SER	2.4
1	A	478	ASP	2.4
1	A	4	THR	2.3
1	A	149	GLY	2.3
1	B	35	VAL	2.3
1	B	312	GLU	2.3
1	B	317	GLY	2.3
1	A	364	GLY	2.2
1	B	5	ARG	2.2
1	B	229	ALA	2.2
1	B	165	GLY	2.1
1	A	208	TYR	2.0
1	B	152	GLY	2.0
1	A	150	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	YK6	A	502	21/21	0.67	0.29	58,83,93,99	0
4	GOL	A	503	6/6	0.87	0.13	55,58,69,75	0
2	FAD	B	700	53/53	0.95	0.08	41,53,71,81	0
2	FAD	A	501	53/53	0.96	0.08	21,31,48,58	0
5	CL	A	504	1/1	0.97	0.10	42,42,42,42	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









Electron density around FAD B 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.