



# wwPDB X-ray Structure Validation Summary Report

Mar 28, 2026 – 04:49 PM UTC

PDB ID : 8URO / pdb\_00008uro  
Title : Crystal structure of IgG1-Fc fragment (E382S) in complex with Corynebacterial ENGase CU43 (D187A-E189A)  
Authors : Sastre, D.E.; Sultana, N.; Sundberg, E.J.  
Deposited on : 2023-10-26  
Resolution : 3.62 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1062 (3.72-3.52)
Clashscore	190562	1092 (3.72-3.52)
Ramachandran outliers	187476	1057 (3.72-3.52)
Sidechain outliers	187428	1055 (3.72-3.52)
RSRZ outliers	180081	1060 (3.72-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	 7% 70% 10% 20%
1	D	415	 5% 71% 9% 20%
2	B	232	 12% 80% 7% 12%
2	C	232	 22% 63% 15% 34%
2	E	232	 19% 82% 5% 12%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	232	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '34%', a green segment in the middle labeled '64%', and a grey segment on the right labeled '31%'. There are two small black dots on the grey segment.</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Corynebacterial protease CP40.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2328	1460	405	456	7	0	0	0
1	D	332	2399	1506	418	469	6	0	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP A0A830QWM5
A	-1	GLY	-	expression tag	UNP A0A830QWM5
A	0	SER	-	expression tag	UNP A0A830QWM5
A	1	SER	-	expression tag	UNP A0A830QWM5
A	2	HIS	-	expression tag	UNP A0A830QWM5
A	3	HIS	-	expression tag	UNP A0A830QWM5
A	4	HIS	-	expression tag	UNP A0A830QWM5
A	5	HIS	-	expression tag	UNP A0A830QWM5
A	6	HIS	-	expression tag	UNP A0A830QWM5
A	7	HIS	-	expression tag	UNP A0A830QWM5
A	8	SER	-	expression tag	UNP A0A830QWM5
A	9	SER	-	expression tag	UNP A0A830QWM5
A	10	GLY	-	expression tag	UNP A0A830QWM5
A	11	LEU	-	expression tag	UNP A0A830QWM5
A	12	VAL	-	expression tag	UNP A0A830QWM5
A	13	PRO	-	expression tag	UNP A0A830QWM5
A	14	ARG	-	expression tag	UNP A0A830QWM5
A	15	GLY	-	expression tag	UNP A0A830QWM5
A	16	SER	-	expression tag	UNP A0A830QWM5
A	17	HIS	-	expression tag	UNP A0A830QWM5
A	18	MET	-	expression tag	UNP A0A830QWM5
A	19	ALA	-	expression tag	UNP A0A830QWM5
A	20	SER	-	expression tag	UNP A0A830QWM5
A	21	MET	-	expression tag	UNP A0A830QWM5
A	22	THR	-	expression tag	UNP A0A830QWM5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLY	-	expression tag	UNP A0A830QWM5
A	24	GLY	-	expression tag	UNP A0A830QWM5
A	25	GLN	-	expression tag	UNP A0A830QWM5
A	26	GLN	-	expression tag	UNP A0A830QWM5
A	27	MET	-	expression tag	UNP A0A830QWM5
A	28	GLY	-	expression tag	UNP A0A830QWM5
A	29	ARG	-	expression tag	UNP A0A830QWM5
A	30	GLY	-	expression tag	UNP A0A830QWM5
A	31	SER	-	expression tag	UNP A0A830QWM5
A	187	ALA	ASP	engineered mutation	UNP A0A830QWM5
A	189	ALA	GLU	engineered mutation	UNP A0A830QWM5
D	-2	MET	-	expression tag	UNP A0A830QWM5
D	-1	GLY	-	expression tag	UNP A0A830QWM5
D	0	SER	-	expression tag	UNP A0A830QWM5
D	1	SER	-	expression tag	UNP A0A830QWM5
D	2	HIS	-	expression tag	UNP A0A830QWM5
D	3	HIS	-	expression tag	UNP A0A830QWM5
D	4	HIS	-	expression tag	UNP A0A830QWM5
D	5	HIS	-	expression tag	UNP A0A830QWM5
D	6	HIS	-	expression tag	UNP A0A830QWM5
D	7	HIS	-	expression tag	UNP A0A830QWM5
D	8	SER	-	expression tag	UNP A0A830QWM5
D	9	SER	-	expression tag	UNP A0A830QWM5
D	10	GLY	-	expression tag	UNP A0A830QWM5
D	11	LEU	-	expression tag	UNP A0A830QWM5
D	12	VAL	-	expression tag	UNP A0A830QWM5
D	13	PRO	-	expression tag	UNP A0A830QWM5
D	14	ARG	-	expression tag	UNP A0A830QWM5
D	15	GLY	-	expression tag	UNP A0A830QWM5
D	16	SER	-	expression tag	UNP A0A830QWM5
D	17	HIS	-	expression tag	UNP A0A830QWM5
D	18	MET	-	expression tag	UNP A0A830QWM5
D	19	ALA	-	expression tag	UNP A0A830QWM5
D	20	SER	-	expression tag	UNP A0A830QWM5
D	21	MET	-	expression tag	UNP A0A830QWM5
D	22	THR	-	expression tag	UNP A0A830QWM5
D	23	GLY	-	expression tag	UNP A0A830QWM5
D	24	GLY	-	expression tag	UNP A0A830QWM5
D	25	GLN	-	expression tag	UNP A0A830QWM5
D	26	GLN	-	expression tag	UNP A0A830QWM5
D	27	MET	-	expression tag	UNP A0A830QWM5
D	28	GLY	-	expression tag	UNP A0A830QWM5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	ARG	-	expression tag	UNP A0A830QWM5
D	30	GLY	-	expression tag	UNP A0A830QWM5
D	31	SER	-	expression tag	UNP A0A830QWM5
D	187	ALA	ASP	engineered mutation	UNP A0A830QWM5
D	189	ALA	GLU	engineered mutation	UNP A0A830QWM5


- Molecule 2 is a protein called Immunoglobulin gamma-1 heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	B	204	Total	C	N	O	S	0	0	0
			1401	899	233	263	6			
2	C	152	Total	C	N	O	S	0	0	0
			893	554	162	175	2			
2	E	203	Total	C	N	O	S	0	0	0
			1317	839	228	244	6			
2	F	160	Total	C	N	O	S	0	0	0
			852	512	163	173	4			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	382	SER	GLU	engineered mutation	UNP P0DOX5
C	382	SER	GLU	engineered mutation	UNP P0DOX5
E	382	SER	GLU	engineered mutation	UNP P0DOX5
F	382	SER	GLU	engineered mutation	UNP P0DOX5








• Molecule 2: Immunoglobulin gamma-1 heavy chain




• Molecule 2: Immunoglobulin gamma-1 heavy chain



• Molecule 2: Immunoglobulin gamma-1 heavy chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.44Å 81.99Å 148.10Å 90.00° 113.20° 90.00°	Depositor
Resolution (Å)	33.96 – 3.62 33.96 – 3.62	Depositor EDS
% Data completeness (in resolution range)	95.6 (33.96-3.62) 95.4 (33.96-3.62)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 3.65Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.334 , 0.369 0.334 , 0.369	Depositor DCC
$R_{free}$ test set	1949 reflections (6.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	96.4	Xtrriage
Anisotropy	0.491	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 124.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.049 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.78	EDS
Total number of atoms	9190	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	108.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.09	0/2377	0.29	0/3254
1	D	0.09	0/2447	0.28	0/3340
2	B	0.12	0/1438	0.38	0/1985
2	C	0.10	0/897	0.30	0/1227
2	E	0.10	0/1348	0.31	0/1867
2	F	0.14	0/837	0.33	0/1132
All	All	0.10	0/9344	0.31	0/12805

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	1949	23	0
1	D	2399	0	2099	24	0
2	B	1401	0	1158	8	0
2	C	893	0	567	2	0
2	E	1317	0	1044	7	0
2	F	852	0	430	3	0
All	All	9190	0	7247	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:120:PHE:HA	1:D:124:TYR:HB2	1.70	0.73
1:A:120:PHE:HA	1:A:124:TYR:HB2	1.72	0.72
1:D:66:TYR:HB2	1:D:332:TYR:HA	1.76	0.67
1:A:66:TYR:HD2	1:A:332:TYR:HD1	1.41	0.67
1:D:191:ARG:HG2	1:D:244:LEU:HD22	1.79	0.64

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/415 (79%)	310 (94%)	14 (4%)	4 (1%)	10	39
1	D	326/415 (79%)	312 (96%)	14 (4%)	0	100	100
2	B	196/232 (84%)	180 (92%)	9 (5%)	7 (4%)	2	20
2	C	121/232 (52%)	114 (94%)	5 (4%)	2 (2%)	7	32
2	E	197/232 (85%)	192 (98%)	2 (1%)	3 (2%)	8	34
2	F	115/232 (50%)	104 (90%)	5 (4%)	6 (5%)	1	14
All	All	1283/1758 (73%)	1212 (94%)	49 (4%)	22 (2%)	7	32

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	SER
1	A	45	PRO
2	B	291	PRO
2	B	297	ASN
2	B	374	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/338 (56%)	188 (100%)	1 (0%)	81	78
1	D	211/338 (62%)	211 (100%)	0	100	100
2	B	124/215 (58%)	124 (100%)	0	100	100
2	C	49/215 (23%)	49 (100%)	0	100	100
2	E	102/215 (47%)	102 (100%)	0	100	100
2	F	27/215 (13%)	26 (96%)	1 (4%)	30	52
All	All	702/1536 (46%)	700 (100%)	2 (0%)	86	81

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	CYS
2	F	425	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	GLN
1	A	246	GLN
1	A	292	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	332/415 (80%)	0.83	28 (8%) 17 13	56, 87, 128, 164	0
1	D	332/415 (80%)	0.76	19 (5%) 29 19	57, 86, 134, 154	0
2	B	204/232 (87%)	1.08	29 (14%) 6 7	59, 98, 147, 171	0
2	C	152/232 (65%)	1.68	50 (32%) 1 1	110, 163, 202, 207	0
2	E	203/232 (87%)	1.41	43 (21%) 2 3	65, 122, 177, 225	0
2	F	160/232 (68%)	2.12	79 (49%) 0 0	89, 170, 194, 208	0
All	All	1383/1758 (78%)	1.18	248 (17%) 3 4	56, 99, 184, 225	0

The worst 5 of 248 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	280	ASP	7.0
1	D	341	TYR	6.2
2	E	294	GLU	5.8
2	F	305	VAL	5.7
2	C	380	GLU	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.