



Full wwPDB X-ray Structure Validation Report i

Oct 17, 2024 – 04:10 PM EDT

PDB ID : 8W1X

Title : 2.35-angstrom resolution intermediate crystal structure of KatG from Mycobacterium tuberculosis with an MYW-OOH cofactor soaked with peroxide for 5 minutes

Authors : Liu, A.; Li, J.

Deposited on : 2024-02-19

Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1460 (2.36-2.36)
Clashscore	180529	1571 (2.36-2.36)
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 8 unique types of molecules in this entry. The entry contains 11790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase-peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	717	5529	3515	953	1042	19	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A0A0D5ZBI4
A	1	HIS	-	expression tag	UNP A0A0D5ZBI4


- Molecule 2 is a protein called Catalase-peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	717	5531	3515	953	1044	19	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP A0A0D5ZBI4
B	1	HIS	-	expression tag	UNP A0A0D5ZBI4

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total		C	Fe	N	O	
			43		34	1	4	4	0 0
3	B	1	Total		C	Fe	N	O	
			43		34	1	4	4	0 0

- Molecule 4 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			2	2		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	245	Total O 245 245	0	0
8	B	376	Total O 376 376	0	0

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	150.46 Å 150.46 Å 155.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 2.35 47.72 – 2.35	Depositor EDS
% Data completeness (in resolution range)	90.3 (47.72-2.35) 82.9 (47.72-2.35)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.30 (at 2.34 Å)	Xtriage
Refinement program	PHENIX 1.21_5207	Depositor
R , R_{free}	0.191 , 0.256 0.191 , 0.255	Depositor DCC
R_{free} test set	65480 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.027 for -h,l,k 0.018 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11790	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TOX, GOL, NA, OXY, HEM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/5680	0.63	0/7732
2	B	0.50	0/5663	0.65	0/7706
All	All	0.49	0/11343	0.64	0/15438

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5529	0	5355	88	0
2	B	5531	0	5354	83	0
3	A	43	0	30	3	0
3	B	43	0	30	2	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	4	0	3	0	0
6	B	4	0	3	0	0
7	B	12	0	16	2	0
8	A	245	0	0	5	0
8	B	376	0	0	7	0
All	All	11790	0	10791	167	0

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:577:GLU:H	2:B:577:GLU:CD	2.11	0.54
2:B:366:ASP:HB3	2:B:370:GLY:HA3	1.90	0.53
1:A:107:TRP:HZ3	1:A:229:TYR:HH	1.54	0.53
1:A:498:ARG:HD3	1:A:540:PHE:CD2	2.44	0.53
1:A:420:MET:O	1:A:425:ARG:HD3	2.09	0.52
2:B:46:LYS:HG2	2:B:49:HIS:CE1	2.44	0.52
1:A:239:PRO:O	1:A:241:PRO:HD3	2.10	0.52
1:A:504:GLY:HA2	1:A:510:PRO:HG3	1.92	0.52
1:A:43:LEU:HG	1:A:611:LEU:HD11	1.90	0.51
1:A:458:LEU:O	1:A:462:ILE:CG1	2.55	0.51
1:A:692:SER:N	1:A:695:ASP:OD2	2.37	0.51
1:A:459:LYS:HG2	1:A:550:ALA:HB2	1.93	0.51
2:B:533:PRO:HG2	2:B:536:ILE:HD12	1.93	0.51
1:A:694:VAL:O	1:A:697:VAL:HG12	2.11	0.51
1:A:450:VAL:CG1	1:A:538:VAL:HG12	2.41	0.51
1:A:502:GLN:HG2	1:A:505:TRP:CZ3	2.46	0.50
1:A:598:LEU:HD22	1:A:606:ALA:HB2	1.94	0.50
2:B:366:ASP:HB2	8:B:1126:HOH:O	2.10	0.50
2:B:46:LYS:HG2	2:B:49:HIS:NE2	2.27	0.50
1:A:27:LYS:NZ	2:B:40:PRO:O	2.42	0.50
2:B:463:ARG:HD3	2:B:554:LYS:HB2	1.94	0.50
2:B:41:ASN:O	2:B:608:TYR:OH	2.28	0.49
1:A:195:GLU:HA	1:A:195:GLU:OE2	2.12	0.49
1:A:463:ARG:HH21	1:A:554:LYS:HD2	1.76	0.49
1:A:78:ARG:NH1	8:A:907:HOH:O	2.35	0.49
2:B:274:LYS:CB	2:B:320:VAL:HG22	2.43	0.48
2:B:174:GLU:HA	2:B:178:PHE:O	2.13	0.48
1:A:450:VAL:HG11	1:A:538:VAL:HG12	1.94	0.48
2:B:509:ASP:CG	2:B:509:ASP:O	2.50	0.48
1:A:239:PRO:HG3	1:A:351:TRP:CG	2.48	0.48
2:B:364:ILE:HD12	2:B:375:PRO:HA	1.96	0.48
1:A:498:ARG:HG3	1:A:499:LEU:HD23	1.95	0.48
2:B:139:ALA:HA	2:B:300:TRP:CH2	2.48	0.48
1:A:542:ASP:OD1	1:A:571:ARG:HB2	2.14	0.48
2:B:532:ALA:HB1	2:B:533:PRO:HD2	1.95	0.47
2:B:139:ALA:HA	2:B:300:TRP:CZ3	2.49	0.47
2:B:694:VAL:O	2:B:697:VAL:HG12	2.14	0.47
1:A:443:PRO:HB2	1:A:572:THR:CG2	2.44	0.47
1:A:673:ALA:HB3	1:A:675:ASP:OD1	2.15	0.47
2:B:731:VAL:HA	2:B:734:LEU:HG	1.96	0.47
2:B:279:GLY:O	2:B:348:ALA:HB2	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:500:GLN:HG3	1:A:501:PRO:HA	2.03	0.40
1:A:529:ASN:O	1:A:532:ALA:CB	2.69	0.40
2:B:149:TRP:NE1	2:B:153:LYS:HD2	2.35	0.40
2:B:471:GLN:HE21	2:B:513:ASP:CG	2.24	0.40
1:A:216:LEU:HB3	1:A:254:ARG:CZ	2.51	0.40
1:A:320:VAL:HB	1:A:364:ILE:HG21	2.03	0.40
2:B:174:GLU:HG2	2:B:178:PHE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	715/741 (96%)	688 (96%)	27 (4%)	0	100 100
2	B	714/741 (96%)	691 (97%)	22 (3%)	1 (0%)	48 59
All	All	1429/1482 (96%)	1379 (96%)	49 (3%)	1 (0%)	48 59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	510	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	565/584 (97%)	551 (98%)	14 (2%)	42 53
2	B	564/583 (97%)	551 (98%)	13 (2%)	45 56
All	All	1129/1167 (97%)	1102 (98%)	27 (2%)	44 55


All (27) residues with a non-rotameric sidechain are listed below:


Mol	Chain	Res	Type
1	A	42	ARG
1	A	54	VAL
1	A	137	ASP
1	A	143	LYS
1	A	253	ARG
1	A	392	ARG
1	A	416	ILE
1	A	435	THR
1	A	458	LEU
1	A	462	ILE
1	A	465	SER
1	A	481	SER
1	A	513	ASP
1	A	675	ASP
2	B	47	VAL
2	B	54	VAL
2	B	68	VAL
2	B	82	GLU
2	B	137	ASP
2	B	143	LYS
2	B	301	LYS
2	B	416	ILE
2	B	426	TYR
2	B	442	VAL
2	B	511	ASP
2	B	557	LYS
2	B	626	VAL


Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	596	ASN
1	A	660	ASN
2	B	722	GLN

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	717/741 (96%)	0.10	10 (1%) 73 77	16, 33, 57, 79	0
2	B	716/741 (96%)	-0.29	5 (0%) 84 86	14, 26, 43, 69	0
All	All	1433/1482 (96%)	-0.10	15 (1%) 79 83	14, 30, 52, 79	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	510	PRO	4.6
1	A	467	LEU	3.4
1	A	465	SER	3.1
1	A	464	ALA	2.8
1	A	524	ILE	2.8
2	B	511	ASP	2.7
1	A	24	GLY	2.6
1	A	533	PRO	2.6
2	B	24	GLY	2.5
1	A	512	GLY	2.5
1	A	536	ILE	2.3
2	B	371	PRO	2.2
1	A	511	ASP	2.2
2	B	213	LYS	2.2
1	A	316	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TOX	B	107	16/17	0.95	0.08	18,21,29,31	0

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	B	803	6/6	0.89	0.13	25,26,34,36	0
6	ACT	B	804	4/4	0.90	0.11	20,26,34,40	0
6	ACT	A	804	4/4	0.90	0.10	22,29,34,37	0
7	GOL	B	802	6/6	0.91	0.10	22,27,32,34	0
4	OXY	A	802	2/2	0.92	0.11	34,34,34,39	0
5	NA	A	803	1/1	0.95	0.05	23,23,23,23	0
3	HEM	B	801	43/43	0.96	0.09	17,22,29,40	0
3	HEM	A	801	43/43	0.97	0.07	19,25,30,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.