



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:48 PM UTC

PDB ID : 9W8G / pdb\_00009w8g  
Title : Crystal structure of Staphylococcus aureus cysteine-free ScdA with bound iron, determined by molecular replacement and Fe anomalous signal  
Authors : Yang, W.K.  
Deposited on : 2025-08-07  
Resolution : 3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| $R_{free}$            | 180053                      | 1085 (3.54-3.46)                                      |
| Clashscore            | 190562                      | 1140 (3.54-3.46)                                      |
| Ramachandran outliers | 187476                      | 1113 (3.54-3.46)                                      |
| Sidechain outliers    | 187428                      | 1114 (3.54-3.46)                                      |
| RSRZ outliers         | 180081                      | 1084 (3.54-3.46)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 224    | <br>54% 17% 29%  |
| 1   | B     | 224    | <br>53% 17% 29%  |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 3   | O    | B     | 303 | -         | -        | X       | -                |

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron-sulfur cluster repair protein ScdA.

| Mol | Chain | Residues | Atoms |     |     |     |   | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
|     |       |          | Total | C   | N   | O   | S |         |         |       |
| 1   | A     | 158      | 1289  | 830 | 211 | 245 | 3 | 0       | 0       | 0     |
| 1   | B     | 158      | 1291  | 830 | 211 | 247 | 3 | 0       | 0       | 0     |

There are 6 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| A     | 30      | ALA      | CYS    | engineered mutation | UNP Q2FK11 |
| A     | 31      | ALA      | CYS    | engineered mutation | UNP Q2FK11 |
| A     | 191     | ALA      | CYS    | engineered mutation | UNP Q2FK11 |
| B     | 30      | ALA      | CYS    | engineered mutation | UNP Q2FK11 |
| B     | 31      | ALA      | CYS    | engineered mutation | UNP Q2FK11 |
| B     | 191     | ALA      | CYS    | engineered mutation | UNP Q2FK11 |

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms |    | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 2   | A     | 2        | Total | Fe | 0       | 0       |
|     |       |          | 2     | 2  |         |         |
| 2   | B     | 2        | Total | Fe | 0       | 0       |
|     |       |          | 2     | 2  |         |         |

- Molecule 3 is OXYGEN ATOM (CCD ID: O) (formula: O) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms |   | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---------|---------|
| 3   | A     | 1        | Total | O | 0       | 0       |
|     |       |          | 1     | 1 |         |         |
| 3   | B     | 1        | Total | O | 0       | 0       |
|     |       |          | 1     | 1 |         |         |

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms      |         | ZeroOcc | AltConf |
|-----|-------|----------|------------|---------|---------|---------|
| 4   | B     | 1        | Total<br>1 | Zn<br>1 | 0       | 0       |

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Iron-sulfur cluster repair protein ScdA

Chain A:  54% 17% 29%

MET ILE ASN LYS ASP ILE VAL ALA ASP VAL VAL THR ASP TYR PRO LYS ALA ALA ASP PHE ARG SER VAL GLY ILE ASP PHE ALA ALA GLY GLN VAL SER ILE GLU ALA LEU LEU LYS LYS ASN VAL ASP ASN GLU LEU LEU GLN ARG LEU ASN VAL ASP VAL GLU

GLN THR ASN THR PRO GLY S67 L68 R69 P70 V75 L78 I82 Y86 H87 L90 R91 L104 S105 K106 V107 H108 G109 M129 L144 D154 S155 I156 I160 D161 D162 L163 V164 S165 G171 M177 T181 S182 S183 Y184 I188 W194 Y198 Q199

R200 L201 A203 L207 T208 V212 H213 L214 L219 S224

- Molecule 1: Iron-sulfur cluster repair protein ScdA

Chain B:  53% 17% 29%

MET ILE ASN LYS ASP ILE VAL ALA ASP VAL VAL THR ASP TYR PRO LYS ALA ALA ASP PHE ARG SER VAL GLY ILE ASP PHE ALA ALA GLY GLN VAL SER ILE GLU ALA ALA ALA LEU LEU LYS LYS ASN VAL ASP ASN GLU LEU LEU GLN ARG LEU ASN VAL ASP VAL GLU

GLN THR ASN THR PRO GLY S67 L68 M69 P70 K71 M74 V75 S76 I79 A85 Y86 H87 E88 P89 L90 R91 T102 G109 P110 K119 D123 T124 F125 M129 L130 M133 E136 F141 D162 L163 V164 S165 D166 H167 I168 A169 T170 L174 M177

S182 E185 V197 L201 H211 V212 H213 L214 V218 S224

## 4 Data and refinement statistics

| Property  | Value   | Source           |
|---|---|------------------|
| Space group   | P 21 21 21  | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 69.02Å 92.53Å 110.92Å<br>90.00° 90.00° 90.00°               | Depositor        |
| Resolution (Å)  | 49.51 – 3.50<br>49.51 – 3.50                                | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 97.7 (49.51-3.50)<br>97.7 (49.51-3.50)                      | Depositor<br>EDS |
| $R_{merge}$   | (Not available)   | Depositor        |
| $R_{sym}$   | (Not available)   | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 1.62 (at 3.48Å)   | Xtrriage         |
| Refinement program  | PHENIX 1.21.1_5286  | Depositor        |
| R, $R_{free}$   | 0.185 , 0.238<br>0.185 , 0.238                              | Depositor<br>DCC |
| $R_{free}$ test set   | 936 reflections (9.92%)                                     | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 50.9  | Xtrriage         |
| Anisotropy  | 0.475   | Xtrriage         |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.35 , 44.9   | EDS              |
| L-test for twinning <sup>2</sup>  | $\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$ | Xtrriage         |
| Estimated twinning fraction   | No twinning to report.                                      | Xtrriage         |
| $F_o, F_c$ correlation  | 0.92  | EDS              |
| Total number of atoms   | 2587  | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 45.0  | wwPDB-VP         |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: O, FE, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |         | Bond angles |         |
|-----|-------|--------------|---------|-------------|---------|
|     |       | RMSZ         | # Z  >5 | RMSZ        | # Z  >5 |
| 1   | A     | 0.39         | 0/1320  | 0.57        | 0/1793  |
| 1   | B     | 0.40         | 0/1322  | 0.55        | 0/1795  |
| All | All   | 0.39         | 0/2642  | 0.56        | 0/3588  |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 1289  | 0        | 1273     | 25      | 0            |
| 1   | B     | 1291  | 0        | 1279     | 25      | 0            |
| 2   | A     | 2     | 0        | 0        | 0       | 0            |
| 2   | B     | 2     | 0        | 0        | 0       | 0            |
| 3   | A     | 1     | 0        | 0        | 1       | 0            |
| 3   | B     | 1     | 0        | 0        | 2       | 0            |
| 4   | B     | 1     | 0        | 0        | 0       | 0            |
| All | All   | 2587  | 0        | 2552     | 47      | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1           | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:B:214:LEU:HD12 | 1:B:218:VAL:HG21 | 1.73                     | 0.70              |
| 1:A:181:THR:O    | 1:A:184:TYR:N    | 2.26                     | 0.69              |
| 1:B:211:HIS:CE1  | 3:B:303:O:O      | 2.46                     | 0.68              |
| 1:A:160:ILE:O    | 1:A:164:VAL:HG23 | 1.99                     | 0.62              |
| 1:B:102:THR:HG23 | 1:B:119:LYS:HE2  | 1.84                     | 0.58              |
| 1:A:78:LEU:O     | 1:A:82:ILE:HG12  | 2.04                     | 0.58              |
| 1:A:68:LEU:HB2   | 1:B:68:LEU:HB2   | 1.86                     | 0.57              |
| 1:A:68:LEU:HD11  | 1:B:70:PRO:HB3   | 1.89                     | 0.55              |
| 1:A:144:LEU:HD23 | 1:A:156:ILE:HD11 | 1.89                     | 0.55              |
| 1:A:87:HIS:O     | 1:A:91:ARG:HG3   | 2.08                     | 0.53              |
| 1:B:125:PHE:CD1  | 1:B:170:THR:HG23 | 2.44                     | 0.53              |
| 1:A:181:THR:HB   | 1:A:198:TYR:CE2  | 2.44                     | 0.52              |
| 1:B:85:ALA:O     | 1:B:89:PRO:HG2   | 2.10                     | 0.51              |
| 1:A:129:MET:HE1  | 3:A:303:O:O      | 2.11                     | 0.51              |
| 1:A:87:HIS:HB3   | 1:A:91:ARG:NH1   | 2.27                     | 0.49              |
| 1:A:177:MET:HE1  | 1:A:201:LEU:HD22 | 1.93                     | 0.49              |
| 1:A:200:ARG:O    | 1:A:203:ALA:HB3  | 2.12                     | 0.48              |
| 1:B:74:ASN:OD1   | 1:B:76:SER:N     | 2.47                     | 0.48              |
| 1:A:208:THR:O    | 1:A:212:VAL:HG22 | 2.14                     | 0.47              |
| 1:B:79:ILE:HD13  | 1:B:141:PHE:HB3  | 1.96                     | 0.47              |
| 1:B:182:SER:O    | 1:B:185:GLU:HG2  | 2.14                     | 0.47              |
| 1:B:197:VAL:O    | 1:B:201:LEU:HG   | 2.15                     | 0.47              |
| 1:A:154:ASP:OD1  | 1:A:154:ASP:N    | 2.43                     | 0.47              |
| 1:B:87:HIS:O     | 1:B:91:ARG:HG3   | 2.15                     | 0.47              |
| 1:A:86:TYR:O     | 1:A:90:LEU:HG    | 2.14                     | 0.46              |
| 1:A:70:PRO:HB3   | 1:B:68:LEU:HD21  | 1.97                     | 0.46              |
| 1:B:119:LYS:NZ   | 1:B:123:ASP:OD2  | 2.40                     | 0.46              |
| 1:B:125:PHE:HD1  | 1:B:170:THR:HG23 | 1.79                     | 0.46              |
| 1:A:194:TRP:HD1  | 1:A:194:TRP:O    | 1.98                     | 0.45              |
| 1:B:162:ASP:O    | 1:B:165:SER:OG   | 2.32                     | 0.45              |
| 1:B:129:MET:HE1  | 3:B:303:O:O      | 2.16                     | 0.44              |
| 1:A:214:LEU:HD12 | 1:A:214:LEU:HA   | 1.65                     | 0.44              |
| 1:B:167:HIS:ND1  | 1:B:212:VAL:HG13 | 2.32                     | 0.44              |
| 1:A:105:SER:O    | 1:A:109:GLY:HA3  | 2.18                     | 0.44              |
| 1:A:75:VAL:HG13  | 1:A:219:LEU:HD11 | 1.99                     | 0.43              |
| 1:B:164:VAL:O    | 1:B:168:ILE:HG13 | 2.18                     | 0.43              |
| 1:B:174:LEU:HA   | 1:B:177:MET:HE3  | 2.01                     | 0.43              |
| 1:A:162:ASP:OD1  | 1:A:165:SER:OG   | 2.36                     | 0.42              |
| 1:A:104:LEU:HD22 | 1:A:194:TRP:HA   | 2.01                     | 0.42              |
| 1:B:109:GLY:N    | 1:B:110:PRO:HD2  | 2.34                     | 0.42              |

*Continued on next page...*

Continued from previous page...

| Atom-1           | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:A:188:ILE:H    | 1:A:188:ILE:HG12 | 1.63                     | 0.42              |
| 1:B:86:TYR:C     | 1:B:89:PRO:HD2   | 2.46                     | 0.41              |
| 1:B:136:GLU:HB3  | 1:B:141:PHE:CE2  | 2.55                     | 0.41              |
| 1:B:130:LEU:HD23 | 1:B:133:MET:HE3  | 2.02                     | 0.41              |
| 1:A:82:ILE:HD13  | 1:A:214:LEU:HD23 | 2.03                     | 0.40              |
| 1:A:90:LEU:HD23  | 1:A:207:LEU:HD23 | 2.03                     | 0.40              |
| 1:B:69:ASN:ND2   | 1:B:71:LYS:HB3   | 2.36                     | 0.40              |

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Favoured  | Allowed | Outliers | Percentiles |     |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1   | A     | 156/224 (70%) | 139 (89%) | 14 (9%) | 3 (2%)   | 6           | 33  |
| 1   | B     | 156/224 (70%) | 150 (96%) | 6 (4%)  | 0        | 100         | 100 |
| All | All   | 312/448 (70%) | 289 (93%) | 20 (6%) | 3 (1%)   | 12          | 44  |

All (3) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 182 | SER  |
| 1   | A     | 171 | GLY  |
| 1   | A     | 107 | VAL  |

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

| Mol | Chain | Analysed      | Rotameric  | Outliers | Percentiles |     |
|-----|-------|---------------|------------|----------|-------------|-----|
| 1   | A     | 146/202 (72%) | 146 (100%) | 0        | 100         | 100 |
| 1   | B     | 148/202 (73%) | 148 (100%) | 0        | 100         | 100 |
| All | All   | 294/404 (73%) | 294 (100%) | 0        | 100         | 100 |

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 80  | GLN  |
| 1   | A     | 217 | HIS  |
| 1   | B     | 96  | ASN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed      | <RSRZ> | #RSRZ>2      | OWAB(Å <sup>2</sup> ) | Q<0.9 |
|-----|-------|---------------|--------|--------------|-----------------------|-------|
| 1   | A     | 158/224 (70%) | -0.11  | 1 (0%) 85 65 | 26, 45, 69, 87        | 0     |
| 1   | B     | 158/224 (70%) | -0.34  | 0 100 100    | 25, 40, 64, 87        | 0     |
| All | All   | 316/448 (70%) | -0.22  | 1 (0%) 90 76 | 25, 42, 68, 87        | 0     |

All (1) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1   | A     | 155 | ASP  | 2.2  |

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


| Mol | Type | Chain | Res | Atoms | RSCC | RSR  | B-factors(Å <sup>2</sup> ) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 3   | O    | A     | 303 | 1/1   | 0.95 | 0.12 | 37,37,37,37                | 0     |
| 3   | O    | B     | 303 | 1/1   | 0.96 | 0.18 | 32,32,32,32                | 0     |
| 2   | FE   | A     | 301 | 1/1   | 0.98 | 0.13 | 44,44,44,44                | 1     |
| 2   | FE   | B     | 302 | 1/1   | 0.99 | 0.15 | 41,41,41,41                | 1     |


*Continued on next page...*

*Continued from previous page...*

| Mol | Type | Chain | Res | Atoms | RSCC | RSR  | B-factors( $\text{\AA}^2$ ) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|-----------------------------|-------|
| 2   | FE   | A     | 302 | 1/1   | 0.99 | 0.15 | 43,43,43,43                 | 0     |
| 2   | FE   | B     | 301 | 1/1   | 0.99 | 0.12 | 42,42,42,42                 | 0     |
| 4   | ZN   | B     | 304 | 1/1   | 0.99 | 0.06 | 63,63,63,63                 | 0     |




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around FE A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE B 302:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.