



wwPDB EM Validation Summary Report i

Dec 18, 2025 – 04:31 PM JST

PDB ID : 9WMV / pdb_00009wmv
EMDB ID : EMD-66106
Title : Co-transcriptional histone H3K36 methylation complex containing RNA polymerase II elongation complex, Set2, and the upstream nucleosome. (temp130, type B)
Authors : Kujirai, T.; Ehara, H.; Ito, T.; Henmi, M.; Sekine, S.; Kurumizaka, H.
Deposited on : 2025-09-03
Resolution : 4.66 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev129
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 4.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
RNA backbone	6643	2191	-
Q-score	-	25397	2028 (4.16 - 5.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.




Continued on next page...

Continued from previous page...



Continued on next page...

Continued from previous page...



2 Entry composition [\(i\)](#)

There are 33 unique types of molecules in this entry. The entry contains 80400 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1404	Total	C	N	O	S	0	0

11064 6975 1930 2089 70

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1164	Total	C	N	O	S	0	0

9284 5848 1639 1739 58

- Molecule 3 is a protein called RNA polymerase II third largest subunit B44, part of central core.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	263	Total	C	N	O	S	0	0

2098 1319 354 413 12

- Molecule 4 is a protein called RNA polymerase II subunit B32.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	174	Total	C	N	O	S	0	0

1349 828 244 274 3

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	213	Total	C	N	O	S	0	0

1741 1094 312 325 10

- Molecule 6 is a protein called RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	F	84	Total	C	N	O	S	0
			677	429	114	131	3	0

- Molecule 7 is a protein called RNA polymerase II subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	G	171	Total	C	N	O	S	0
			1325	858	214	248	5	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	H	133	Total	C	N	O	S	0
			1053	671	169	209	4	0

- Molecule 9 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
9	I	111	Total	C	N	O	S	0
			917	565	161	180	11	0

- Molecule 10 is a protein called RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms				AltConf	Trace
10	J	67	Total	C	N	O	S	0
			554	355	97	96	6	0

- Molecule 11 is a protein called RNA polymerase II subunit B12.5.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	K	113	Total	C	N	O	S	0
			932	599	160	169	4	0

- Molecule 12 is a protein called RNA polymerase subunit ABC10-alpha.

Mol	Chain	Residues	Atoms				AltConf	Trace
12	L	45	Total	C	N	O	S	0
			359	221	72	61	5	0

- Molecule 13 is a protein called Transcription elongation factor 1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	64	Total	C	N	O	S	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-2	GLY	-	expression tag	UNP C4QZ45
M	-1	PRO	-	expression tag	UNP C4QZ45
M	0	GLY	-	expression tag	UNP C4QZ45

- Molecule 14 is a DNA chain called DNA (198-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	168	Total	C	N	O	P	0	0

- Molecule 15 is a RNA chain called RNA (5'-R(P*GP*UP*CP*CP*UP*UP*UP*UP*GP*UP*GP*CP*GP*GP*CP*CP*UP*CP*G)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
15	P	19	Total	C	N	O	P	0	0

- Molecule 16 is a DNA chain called DNA (198-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
16	T	179	Total	C	N	O	P	0	0

- Molecule 17 is a protein called Transcription elongation factor SPT4.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	V	106	Total	C	N	O	S	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
V	7	MET	-	initiating methionine	UNP C4R0E6

- Molecule 18 is a protein called Transcription elongation factor SPT5.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	W	535	Total	C	N	O	S	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
W	-2	GLY	-	expression tag	UNP C4R370
W	-1	PRO	-	expression tag	UNP C4R370
W	0	GLY	-	expression tag	UNP C4R370

- Molecule 19 is a protein called Histone H3.3.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	a	104	Total	C	N	O	S	0	0
			841	530	163	145	3		
19	e	91	Total	C	N	O	S	0	0
			739	466	141	130	2		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	-3	GLY	-	expression tag	UNP P84243
a	-2	SER	-	expression tag	UNP P84243
a	-1	HIS	-	expression tag	UNP P84243
a	36	MET	LYS	variant	UNP P84243
e	-3	GLY	-	expression tag	UNP P84243
e	-2	SER	-	expression tag	UNP P84243
e	-1	HIS	-	expression tag	UNP P84243
e	36	MET	LYS	variant	UNP P84243

- Molecule 20 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	b	81	Total	C	N	O	S	0	0
			646	407	126	112	1		
20	f	81	Total	C	N	O	S	0	0
			646	407	126	112	1		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
b	-3	GLY	-	expression tag	UNP P62805

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
b	-2	SER	-	expression tag	UNP P62805
b	-1	HIS	-	expression tag	UNP P62805
f	-3	GLY	-	expression tag	UNP P62805
f	-2	SER	-	expression tag	UNP P62805
f	-1	HIS	-	expression tag	UNP P62805

- Molecule 21 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
21	c	104	Total	C	N	O	0	0
			805	508	157	140		
21	g	106	Total	C	N	O	0	0
			819	517	160	142		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
c	-3	GLY	-	expression tag	UNP P04908
c	-2	SER	-	expression tag	UNP P04908
c	-1	HIS	-	expression tag	UNP P04908
g	-3	GLY	-	expression tag	UNP P04908
g	-2	SER	-	expression tag	UNP P04908
g	-1	HIS	-	expression tag	UNP P04908

- Molecule 22 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms				AltConf	Trace
22	d	93	Total	C	N	O	S	0
			725	456	130	137	2	
22	h	94	Total	C	N	O	S	0
			735	462	132	139	2	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
d	-3	GLY	-	expression tag	UNP P06899
d	-2	SER	-	expression tag	UNP P06899
d	-1	HIS	-	expression tag	UNP P06899
h	-3	GLY	-	expression tag	UNP P06899
h	-2	SER	-	expression tag	UNP P06899
h	-1	HIS	-	expression tag	UNP P06899

- Molecule 23 is a protein called Transcription elongation factor Spt6.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	m	1178	Total	C	N	O	S	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
m	-2	GLY	-	expression tag	UNP C4R7H2
m	-1	PRO	-	expression tag	UNP C4R7H2
m	0	GLY	-	expression tag	UNP C4R7H2

- Molecule 24 is a protein called Protein that interacts with Spt6p and copurifies with Spt5p and RNA polymerase II.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	n	139	Total	C	N	O	S	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
n	-2	GLY	-	expression tag	UNP C4R7L8
n	-1	PRO	-	expression tag	UNP C4R7L8
n	0	GLY	-	expression tag	UNP C4R7L8

- Molecule 25 is a protein called Component of the Paf1p complex.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	q	930	Total	C	N	O	S	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
q	-39	MET	-	initiating methionine	UNP C4R6B2
q	-38	LYS	-	expression tag	UNP C4R6B2
q	-37	ASP	-	expression tag	UNP C4R6B2
q	-36	HIS	-	expression tag	UNP C4R6B2
q	-35	LEU	-	expression tag	UNP C4R6B2
q	-34	ILE	-	expression tag	UNP C4R6B2
q	-33	HIS	-	expression tag	UNP C4R6B2
q	-32	ASN	-	expression tag	UNP C4R6B2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
q	-31	HIS	-	expression tag	UNP C4R6B2
q	-30	HIS	-	expression tag	UNP C4R6B2
q	-29	LYS	-	expression tag	UNP C4R6B2
q	-28	HIS	-	expression tag	UNP C4R6B2
q	-27	GLU	-	expression tag	UNP C4R6B2
q	-26	HIS	-	expression tag	UNP C4R6B2
q	-25	ALA	-	expression tag	UNP C4R6B2
q	-24	HIS	-	expression tag	UNP C4R6B2
q	-23	ALA	-	expression tag	UNP C4R6B2
q	-22	GLU	-	expression tag	UNP C4R6B2
q	-21	HIS	-	expression tag	UNP C4R6B2
q	-20	ASP	-	expression tag	UNP C4R6B2
q	-19	TYR	-	expression tag	UNP C4R6B2
q	-18	LYS	-	expression tag	UNP C4R6B2
q	-17	ASP	-	expression tag	UNP C4R6B2
q	-16	ASP	-	expression tag	UNP C4R6B2
q	-15	ASP	-	expression tag	UNP C4R6B2
q	-14	ASP	-	expression tag	UNP C4R6B2
q	-13	LYS	-	expression tag	UNP C4R6B2
q	-12	GLU	-	expression tag	UNP C4R6B2
q	-11	HIS	-	expression tag	UNP C4R6B2
q	-10	LEU	-	expression tag	UNP C4R6B2
q	-9	TYR	-	expression tag	UNP C4R6B2
q	-8	PHE	-	expression tag	UNP C4R6B2
q	-7	GLN	-	expression tag	UNP C4R6B2
q	-6	GLY	-	expression tag	UNP C4R6B2
q	-5	SER	-	expression tag	UNP C4R6B2
q	-4	SER	-	expression tag	UNP C4R6B2
q	-3	GLY	-	expression tag	UNP C4R6B2
q	-2	SER	-	expression tag	UNP C4R6B2
q	-1	SER	-	expression tag	UNP C4R6B2
q	0	GLY	-	expression tag	UNP C4R6B2

- Molecule 26 is a protein called RNAPII-associated chromatin remodeling Paf1 complex sub-unit.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	r	266	Total	C	N	O	S	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
r	-29	MET	-	initiating methionine	UNP F2QQ42
r	-28	LYS	-	expression tag	UNP F2QQ42
r	-27	ASP	-	expression tag	UNP F2QQ42
r	-26	HIS	-	expression tag	UNP F2QQ42
r	-25	LEU	-	expression tag	UNP F2QQ42
r	-24	ILE	-	expression tag	UNP F2QQ42
r	-23	HIS	-	expression tag	UNP F2QQ42
r	-22	ASN	-	expression tag	UNP F2QQ42
r	-21	HIS	-	expression tag	UNP F2QQ42
r	-20	HIS	-	expression tag	UNP F2QQ42
r	-19	LYS	-	expression tag	UNP F2QQ42
r	-18	HIS	-	expression tag	UNP F2QQ42
r	-17	GLU	-	expression tag	UNP F2QQ42
r	-16	HIS	-	expression tag	UNP F2QQ42
r	-15	ALA	-	expression tag	UNP F2QQ42
r	-14	HIS	-	expression tag	UNP F2QQ42
r	-13	ALA	-	expression tag	UNP F2QQ42
r	-12	GLU	-	expression tag	UNP F2QQ42
r	-11	HIS	-	expression tag	UNP F2QQ42
r	-10	LEU	-	expression tag	UNP F2QQ42
r	-9	TYR	-	expression tag	UNP F2QQ42
r	-8	PHE	-	expression tag	UNP F2QQ42
r	-7	GLN	-	expression tag	UNP F2QQ42
r	-6	GLY	-	expression tag	UNP F2QQ42
r	-5	SER	-	expression tag	UNP F2QQ42
r	-4	SER	-	expression tag	UNP F2QQ42
r	-3	GLY	-	expression tag	UNP F2QQ42
r	-2	SER	-	expression tag	UNP F2QQ42
r	-1	SER	-	expression tag	UNP F2QQ42
r	0	GLY	-	expression tag	UNP F2QQ42

- Molecule 27 is a protein called Histone-lysine N-methyltransferase, H3 lysine-36 specific.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	s	431	3419	2126	593	677	23	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
s	-2	GLY	-	expression tag	UNP C4QY01
s	-1	PRO	-	expression tag	UNP C4QY01
s	0	GLY	-	expression tag	UNP C4QY01

- Molecule 28 is a protein called Leo1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
28	u	188	1522	955	262	302	3	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
u	-29	MET	-	initiating methionine	UNP C4R3K1
u	-28	LYS	-	expression tag	UNP C4R3K1
u	-27	ASP	-	expression tag	UNP C4R3K1
u	-26	HIS	-	expression tag	UNP C4R3K1
u	-25	LEU	-	expression tag	UNP C4R3K1
u	-24	ILE	-	expression tag	UNP C4R3K1
u	-23	HIS	-	expression tag	UNP C4R3K1
u	-22	ASN	-	expression tag	UNP C4R3K1
u	-21	HIS	-	expression tag	UNP C4R3K1
u	-20	HIS	-	expression tag	UNP C4R3K1
u	-19	LYS	-	expression tag	UNP C4R3K1
u	-18	HIS	-	expression tag	UNP C4R3K1
u	-17	GLU	-	expression tag	UNP C4R3K1
u	-16	HIS	-	expression tag	UNP C4R3K1
u	-15	ALA	-	expression tag	UNP C4R3K1
u	-14	HIS	-	expression tag	UNP C4R3K1
u	-13	ALA	-	expression tag	UNP C4R3K1
u	-12	GLU	-	expression tag	UNP C4R3K1
u	-11	HIS	-	expression tag	UNP C4R3K1
u	-10	LEU	-	expression tag	UNP C4R3K1
u	-9	TYR	-	expression tag	UNP C4R3K1
u	-8	PHE	-	expression tag	UNP C4R3K1
u	-7	GLN	-	expression tag	UNP C4R3K1
u	-6	GLY	-	expression tag	UNP C4R3K1
u	-5	SER	-	expression tag	UNP C4R3K1
u	-4	SER	-	expression tag	UNP C4R3K1
u	-3	GLY	-	expression tag	UNP C4R3K1
u	-2	SER	-	expression tag	UNP C4R3K1
u	-1	SER	-	expression tag	UNP C4R3K1
u	0	GLY	-	expression tag	UNP C4R3K1

- Molecule 29 is a protein called RNAP II-associated protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	v	349	Total	C	N	O	S	0	0
			2878	1835	510	528	5		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
v	-2	GLY	-	expression tag	UNP C4R997
v	-1	SER	-	expression tag	UNP C4R997
v	0	ALA	-	expression tag	UNP C4R997

- Molecule 30 is a protein called Constituent of Paf1 complex with RNA polymerase II, Paf1p, Hpr1p, Ctr9, Leo1, Rtf1 and Ccr4p.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	x	205	Total	C	N	O	S	0	0
			1682	1086	287	307	2		

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
x	-29	MET	-	initiating methionine	UNP C4R1E6
x	-28	LYS	-	expression tag	UNP C4R1E6
x	-27	ASP	-	expression tag	UNP C4R1E6
x	-26	HIS	-	expression tag	UNP C4R1E6
x	-25	LEU	-	expression tag	UNP C4R1E6
x	-24	ILE	-	expression tag	UNP C4R1E6
x	-23	HIS	-	expression tag	UNP C4R1E6
x	-22	ASN	-	expression tag	UNP C4R1E6
x	-21	HIS	-	expression tag	UNP C4R1E6
x	-20	HIS	-	expression tag	UNP C4R1E6
x	-19	LYS	-	expression tag	UNP C4R1E6
x	-18	HIS	-	expression tag	UNP C4R1E6
x	-17	GLU	-	expression tag	UNP C4R1E6
x	-16	HIS	-	expression tag	UNP C4R1E6
x	-15	ALA	-	expression tag	UNP C4R1E6
x	-14	HIS	-	expression tag	UNP C4R1E6
x	-13	ALA	-	expression tag	UNP C4R1E6
x	-12	GLU	-	expression tag	UNP C4R1E6
x	-11	HIS	-	expression tag	UNP C4R1E6
x	-10	LEU	-	expression tag	UNP C4R1E6
x	-9	TYR	-	expression tag	UNP C4R1E6
x	-8	PHE	-	expression tag	UNP C4R1E6
x	-7	GLN	-	expression tag	UNP C4R1E6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
x	-6	GLY	-	expression tag	UNP C4R1E6
x	-5	SER	-	expression tag	UNP C4R1E6
x	-4	SER	-	expression tag	UNP C4R1E6
x	-3	GLY	-	expression tag	UNP C4R1E6
x	-2	SER	-	expression tag	UNP C4R1E6
x	-1	SER	-	expression tag	UNP C4R1E6
x	0	GLY	-	expression tag	UNP C4R1E6


- Molecule 31 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
31	A	2	Total Zn 2 2	0
31	B	1	Total Zn 1 1	0
31	C	1	Total Zn 1 1	0
31	I	2	Total Zn 2 2	0
31	J	1	Total Zn 1 1	0
31	L	1	Total Zn 1 1	0
31	M	1	Total Zn 1 1	0
31	V	1	Total Zn 1 1	0
31	s	3	Total Zn 3 3	0

- Molecule 32 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
32	A	1	Total Mg 1 1	0

- Molecule 33 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	S	
33	S	1	26	14	6	5	1	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase subunit




- Molecule 2: DNA-directed RNA polymerase subunit beta






- Molecule 3: RNA polymerase II third largest subunit B44, part of central core




- Molecule 4: RNA polymerase II subunit B32



- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




- Molecule 6: RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III







- Molecule 7: RNA polymerase II subunit



- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 9: DNA-directed RNA polymerase subunit




- Molecule 10: RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III




- Molecule 11: RNA polymerase II subunit B12.5




- Molecule 12: RNA polymerase subunit ABC10-alpha



- Molecule 13: Transcription elongation factor 1 homolog




- Molecule 14: DNA (198-MER)




- Molecule 15: RNA (5'-R(P*GP*UP*CP*CP*UP*UP*UP*UP*GP*UP*GP*GP*CP*CP*UP*CP*G)-3')



- Molecule 16: DNA (198-MER)








- Molecule 17: Transcription elongation factor SPT4





- Molecule 18: Transcription elongation factor SPT5





- Molecule 19: Histone H3.3







- Molecule 19: Histone H3.3





- Molecule 20: Histone H4



- Molecule 20: Histone H4





- Molecule 21: Histone H2A type 1-B/E






- Molecule 21: Histone H2A type 1-B/E





- Molecule 22: Histone H2B type 1-J




- Molecule 22: Histone H2B type 1-J





- Molecule 23: Transcription elongation factor Spt6









- Molecule 24: Protein that interacts with Spt6p and copurifies with Spt5p and RNA polymerase II






• Molecule 26: RNAPII-associated chromatin remodeling Paf1 complex subunit




• Molecule 27: Histone-lysine N-methyltransferase, H3 lysine-36 specific







- Molecule 28: Leo1

Chain u: 23% 36% 5% 59%







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	10768	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	59.1	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.070	Depositor
Minimum map value	-0.026	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.015	Depositor
Map size (Å)	531.0, 531.0, 531.0	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.475, 1.475, 1.475	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/11267	0.50	0/15222
2	B	0.23	0/9464	0.51	0/12763
3	C	0.22	0/2139	0.46	0/2895
4	D	0.17	0/1361	0.44	0/1837
5	E	0.22	0/1773	0.54	0/2385
6	F	0.21	0/687	0.48	0/931
7	G	0.17	0/1354	0.39	0/1837
8	H	0.22	0/1070	0.43	0/1444
9	I	0.15	0/934	0.41	0/1257
10	J	0.22	0/563	0.52	0/753
11	K	0.22	0/953	0.49	0/1291
12	L	0.23	0/365	0.58	0/484
13	M	0.17	0/513	0.37	0/693
14	N	0.28	0/3837	0.60	0/5930
15	P	0.26	0/441	0.63	0/684
16	T	0.25	0/4142	0.53	0/6378
17	V	0.17	0/840	0.47	0/1140
18	W	0.20	0/4319	0.51	0/5838
19	a	0.24	0/853	0.63	0/1144
19	e	0.26	0/747	0.65	0/1001
20	b	0.26	0/653	0.67	0/873
20	f	0.27	0/653	0.67	0/873
21	c	0.23	0/815	0.60	0/1100
21	g	0.26	0/829	0.62	0/1118
22	d	0.24	0/736	0.61	0/990
22	h	0.27	0/746	0.60	0/1001
23	m	0.20	0/9847	0.46	0/13321
24	n	0.19	0/1132	0.45	0/1526
25	q	0.18	0/7689	0.40	0/10368
26	r	0.19	0/2169	0.43	0/2901
27	s	0.33	0/3473	0.62	0/4676
28	u	0.17	0/1554	0.38	0/2102

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
29	v	0.18	0/2944	0.44	0/3973
30	x	0.16	0/1716	0.39	0/2310
All	All	0.22	0/82578	0.50	0/113039

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11064	0	11090	62	0
2	B	9284	0	9282	63	0
3	C	2098	0	2057	10	0
4	D	1349	0	1345	5	0
5	E	1741	0	1754	10	0
6	F	677	0	693	3	0
7	G	1325	0	1342	9	0
8	H	1053	0	1050	4	0
9	I	917	0	864	10	0
10	J	554	0	573	1	0
11	K	932	0	944	11	0
12	L	359	0	358	1	0
13	M	505	0	495	3	0
14	N	3443	0	1927	22	0
15	P	398	0	203	6	0
16	T	3671	0	1976	20	0
17	V	824	0	795	11	0
18	W	4250	0	4296	52	0
19	a	841	0	883	7	0
19	e	739	0	778	8	0
20	b	646	0	687	5	0
20	f	646	0	687	5	0
21	c	805	0	861	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
21	g	819	0	879	9	0
22	d	725	0	745	8	0
22	h	735	0	758	9	0
23	m	9653	0	9500	63	0
24	n	1115	0	1186	15	0
25	q	7552	0	7545	43	0
26	r	2139	0	2155	19	0
27	s	3419	0	3351	20	0
28	u	1522	0	1492	19	0
29	v	2878	0	2873	25	0
30	x	1682	0	1731	7	0
31	A	2	0	0	0	0
31	B	1	0	0	0	0
31	C	1	0	0	0	0
31	I	2	0	0	0	0
31	J	1	0	0	0	0
31	L	1	0	0	0	0
31	M	1	0	0	0	0
31	V	1	0	0	0	0
31	s	3	0	0	0	0
32	A	1	0	0	0	0
33	s	26	0	19	1	0
All	All	80400	0	77174	456	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 456 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:445:PHE:CE2	1:A:471:LEU:HD21	2.24	0.72
14:N:27:DG:H2'	14:N:28:DT:H71	1.74	0.69
18:W:461:LEU:HD13	23:m:278:THR:HA	1.74	0.69
18:W:444:LEU:HG	23:m:284:PHE:CE1	2.28	0.69
18:W:218:ILE:CD1	24:n:289:MET:HE1	2.23	0.68

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1392/1743 (80%)	1355 (97%)	36 (3%)	1 (0%)	48 83
2	B	1154/1227 (94%)	1127 (98%)	27 (2%)	0	100 100
3	C	261/304 (86%)	259 (99%)	2 (1%)	0	100 100
4	D	170/186 (91%)	168 (99%)	2 (1%)	0	100 100
5	E	211/214 (99%)	207 (98%)	4 (2%)	0	100 100
6	F	82/155 (53%)	79 (96%)	3 (4%)	0	100 100
7	G	169/171 (99%)	166 (98%)	3 (2%)	0	100 100
8	H	129/145 (89%)	125 (97%)	4 (3%)	0	100 100
9	I	109/115 (95%)	105 (96%)	4 (4%)	0	100 100
10	J	65/72 (90%)	65 (100%)	0	0	100 100
11	K	111/118 (94%)	109 (98%)	2 (2%)	0	100 100
12	L	43/72 (60%)	41 (95%)	2 (5%)	0	100 100
13	M	62/113 (55%)	61 (98%)	1 (2%)	0	100 100
17	V	104/108 (96%)	102 (98%)	2 (2%)	0	100 100
18	W	529/911 (58%)	506 (96%)	23 (4%)	0	100 100
19	a	102/139 (73%)	100 (98%)	2 (2%)	0	100 100
19	e	89/139 (64%)	88 (99%)	1 (1%)	0	100 100
20	b	79/106 (74%)	76 (96%)	3 (4%)	0	100 100
20	f	79/106 (74%)	76 (96%)	3 (4%)	0	100 100
21	c	102/133 (77%)	100 (98%)	2 (2%)	0	100 100
21	g	104/133 (78%)	101 (97%)	3 (3%)	0	100 100
22	d	91/129 (70%)	88 (97%)	3 (3%)	0	100 100
22	h	92/129 (71%)	90 (98%)	2 (2%)	0	100 100
23	m	1170/1503 (78%)	1152 (98%)	18 (2%)	0	100 100
24	n	137/417 (33%)	136 (99%)	1 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
25	q	928/1084 (86%)	923 (100%)	5 (0%)	0	100 100
26	r	260/544 (48%)	254 (98%)	6 (2%)	0	100 100
27	s	429/725 (59%)	417 (97%)	12 (3%)	0	100 100
28	u	186/459 (40%)	186 (100%)	0	0	100 100
29	v	341/396 (86%)	330 (97%)	11 (3%)	0	100 100
30	x	201/395 (51%)	200 (100%)	1 (0%)	0	100 100
All	All	8981/12191 (74%)	8792 (98%)	188 (2%)	1 (0%)	100 100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	960	VAL

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1219/1528 (80%)	1219 (100%)	0	100 100
2	B	1018/1077 (94%)	1014 (100%)	4 (0%)	89 91
3	C	236/264 (89%)	236 (100%)	0	100 100
4	D	149/160 (93%)	148 (99%)	1 (1%)	81 87
5	E	196/197 (100%)	196 (100%)	0	100 100
6	F	75/137 (55%)	75 (100%)	0	100 100
7	G	148/148 (100%)	148 (100%)	0	100 100
8	H	120/130 (92%)	120 (100%)	0	100 100
9	I	106/109 (97%)	106 (100%)	0	100 100
10	J	61/66 (92%)	61 (100%)	0	100 100
11	K	104/109 (95%)	104 (100%)	0	100 100
12	L	38/56 (68%)	38 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
13	M	61/99 (62%)	61 (100%)	0	100	100
17	V	90/92 (98%)	90 (100%)	0	100	100
18	W	482/796 (61%)	482 (100%)	0	100	100
19	a	87/112 (78%)	87 (100%)	0	100	100
19	e	77/112 (69%)	77 (100%)	0	100	100
20	b	66/81 (82%)	66 (100%)	0	100	100
20	f	66/81 (82%)	66 (100%)	0	100	100
21	c	83/102 (81%)	83 (100%)	0	100	100
21	g	84/102 (82%)	84 (100%)	0	100	100
22	d	79/107 (74%)	79 (100%)	0	100	100
22	h	80/107 (75%)	79 (99%)	1 (1%)	65	77
23	m	1079/1354 (80%)	1078 (100%)	1 (0%)	92	94
24	n	125/361 (35%)	125 (100%)	0	100	100
25	q	824/962 (86%)	824 (100%)	0	100	100
26	r	239/485 (49%)	239 (100%)	0	100	100
27	s	380/649 (59%)	380 (100%)	0	100	100
28	u	173/406 (43%)	173 (100%)	0	100	100
29	v	325/369 (88%)	325 (100%)	0	100	100
30	x	190/354 (54%)	190 (100%)	0	100	100
All	All	8060/10712 (75%)	8053 (100%)	7 (0%)	92	94

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	904	ARG
4	D	11	ARG
23	m	770	PHE
22	h	42	TYR
2	B	879	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 100 such sidechains are listed below:

Mol	Chain	Res	Type
23	m	819	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
25	q	147	ASN
30	x	251	GLN
23	m	886	HIS
23	m	1338	ASN

5.3.3 RNA [\(i\)](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
15	P	18/19 (94%)	8 (44%)	1 (5%)

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
15	P	-6	U
15	P	-5	C
15	P	-4	C
15	P	-3	U
15	P	-2	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
15	P	-4	C

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 15 ligands modelled in this entry, 14 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
33	SAH	s	1904	-	24,28,28	0.83	1 (4%)	25,40,40	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	SAH	s	1904	-	-	2/11/31/31	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
33	s	1904	SAH	C8-N7	-2.44	1.30	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
33	s	1904	SAH	O-C-CA-N
33	s	1904	SAH	OXT-C-CA-N


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
33	s	1904	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-66106. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [\(i\)](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 181




Y Index: 192




Z Index: 178


6.3.2 Raw map



X Index: 187



Y Index: 186




Z Index: 178


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [\(i\)](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [\(i\)](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1042 nm³; this corresponds to an approximate mass of 941 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.215 \AA^{-1}

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.215 \AA^{-1}

8.2 Resolution estimates [\(i\)](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.66	-	-
Author-provided FSC curve	4.60	8.60	5.31
Unmasked-calculated*	8.77	20.37	9.51

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.77 differs from the reported value 4.66 by more than 10 %


9 Map-model fit [\(i\)](#)

This section contains information regarding the fit between EMDB map EMD-66106 and PDB model 9WMV. Per-residue inclusion information can be found in section [3](#) on page [17](#).


9.1 Map-model overlay [\(i\)](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).

9.4 Atom inclusion [\(i\)](#)



At the recommended contour level, 74% of all backbone atoms, 66% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6600	0.1680
A	0.8930	0.3470
B	0.8700	0.3400
C	0.8790	0.3480
D	0.7450	0.1530
E	0.9270	0.3100
F	0.8980	0.3800
G	0.8150	0.2200
H	0.9040	0.3550
I	0.8330	0.1930
J	0.8870	0.3580
K	0.9160	0.3730
L	0.9110	0.3080
M	0.7820	0.1120
N	0.4570	0.0640
P	0.8590	0.2480
T	0.4800	0.0760
V	0.8210	0.0690
W	0.6620	0.1200
a	0.5960	-0.0130
b	0.6050	0.0130
c	0.3180	-0.0050
d	0.3090	-0.0140
e	0.2680	0.0020
f	0.4180	0.0560
g	0.4030	-0.0170
h	0.3870	0.0050
m	0.5280	0.0720
n	0.5490	0.0820
q	0.6900	0.0710
r	0.3420	0.0740
s	0.3160	0.0210
u	0.3750	0.0760
v	0.4040	0.0450
x	0.6370	0.1580

