

wwPDB EM Validation Summary Report (i)

Oct 28, 2024 – 12:27 PM JST

PDB ID : 8WP8

EMDB ID : EMD-37711

Title: Cryo-EM structure of SARS-CoV-2 Omicron BA.2.86 RBD in complex with

human ACE2

Authors : Li, L.J.; Gu, Y.H.; Qi, J.X.; Gao, G.F.

Deposited on : 2023-10-09

Resolution : 2.89 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain						
1	A	606	90%		8% •				
2	В	228	63%	18%	19%				
3	С	3	100%						
3	Е	3	100%						
4	D	2	100%						



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6539 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Processed angiotensin-converting enzyme 2.

\mathbf{Mol}	Chain	Residues	Atoms					AltConf	Trace
1	A	596	Total 4869	C 3116	N 807	O 917	S 29	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	616	HIS	-	expression tag	UNP Q9BYF1
A	617	HIS	-	expression tag	UNP Q9BYF1
A	618	HIS	-	expression tag	UNP Q9BYF1
A	619	HIS	-	expression tag	UNP Q9BYF1
A	620	HIS	-	expression tag	UNP Q9BYF1
A	621	HIS	-	expression tag	UNP Q9BYF1
A	622	HIS	-	expression tag	UNP Q9BYF1
A	623	HIS	-	expression tag	UNP Q9BYF1

• Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
2	В	184	Total 1493	C 969	N 250	O 267	S 7	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	332	VAL	ILE	$\operatorname{conflict}$	UNP P0DTC2
В	339	HIS	GLY	variant	UNP P0DTC2
В	356	THR	LYS	conflict	UNP P0DTC2
В	371	PHE	SER	variant	UNP P0DTC2
В	373	PRO	SER	variant	UNP P0DTC2
В	375	PHE	SER	variant	UNP P0DTC2
В	376	ALA	THR	variant	UNP P0DTC2
В	403	LYS	ARG	conflict	UNP P0DTC2
В	405	ASN	ASP	variant	UNP P0DTC2

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	408	SER	ARG	variant	UNP P0DTC2
В	417	ASN	LYS	variant	UNP P0DTC2
В	440	LYS	ASN	variant	UNP P0DTC2
В	445	HIS	VAL	conflict	UNP P0DTC2
В	446	SER	GLY	variant	UNP P0DTC2
В	450	ASP	ASN	conflict	UNP P0DTC2
В	452	TRP	LEU	conflict	UNP P0DTC2
В	460	LYS	ASN	variant	UNP P0DTC2
В	477	ASN	SER	variant	UNP P0DTC2
В	478	LYS	THR	variant	UNP P0DTC2
В	481	LYS	ASN	conflict	UNP P0DTC2
В	?	-	VAL	deletion	UNP P0DTC2
В	484	LYS	GLU	variant	UNP P0DTC2
В	486	PRO	PHE	variant	UNP P0DTC2
В	498	ARG	GLN	variant	UNP P0DTC2
В	501	TYR	ASN	variant	UNP P0DTC2
В	505	HIS	TYR	variant	UNP P0DTC2
В	542	HIS	-	expression tag	UNP P0DTC2
В	543	HIS	-	expression tag	UNP P0DTC2
В	544	HIS	-	expression tag	UNP P0DTC2
В	545	HIS	-	expression tag	UNP P0DTC2
В	546	HIS	-	expression tag	UNP P0DTC2
В	547	HIS	-	expression tag	UNP P0DTC2

• Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			AltConf	Trace	
3	С	2	Total	С	N	О	0	0
9		5	39	22	2	15	0	
2	E	2	Total	С	N	О	0	0
3	Ľ	3	39	22	2	15	0	U

 \bullet Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.






Mol	Chain	Residues	Atoms				AltConf	Trace
4	D	2	Total 28	C 16	N 2	O 10	0	0

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
5	A	1	Total Zn 1 1	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	AltConf
6	Δ	1	Total C N O	0
0	Λ	1	14 8 1 5	U
6	Δ	1	Total C N O	0
0	Λ	1	14 8 1 5	0
6	Δ	1	Total C N O	0
0	Λ	1	14 8 1 5	U
6	R	1	Total C N O	0
	Ъ	1	14 8 1 5	

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms				AltConf
6	D	1	Total	С	N	О	0
U	Б	1	14	8	1	5	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Processed angiotensin-converting enzyme 2



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



Chain D: 100%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	341631	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	60	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.25	0/5010	0.43	0/6807	
2	В	0.27	0/1543	0.50	0/2097	
All	All	0.25	0/6553	0.45	0/8904	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4869	0	4640	34	0
2	В	1493	0	1413	25	0
3	С	39	0	34	0	0
3	Ε	39	0	34	0	0
4	D	28	0	25	0	0
5	A	1	0	0	0	0
6	A	42	0	39	0	0
6	В	28	0	26	0	0
All	All	6539	0	6211	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



The worst 5 of 58 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\mbox{\normalfont\AA}) \end{aligned}$
1:A:152:MET:O	1:A:161:ARG:NH2	2.31	0.63
1:A:134:ASN:OD1	1:A:137:ASN:ND2	2.34	0.61
1:A:351:LEU:HB2	1:A:355:ASP:HB3	1.85	0.59
1:A:35:GLU:OE2	2:B:493:GLN:NE2	2.36	0.59
1:A:177:ARG:HB2	1:A:498:CYS:HB2	1.85	0.59

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	595/606 (98%)	578 (97%)	17 (3%)	0	100	100
2	В	182/228 (80%)	163 (90%)	19 (10%)	0	100	100
All	All	777/834 (93%)	741 (95%)	36 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

\mathbf{Mol}	Chain	Analysed	Analysed Rotameric Ou		Percentiles	
1	A	527/536 (98%)	527 (100%)	0	100 100	

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	В	160/202 (79%)	160 (100%)	0	100	100	
All	All	687/738 (93%)	687 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	401	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Type Chain	nain Res	Link	Во	ond leng	ths	Bond angles		
MIOI	Type	Chain			Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	С	1	1,3	14,14,15	0.24	0	17,19,21	0.37	0
3	NAG	С	2	3	14,14,15	0.25	0	17,19,21	0.41	0
3	BMA	С	3	3	11,11,12	0.65	0	15,15,17	0.85	0
4	NAG	D	1	1,4	14,14,15	0.26	0	17,19,21	0.48	0
4	NAG	D	2	4	14,14,15	0.28	0	17,19,21	0.33	0
3	NAG	Е	1	1,3	14,14,15	0.34	0	17,19,21	0.48	0
3	NAG	Е	2	3	14,14,15	0.24	0	17,19,21	0.52	0
3	BMA	Е	3	3	11,11,12	0.64	0	15,15,17	0.84	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	С	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	С	2	3	-	0/6/23/26	0/1/1/1
3	BMA	С	3	3	-	1/2/19/22	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
3	NAG	Е	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	Ε	2	3	-	0/6/23/26	0/1/1/1
3	BMA	Е	3	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	Е	1	NAG	C8-C7-N2-C2
3	Е	1	NAG	O7-C7-N2-C2
4	D	2	NAG	O5-C5-C6-O6
3	С	3	BMA	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain		their Dec	Res Link	Bo	Bond lengths			Bond angles		
MIOI	wioi Type Chain	nes	Counts		RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
6	NAG	В	602	2	14,14,15	0.32	0	17,19,21	0.68	0	
6	NAG	A	703	1	14,14,15	0.23	0	17,19,21	0.34	0	
6	NAG	A	704	1	14,14,15	0.23	0	17,19,21	0.53	0	
6	NAG	A	702	1	14,14,15	0.22	0	17,19,21	0.55	0	



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	В	601	2	14,14,15	0.39	0	17,19,21	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	В	602	2	-	2/6/23/26	0/1/1/1
6	NAG	A	703	1	-	2/6/23/26	0/1/1/1
6	NAG	A	704	1	-	3/6/23/26	0/1/1/1
6	NAG	A	702	1	-	2/6/23/26	0/1/1/1
6	NAG	В	601	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	В	602	NAG	C8-C7-N2-C2
6	В	602	NAG	O7-C7-N2-C2
6	A	702	NAG	O5-C5-C6-O6
6	A	702	NAG	C4-C5-C6-O6
6	A	703	NAG	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

