



## wwPDB EM Validation Summary Report i

Dec 7, 2025 – 12:41 AM JST

PDB ID : 9WXM / pdb\_00009wxm  
EMDB ID : EMD-66350  
Title : Cryo-EM structure of the full-length GPR15L bound GPR15-Gi complex  
Authors : Han, S.; Wu, B.; Zhao, Q.  
Deposited on : 2025-09-25  
Resolution : 3.30 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:


EMDB validation analysis : **FAILED**  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
EM percentile statistics : **NOT EXECUTED**  
MapQ : **FAILED**  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
**ELECTRON MICROSCOPY**

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .



## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 8643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein GPR15LG.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
1	L	14	105	71	16	18	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	0	GLY	-	expression tag	UNP Q6UWK7

- Molecule 2 is a protein called G-protein coupled receptor 15.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	R	287	2233	1485	366	363	19	0	0

There are 49 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-2	GLY	-	expression tag	UNP P49685
R	-1	ALA	-	expression tag	UNP P49685
R	0	PRO	-	expression tag	UNP P49685
R	331	GLU	-	expression tag	UNP P49685
R	332	PHE	-	expression tag	UNP P49685
R	333	LEU	-	expression tag	UNP P49685
R	334	GLU	-	expression tag	UNP P49685
R	335	VAL	-	expression tag	UNP P49685
R	336	LEU	-	expression tag	UNP P49685
R	337	PHE	-	expression tag	UNP P49685
R	338	GLN	-	expression tag	UNP P49685
R	339	GLY	-	expression tag	UNP P49685
R	340	PRO	-	expression tag	UNP P49685
R	341	TRP	-	expression tag	UNP P49685
R	342	SER	-	expression tag	UNP P49685
R	343	HIS	-	expression tag	UNP P49685

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
R	344	PRO	-	expression tag	UNP P49685
R	345	GLN	-	expression tag	UNP P49685
R	346	PHE	-	expression tag	UNP P49685
R	347	GLU	-	expression tag	UNP P49685
R	348	LYS	-	expression tag	UNP P49685
R	349	GLY	-	expression tag	UNP P49685
R	350	GLY	-	expression tag	UNP P49685
R	351	GLY	-	expression tag	UNP P49685
R	352	SER	-	expression tag	UNP P49685
R	353	GLY	-	expression tag	UNP P49685
R	354	GLY	-	expression tag	UNP P49685
R	355	GLY	-	expression tag	UNP P49685
R	356	SER	-	expression tag	UNP P49685
R	357	GLY	-	expression tag	UNP P49685
R	358	GLY	-	expression tag	UNP P49685
R	359	SER	-	expression tag	UNP P49685
R	360	ALA	-	expression tag	UNP P49685
R	361	TRP	-	expression tag	UNP P49685
R	362	SER	-	expression tag	UNP P49685
R	363	HIS	-	expression tag	UNP P49685
R	364	PRO	-	expression tag	UNP P49685
R	365	GLN	-	expression tag	UNP P49685
R	366	PHE	-	expression tag	UNP P49685
R	367	GLU	-	expression tag	UNP P49685
R	368	LYS	-	expression tag	UNP P49685
R	369	ASP	-	expression tag	UNP P49685
R	370	TYR	-	expression tag	UNP P49685
R	371	LYS	-	expression tag	UNP P49685
R	372	ASP	-	expression tag	UNP P49685
R	373	ASP	-	expression tag	UNP P49685
R	374	ASP	-	expression tag	UNP P49685
R	375	ASP	-	expression tag	UNP P49685
R	376	LYS	-	expression tag	UNP P49685

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	A	220	1699	1094	287	305	13	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	CYS	SER	conflict	UNP P08754
A	202	THR	GLY	conflict	UNP P08754
A	203	ALA	GLY	conflict	UNP P08754
A	245	ALA	GLU	conflict	UNP P08754
A	326	SER	ALA	conflict	UNP P08754

- Molecule 4 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	338	Total	C	N	O	S	0	0
			2496	1546	445	485	20		

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	MET	-	initiating methionine	UNP P62873
B	-9	HIS	-	expression tag	UNP P62873
B	-8	HIS	-	expression tag	UNP P62873
B	-7	HIS	-	expression tag	UNP P62873
B	-6	HIS	-	expression tag	UNP P62873
B	-5	HIS	-	expression tag	UNP P62873
B	-4	HIS	-	expression tag	UNP P62873
B	-3	GLY	-	expression tag	UNP P62873
B	-2	SER	-	expression tag	UNP P62873
B	-1	LEU	-	expression tag	UNP P62873
B	0	LEU	-	expression tag	UNP P62873
B	1	GLN	-	expression tag	UNP P62873

- Molecule 5 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	C	54	Total	C	N	O	S	0	0
			357	226	61	69	1		

- Molecule 6 is a protein called scFv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	D	234	Total	C	N	O	S	0	0
			1753	1115	293	335	10		



### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein GPR15LG



- Molecule 2: G-protein coupled receptor 15



- Molecule 3: Guanine nucleotide-binding protein G(i) subunit alpha-3



- Molecule 4: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1






- Molecule 5: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



- Molecule 6: scFv16



## 4 Experimental information i

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	286873	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	70	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	L	0.18	0/109	0.46	0/151
2	R	0.12	0/2291	0.33	0/3116
3	A	0.09	0/1728	0.25	0/2326
4	B	0.08	0/2543	0.24	0/3461
5	C	0.08	0/363	0.20	0/501
6	D	0.09	0/1796	0.26	0/2439
All	All	0.10	0/8830	0.28	0/11994

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	105	0	103	6	0
2	R	2233	0	2309	38	0
3	A	1699	0	1643	9	0
4	B	2496	0	2326	48	0
5	C	357	0	309	3	0
6	D	1753	0	1678	32	0
All	All	8643	0	8368	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:122:LEU:HB3	2:R:157:ILE:HD11	1.53	0.89
2:R:187:LYS:HG3	2:R:188:ALA:H	1.54	0.70
6:D:166:LEU:HB2	6:D:176:LEU:HD11	1.74	0.70
2:R:178:ASP:HB2	2:R:181:PRO:HG3	1.73	0.69
3:A:241:ASN:ND2	3:A:286:CYS:SG	2.65	0.69

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	12/58 (21%)	7 (58%)	5 (42%)	0	100 100
2	R	285/379 (75%)	268 (94%)	17 (6%)	0	100 100
3	A	214/354 (60%)	211 (99%)	3 (1%)	0	100 100
4	B	336/351 (96%)	323 (96%)	13 (4%)	0	100 100
5	C	52/71 (73%)	52 (100%)	0	0	100 100
6	D	230/267 (86%)	221 (96%)	9 (4%)	0	100 100
All	All	1129/1480 (76%)	1082 (96%)	47 (4%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	11/51 (22%)	11 (100%)	0	100	100
2	R	243/333 (73%)	243 (100%)	0	100	100
3	A	170/311 (55%)	170 (100%)	0	100	100
4	B	254/293 (87%)	254 (100%)	0	100	100
5	C	28/58 (48%)	28 (100%)	0	100	100
6	D	188/216 (87%)	187 (100%)	1 (0%)	86	91
All	All	894/1262 (71%)	893 (100%)	1 (0%)	92	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	D	220	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	241	ASN
4	B	91	HIS
4	B	293	ASN
6	D	39	GLN
6	D	167	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.