wwPDB EM Validation Summary Report

Nov 9, 2026 – 12:48 PM JST

PDB ID	:	9XO7 / pdb_00009xo7
EMDB ID	:	EMD-67072
Title	:	Cryo-EM structure of ABTS-1 in complex with HCO3 bound states
Authors	:	You, Z.L.; Bai, L.
Deposited on	:	2025-11-13
Resolution	:	2.94 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13068 (2.44 - 3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	5% 73% 27%
1	B	591	5% 73% 27%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BCT	A	1105	-	-	X	-
2	BCT	B	1106	-	-	X	-
4	CL	A	1103	-	-	X	-
4	CL	B	1104	-	-	X	-
6	POV	A	1106	-	-	X	-

2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 9606 atoms, of which 168 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anion exchange protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	591	Total 4643	C 3064	N 748	O 797	S 34	0	0
1	B	591	Total 4643	C 3064	N 748	O 797	S 34	0	0

- Molecule 2 is BICARBONATE ION (CCD ID: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total 5	C 1	H 1	O 3	0
2	A	1	Total 5	C 1	H 1	O 3	0
2	B	1	Total 5	C 1	H 1	O 3	0
2	B	1	Total 5	C 1	H 1	O 3	0


- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	AltConf
3	A	1	Total Na 1 1	0
3	B	1	Total Na 1 1	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	AltConf
4	A	1	Total Cl 1 1	0
4	B	1	Total Cl 1 1	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	AltConf
5	A	1	Total C N O 14 8 1 5	0
5	B	1	Total C N O 14 8 1 5	0

- Molecule 6 is (2S)-3-(hexadecanoyloxy)-2-[(9Z)-octadec-9-enoyloxy]propyl 2-(trimethylammonio)ethyl phosphate (CCD ID: POV) (formula: $C_{42}H_{82}NO_8P$).




Mol	Chain	Residues	Atoms						AltConf
6	A	1	Total	C	H	N	O	P	0
			134	42	82	1	8	1	
6	B	1	Total	C	H	N	O	P	0
			134	42	82	1	8	1	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Anion exchange protein


Chain A:



• Molecule 1: Anion exchange protein

Chain B:




L707 K708 K709 M710 R711 N712 Y715 F716 R721 A732 I733 A734 S735 M736 T737 F738 I741 F742 V743 G744 V745 V752 T759 D771 A778 P779 I782 L783 P784 A785 L786 I792 D795 I798 T799 I802 V803 N804 R805 G813 C814 G815 V822 V830
L835 P836 I837 A840 A841 T842 V843 K853 W854 E855 C858 K859 A860 P861 G862 E863 V864 A865 V870 R871 E872 Q873 R874 G877 I878 V879 L882 I883 I884 L894 P898 N899 P900 V901 L902 V905 F906 L907 A913 L914 G915 G916 I917 Q918 L919 R922 T923 L924
L925 M928 Y932 D935 T936 I937 Y938 I939 R940 H941 V942 F950 T951 Q954 A959 V963 I964 K968 S969 I972 L973 M977 I984 R985 K986 M987 M988 E989 D996 L997 L1000 D1001 D1002 P1003 M1004 P1005 H1008 L1009 R1010 E1013 D1014 A1015 K1016 R1017 R1018
Q1019 S1020 E1021 G1022 E1023

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	503518	Depositor
Resolution determination method	OTHER	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI F30	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.292	Depositor
Minimum map value	-3.208	Depositor
Average map value	-0.003	Depositor
Map value standard deviation	0.097	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	291.19998, 291.19998, 291.19998	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.04, 1.04, 1.04	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BCT, NA, NAG, POV, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.18	0/4759	0.34	0/6462
1	B	0.19	0/4759	0.34	0/6462
All	All	0.19	0/9518	0.34	0/12924

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4643	0	4788	159	0
1	B	4643	0	4788	155	0
2	A	8	2	1	3	0
2	B	8	2	1	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	4	0
4	B	1	0	0	4	0
5	A	14	0	13	1	0
5	B	14	0	13	1	0
6	A	52	82	82	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	52	82	82	20	0
All	All	9438	168	9768	302	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 302 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:843:VAL:HG12	4:A:1103:CL:CL	1.47	1.49
1:B:843:VAL:HG12	4:B:1104:CL:CL	1.47	1.49
1:A:695:LEU:CD2	1:B:695:LEU:CD2	2.15	1.23
1:B:843:VAL:CG1	4:B:1104:CL:CL	2.24	1.22
1:A:843:VAL:CG1	4:A:1103:CL:CL	2.24	1.21

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/591 (100%)	561 (95%)	28 (5%)	0	100	100
1	B	589/591 (100%)	561 (95%)	28 (5%)	0	100	100
All	All	1178/1182 (100%)	1122 (95%)	56 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	511/511 (100%)	509 (100%)	2 (0%)	84 92
1	B	511/511 (100%)	509 (100%)	2 (0%)	84 92
All	All	1022/1022 (100%)	1018 (100%)	4 (0%)	81 92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	870	VAL
1	A	924	LEU
1	B	870	VAL
1	B	924	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	611	GLN
1	B	797	GLN
1	B	918	GLN
1	B	808	ASN
1	B	873	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	A	1104	1	14,14,15	0.66	0	17,19,21	1.85	3 (17%)
2	BCT	B	1106	-	2,3,3	1.31	0	2,3,3	4.16	2 (100%)
6	POV	B	1101	-	51,51,51	0.81	1 (1%)	57,59,59	1.04	3 (5%)
6	POV	A	1106	-	51,51,51	0.81	1 (1%)	57,59,59	1.04	3 (5%)
2	BCT	A	1105	-	2,3,3	1.31	0	2,3,3	4.15	2 (100%)
5	NAG	B	1105	1	14,14,15	0.66	0	17,19,21	1.86	3 (17%)
2	BCT	B	1102	3	2,3,3	1.70	1 (50%)	2,3,3	2.23	1 (50%)
2	BCT	A	1101	3	2,3,3	1.71	1 (50%)	2,3,3	2.24	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	POV	A	1106	-	-	22/55/55/55	-
5	NAG	A	1104	1	-	3/6/23/26	0/1/1/1
6	POV	B	1101	-	-	22/55/55/55	-
5	NAG	B	1105	1	-	3/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1101	POV	C22-C21	-2.21	1.44	1.50
6	A	1106	POV	C22-C21	-2.19	1.44	1.50
2	A	1101	BCT	O2-C	-2.07	1.15	1.33
2	B	1102	BCT	O2-C	-2.06	1.15	1.33

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1106	BCT	O2-C-O1	5.52	133.87	119.55
2	A	1105	BCT	O2-C-O1	5.51	133.83	119.55
5	A	1104	NAG	C4-C3-C2	-5.45	103.03	111.02
5	B	1105	NAG	C4-C3-C2	-5.45	103.03	111.02
6	A	1106	POV	O21-C21-O22	4.57	134.74	123.70

There are no chirality outliers.

5 of 50 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	1106	POV	C12-C11-O12-P
6	B	1101	POV	C12-C11-O12-P
5	A	1104	NAG	O5-C5-C6-O6
5	B	1105	NAG	O5-C5-C6-O6
6	A	1106	POV	C11-C12-N-C14

There are no ring outliers.

8 monomers are involved in 50 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1104	NAG	1	0
2	B	1106	BCT	2	0
6	B	1101	POV	20	0
6	A	1106	POV	22	0
2	A	1105	BCT	2	0
5	B	1105	NAG	1	0
2	B	1102	BCT	1	0
2	A	1101	BCT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-67072. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 140




Y Index: 140




Z Index: 140


6.2.2 Raw map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 143




Y Index: 152




Z Index: 143


6.3.2 Raw map



X Index: 140



Y Index: 127




Z Index: 125


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 116 nm^3; this corresponds to an approximate mass of 104 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.340 $Å^{-1}$

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.340 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
	0.143	0.5	Half-bit	Other
Reported by author	-	-	-	2.94
Author-provided FSC curve	-	-	-	-
Unmasked-calculated*	3.52	3.94	3.58	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-67072 and PDB model 9XO7. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

9.4 Atom inclusion (i)



At the recommended contour level, 86% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.



Chain	Atom inclusion	Q-score
All	0.8440	0.4740
A	0.8460	0.4740
B	0.8460	0.4740