



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:41 AM UTC

PDB ID : 5XQO / pdb\_00005xqo  
Title : Crystal structure of a PL 26 exo-rhamnogalacturonan lyase from *Penicillium chrysogenum* complexed with tetrameric substrate  
Authors : Kunishige, Y.; Iwai, M.; Tada, T.; Nishimura, S.; Sakamoto, T.  
Deposited on : 2017-06-07  
Resolution : 3.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	906	 82% 16% ..
1	B	906	 79% 17% ..
2	C	4	 25% 75%
3	D	4	 75% 25%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 14092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Perglx protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	899	7017	4476	1178	1357	3	3	0	0	0
1	B	881	6870	4392	1153	1319	3	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	458	PHE	TYR	engineered mutation	UNP A0A0C6EFY4
B	458	PHE	TYR	engineered mutation	UNP A0A0C6EFY4

- Molecule 2 is an oligosaccharide called 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-2)-alpha-L-rhamnopyranose-(1-4)-alpha-D-galactopyranuronic acid-(1-2)-alpha-L-rhamnopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	4	44	24	20	0	0	0

- Molecule 3 is an oligosaccharide called 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-3)-alpha-L-rhamnopyranose-(1-4)-alpha-D-galactopyranuronic acid-(1-2)-alpha-L-rhamnopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
3	D	4	Total	C O	0	0	0
			44	24 20			


- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	62	Total	O	0	0
			62	62		
5	B	53	Total	O	0	0
			53	53		






- Molecule 2: 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-2)-alpha-L-rhamnopyranose-(1-4)-alpha-D-galactopyranuronic acid-(1-2)-alpha-L-rhamnopyranose



- Molecule 3: 2,6-anhydro-3-deoxy-L-threo-hex-2-enonic acid-(1-3)-alpha-L-rhamnopyranose-(1-4)-alpha-D-galactopyranuronic acid-(1-2)-alpha-L-rhamnopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.44Å 166.44Å 171.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	119.58 – 3.20 119.58 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (119.58-3.20) 99.5 (119.58-3.20)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.40 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, $R_{free}$	0.191 , 0.264 0.195 , 0.260	Depositor DCC
$R_{free}$ test set	2100 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	87.9	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k 0.000 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	14092	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, RAM, GAD, ADA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.74	0/7218	0.95	9/9854 (0.1%)
1	B	0.79	1/7069 (0.0%)	0.98	6/9655 (0.1%)
All	All	0.76	1/14287 (0.0%)	0.97	15/19509 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	278	LEU	CA-C	5.39	1.59	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	302	VAL	N-CA-C	-7.31	99.51	109.37
1	A	661	LEU	N-CA-C	-6.62	100.64	110.23
1	A	24	CYS	N-CA-C	6.42	120.29	109.76
1	A	756	GLY	N-CA-C	5.67	121.22	111.98
1	A	891	VAL	N-CA-C	5.58	113.77	109.19

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7017	0	6566	70	0
1	B	6870	0	6391	73	0
2	C	44	0	32	5	0
3	D	44	0	26	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	62	0	0	3	0
5	B	53	0	0	6	0
All	All	14092	0	13015	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:24:CYS:SG	5:B:1114:HOH:O	2.34	0.84
1:A:157:ILE:HA	1:A:204:ILE:HD11	1.70	0.73
1:B:288:ILE:HD13	1:B:349:TYR:CE1	2.29	0.67
1:A:708:LEU:HG	1:A:755:SER:HA	1.78	0.66
1:B:825:GLU:CB	5:B:1149:HOH:O	2.46	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	893/906 (99%)	805 (90%)	74 (8%)	14 (2%)	7	36
1	B	873/906 (96%)	773 (88%)	77 (9%)	23 (3%)	4	26
All	All	1766/1812 (98%)	1578 (89%)	151 (9%)	37 (2%)	5	31

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	695	GLY
1	A	748	THR
1	A	774	ASN
1	B	231	ASP
1	B	525	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	719/742 (97%)	675 (94%)	44 (6%)	17	49
1	B	694/742 (94%)	655 (94%)	39 (6%)	19	52
All	All	1413/1484 (95%)	1330 (94%)	83 (6%)	18	50

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	252	LYS
1	B	487	THR
1	B	282	GLU
1	B	331	THR
1	B	591	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	178	GLN
1	B	391	GLN
1	B	189	ASN
1	B	774	ASN
1	B	369	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RAM	C	1	2	11,11,11	0.63	0	16,16,16	1.48	5 (31%)
2	ADA	C	2	2	12,12,13	0.76	0	14,17,19	2.50	4 (28%)
2	RAM	C	3	2	10,10,11	0.75	0	14,14,16	1.57	3 (21%)
2	GAD	C	4	2	10,11,11	2.37	2 (20%)	12,15,15	2.65	4 (33%)
3	RAM	D	1	3	11,11,11	0.90	0	16,16,16	2.29	6 (37%)
3	ADA	D	2	3	12,12,13	0.79	0	14,17,19	1.65	2 (14%)
3	RAM	D	3	3	10,10,11	3.06	3 (30%)	14,14,16	5.15	9 (64%)
3	GAD	D	4	3	10,11,11	2.72	3 (30%)	12,15,15	4.19	5 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RAM	C	1	2	-	-	0/1/1/1
2	ADA	C	2	2	-	0/4/21/24	0/1/1/1
2	RAM	C	3	2	-	-	0/1/1/1
2	GAD	C	4	2	-	0/4/17/17	0/1/1/1
3	RAM	D	1	3	-	-	0/1/1/1
3	ADA	D	2	3	-	2/4/21/24	0/1/1/1
3	RAM	D	3	3	-	-	0/1/1/1
3	GAD	D	4	3	-	4/4/17/17	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	3	RAM	C2-C3	8.39	1.65	1.52
3	D	4	GAD	O5-C5	7.56	1.47	1.37
2	C	4	GAD	O5-C5	6.52	1.46	1.37
3	D	3	RAM	O3-C3	3.41	1.51	1.43
3	D	3	RAM	C4-C3	2.60	1.59	1.52

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3	RAM	O3-C3-C2	12.57	135.71	110.05
3	D	3	RAM	C1-C2-C3	9.94	124.11	109.64
3	D	4	GAD	O5-C5-C4	-9.46	116.26	124.94
2	C	2	ADA	O4-C4-C5	7.40	126.65	109.76
2	C	4	GAD	O5-C5-C4	-6.86	118.65	124.94

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	4	GAD	C4-C5-C6-O6B
3	D	4	GAD	O5-C5-C6-O6A
3	D	4	GAD	O5-C5-C6-O6B
3	D	4	GAD	C4-C5-C6-O6A
3	D	2	ADA	O5-C5-C6-O6B

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	RAM	3	0
3	D	4	GAD	1	0
2	C	3	RAM	2	0
2	C	2	ADA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	896/906 (98%)	-0.55	0 <a href="#">100</a>   <a href="#">100</a>	62, 84, 111, 156	0
1	B	878/906 (96%)	-0.43	2 (0%) <a href="#">91</a>   <a href="#">85</a>	60, 90, 119, 159	0
All	All	1774/1812 (97%)	-0.49	2 (0%) <a href="#">92</a>   <a href="#">89</a>	60, 87, 117, 159	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	228	ASP	3.3
1	B	263	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GAD	C	4	11/11	0.72	0.10	89,104,110,113	0
3	GAD	D	4	11/11	0.81	0.07	100,114,134,138	0
3	RAM	D	3	10/11	0.89	0.14	100,139,152,168	0
3	RAM	D	1	11/11	0.89	0.10	99,103,108,120	0
3	ADA	D	2	12/13	0.91	0.07	81,102,118,125	0
2	RAM	C	1	11/11	0.93	0.10	73,76,83,84	0
2	ADA	C	2	12/13	0.94	0.06	73,84,88,103	0
2	RAM	C	3	10/11	0.96	0.10	104,106,115,125	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CA	A	1001	1/1	0.99	0.02	44,44,44,44	0
4	CA	B	1001	1/1	0.99	0.04	57,57,57,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.