



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 12:43 AM UTC

PDB ID : 6XYL / pdb\_00006xyl  
Title : Crystal structure of delta466-491 cystathionine beta-synthase from *Toxoplasma gondii* with L-serine  
Authors : Fernandez-Rodriguez, C.; Oyenarte, I.; Conter, C.; Gonzalez-Recio, I.; Quintana, I.; Martinez-Chantar, M.; Astegno, A.; Martinez-Cruz, L.A.  
Deposited on : 2020-01-30  
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2361 (3.20-3.12)
Clashscore	190562	2486 (3.20-3.12)
Ramachandran outliers	187476	2405 (3.20-3.12)
Sidechain outliers	187428	2404 (3.20-3.12)
RSRZ outliers	180081	2361 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	514	 69% 21% 9%
1	B	514	 68% 23% 9%
1	C	514	 67% 25% 9%
1	D	514	 69% 24% 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	E	514	 70% 22% 7%
1	F	514	 73% 19% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	P1T	A	601	-	-	X	-

## 2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 21482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cystathionine beta-synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	Total 3528	C 2214	N 620	O 668	S 26	0	0	0
1	B	469	Total 3540	C 2222	N 622	O 670	S 26	0	0	0
1	C	470	Total 3546	C 2225	N 622	O 674	S 25	0	0	0
1	D	475	Total 3576	C 2241	N 627	O 682	S 26	0	0	0
1	E	476	Total 3586	C 2247	N 629	O 684	S 26	0	0	0
1	F	475	Total 3580	C 2244	N 628	O 682	S 26	0	0	0

- Molecule 2 is 2-[(3-HYDROXY-2-METHYL-5-[(PHOSPHONOXY)METHYL]PYRIDIN-4-YL)METHYL)AMINO]ACRYLIC ACID (CCD ID: P1T) (formula: C<sub>11</sub>H<sub>15</sub>N<sub>2</sub>O<sub>7</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	21	11	2	7	1	0	0
2	B	1	21	11	2	7	1	0	0
2	C	1	21	11	2	7	1	0	0
2	D	1	21	11	2	7	1	0	0
2	E	1	21	11	2	7	1	0	0
2	F	1	21	11	2	7	1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cystathionine beta-synthase



- Molecule 1: Cystathionine beta-synthase








- Molecule 1: Cystathionine beta-synthase

Chain F:  73%  19%  8%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.44Å 82.44Å 421.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.02 – 3.15 41.02 – 3.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (41.02-3.15) 99.4 (41.02-3.15)	Depositor EDS
$R_{merge}$	0.32	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.18 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.274 , 0.294 0.276 , 0.295	Depositor DCC
$R_{free}$ test set	1980 reflections (3.58%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	85.0	Xtrriage
Anisotropy	0.537	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 40.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.019 for -h,-k,l 0.458 for h,-h-k,-l 0.026 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	21482	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P1T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.16	0/3580	0.37	1/4827 (0.0%)
1	B	0.15	0/3593	0.34	0/4845
1	C	0.14	0/3597	0.36	1/4848 (0.0%)
1	D	0.13	0/3630	0.33	0/4899
1	E	0.12	0/3640	0.32	0/4913
1	F	0.15	0/3634	0.34	0/4903
All	All	0.14	0/21674	0.34	2/29235 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	500	ILE	N-CA-C	6.29	116.92	108.11
1	A	87	THR	CB-CA-C	6.03	122.16	110.46

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3528	0	3560	100	0
1	B	3540	0	3572	88	0
1	C	3546	0	3582	102	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3576	0	3591	88	0
1	E	3586	0	3601	87	0
1	F	3580	0	3602	81	0
2	A	21	0	12	7	0
2	B	21	0	12	2	0
2	C	21	0	12	5	0
2	D	21	0	12	5	0
2	E	21	0	12	5	0
2	F	21	0	12	5	0
All	All	21482	0	21580	523	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 523 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:393:VAL:HG23	1:F:406:VAL:CG1	1.56	1.33
1:B:423:CYS:SG	1:B:432:ILE:HG12	1.85	1.15
1:F:393:VAL:CG2	1:F:406:VAL:HG13	1.83	1.08
1:A:56:LYS:HB3	1:A:87:THR:CG2	1.90	1.00
1:A:56:LYS:HB3	1:A:87:THR:HG22	1.43	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	460/514 (90%)	449 (98%)	11 (2%)	0	100 100
1	B	461/514 (90%)	449 (97%)	12 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	462/514 (90%)	451 (98%)	11 (2%)	0	100	100
1	D	469/514 (91%)	460 (98%)	9 (2%)	0	100	100
1	E	470/514 (91%)	459 (98%)	11 (2%)	0	100	100
1	F	469/514 (91%)	454 (97%)	15 (3%)	0	100	100
All	All	2791/3084 (90%)	2722 (98%)	69 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/417 (89%)	372 (100%)	1 (0%)	86	85
1	B	375/417 (90%)	375 (100%)	0	100	100
1	C	375/417 (90%)	374 (100%)	1 (0%)	86	85
1	D	378/417 (91%)	378 (100%)	0	100	100
1	E	379/417 (91%)	379 (100%)	0	100	100
1	F	379/417 (91%)	378 (100%)	1 (0%)	86	85
All	All	2259/2502 (90%)	2256 (100%)	3 (0%)	88	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	C	497	LEU
1	F	406	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	384	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	153	ASN
1	D	86	ASN
1	C	161	ASN
1	D	318	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	P1T	F	601	-	20,21,21	1.51	3 (15%)	29,30,30	1.35	6 (20%)
2	P1T	B	601	-	20,21,21	1.51	3 (15%)	29,30,30	1.34	6 (20%)
2	P1T	E	601	-	20,21,21	1.51	3 (15%)	29,30,30	1.35	6 (20%)
2	P1T	D	601	-	20,21,21	1.52	3 (15%)	29,30,30	1.35	6 (20%)
2	P1T	A	601	-	20,21,21	1.52	3 (15%)	29,30,30	1.35	6 (20%)
2	P1T	C	601	-	20,21,21	1.52	3 (15%)	29,30,30	1.35	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P1T	F	601	-	-	3/14/15/15	0/1/1/1
2	P1T	B	601	-	-	4/14/15/15	0/1/1/1
2	P1T	E	601	-	-	3/14/15/15	0/1/1/1
2	P1T	D	601	-	-	3/14/15/15	0/1/1/1
2	P1T	A	601	-	-	1/14/15/15	0/1/1/1
2	P1T	C	601	-	-	3/14/15/15	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	P1T	C5-C4	-3.21	1.35	1.40
2	C	601	P1T	C5-C4	-3.20	1.35	1.40
2	D	601	P1T	C5-C4	-3.18	1.36	1.40
2	F	601	P1T	C5-C4	-3.17	1.36	1.40
2	B	601	P1T	C5-C4	-3.17	1.36	1.40

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	P1T	O-C-CA	2.71	120.07	114.25
2	D	601	P1T	C4A-N-CA	-2.70	120.01	125.68
2	E	601	P1T	C4A-N-CA	-2.70	120.02	125.68
2	D	601	P1T	O-C-CA	2.70	120.04	114.25
2	C	601	P1T	C4A-N-CA	-2.70	120.03	125.68

There are no chirality outliers.

5 of 17 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	601	P1T	C5-C4-C4A-N
2	B	601	P1T	C-CA-N-C4A
2	C	601	P1T	C-CA-N-C4A
2	D	601	P1T	C-CA-N-C4A
2	E	601	P1T	C-CA-N-C4A


There are no ring outliers.


6 monomers are involved in 29 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	601	P1T	5	0
2	B	601	P1T	2	0
2	E	601	P1T	5	0
2	D	601	P1T	5	0
2	A	601	P1T	7	0
2	C	601	P1T	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	468/514 (91%)	-0.83	0 100 100	60, 72, 123, 141	0
1	B	469/514 (91%)	-0.69	1 (0%) 91 84	65, 85, 127, 149	0
1	C	470/514 (91%)	-0.70	0 100 100	63, 84, 127, 138	0
1	D	475/514 (92%)	-0.76	0 100 100	65, 83, 126, 142	1 (0%)
1	E	476/514 (92%)	-0.74	0 100 100	65, 85, 124, 140	0
1	F	475/514 (92%)	-0.86	0 100 100	59, 73, 129, 139	0
All	All	2833/3084 (91%)	-0.76	1 (0%) 100 100	59, 81, 126, 149	1 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	406	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	P1T	C	601	21/21	0.98	0.06	71,72,73,74	0
2	P1T	D	601	21/21	0.98	0.06	69,70,72,73	0
2	P1T	E	601	21/21	0.98	0.05	70,71,72,73	0
2	P1T	B	601	21/21	0.99	0.06	69,73,76,77	0
2	P1T	A	601	21/21	0.99	0.05	62,63,63,64	0
2	P1T	F	601	21/21	0.99	0.06	60,61,62,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around P1T D 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around P1T E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around P1T B 601:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around P1T A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.