

wwPDB X-ray Structure Validation Summary Report (i)

Sep 3, 2025 – 10:09 AM EDT

PDB ID : 9Y12 / pdb 00009y12

Title : Crystal Structure of N-Acetyl Transferase Domain-Containing Protein from

Bacteroides fragilis

Authors: Kim, Y.; Bigelow, L.; Clancy, L.; Abdullah, J.; Joachimiak, A.; Midwest

Center for Structural Genomics (MCSG)

Deposited on : 2025-08-29

Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 2.0rc1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.45.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	154	88%	10%	
1	В	154	86%	12%	:
1	С	154	86%	12%	
1	D	154	88%	10%	



Continued from previous page...

001000		v precedue	page		
Mol	Chain	Length	Quality of chain		
1	Е	154	82%	16%	
1	F	154	87%	12%	-
1	G	154	81%	17%	
1	Н	154	81%	17%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 10683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acetyltransferase (GNAT) family protein.

Mol	Chain	Residues		_	Atom	ıs			ZeroOcc	AltConf	Trace	
1	A	152	Total	С	N	О	S	Se	0	1	0	
1	A	152	1277	819	215	235	5	3	0	1		
1	В	152	Total	С	N	О	S	Se	0	0	0	
1	Ъ	152	1266	813	211	234	5	3	0	0		
1	С	152	Total	С	N	О	S	Se	0	0	0	
1		152	1266	813	211	234	5	3	0	U	0	
1	D	152	Total	С	N	О	S	Se	0	0	0	
1	ט	152	1266	813	211	234	5	3	0	O		
1	Е	152	Total	С	N	О	S	Se	0	0	0	
1	l Li	152	1266	813	211	234	5	3	0	0		
1	F	152	Total	С	N	О	S	Se	0	1	0	
1	I.	102	1275	818	212	237	5	3	U	1		
1	G	152	Total	С	N	О	S	Se	0	1	0	
1	G	102	1274	818	212	235	5	4		1	U	
1	Н	152	Total	С	N	О	S	Se	0	0	0	
1	11	102	1266	813	211	234	5	3	U	U	U	

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A0A149N1Y8
A	-1	ASN	-	expression tag	UNP A0A149N1Y8
A	0	ALA	-	expression tag	UNP A0A149N1Y8
В	-2	SER	-	expression tag	UNP A0A149N1Y8
В	-1	ASN	-	expression tag	UNP A0A149N1Y8
В	0	ALA	-	expression tag	UNP A0A149N1Y8
С	-2	SER	-	expression tag	UNP A0A149N1Y8
С	-1	ASN	-	expression tag	UNP A0A149N1Y8
С	0	ALA	-	expression tag	UNP A0A149N1Y8
D	-2	SER	-	expression tag	UNP A0A149N1Y8
D	-1	ASN	-	expression tag	UNP A0A149N1Y8
D	0	ALA	-	expression tag	UNP A0A149N1Y8
Е	-2	SER	-	expression tag	UNP A0A149N1Y8



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-1	ASN	-	expression tag	UNP A0A149N1Y8
Е	0	ALA	-	expression tag	UNP A0A149N1Y8
F	-2	SER	-	expression tag	UNP A0A149N1Y8
F	-1	ASN	-	expression tag	UNP A0A149N1Y8
F	0	ALA	-	expression tag	UNP A0A149N1Y8
G	-2	SER	-	expression tag	UNP A0A149N1Y8
G	-1	ASN	-	expression tag	UNP A0A149N1Y8
G	0	ALA	-	expression tag	UNP A0A149N1Y8
Н	-2	SER	-	expression tag	UNP A0A149N1Y8
Н	-1	ASN	_	expression tag	UNP A0A149N1Y8
Н	0	ALA	-	expression tag	UNP A0A149N1Y8

 \bullet Molecule 2 is COENZYME A (CCD ID: COA) (formula: $\mathrm{C_{21}H_{36}N_7O_{16}P_3S}).$




Mol	Chain	Residues		Atoms						AltConf	
2	A	1	Total	С	N	О	Р	S	0	0	
2	Λ	1	48	21	7	16	3	1	0		
2	В	1	Total	С	N	О	Р	S	0	0	
	Ъ	1	48	21	7	16	3	1	U	U	
2	\mathbf{C}	1	Total	С	N	Ο	Р	\mathbf{S}	0	0	
2		1	48	21	7	16	3	1	U	U	
2	D	1	Total	С	N	Ο	Р	S	0	0	
	D	1	48	21	7	16	3	1	U	U	
2	E	1	Total	С	N	О	Р	S	0	0	
	L	1	48	21	7	16	3	1	U	0	
2	F	1	Total	С	N	Ο	Р	S	0	0	
	1	1	48	21	7	16	3	1	U		



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
9	C	1	Total	С	N	О	Р	S	0	0	
	G	1	48	21	7	16	3	1	U	. 0	
9	П	1	Total	С	N	О	Р	S	0	0	
2	П	1	48	21	7	16	3	1	0		

 $\bullet\,$ Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	С	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
3	D	1	Total	О	S	0	0
	D	1	5	4	1	0	0
3	D	1	Total	Ο	S	0	0
		_	5	4	1	Ü	
3	E	1	Total	O	S	0	0
	_	_	5	4	1		
3	Е	1	Total	O	S	0	0
			5	4	1		_
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	Н	1	Total	O	S	0	0
			5	4	1		
3	Н	1	Total	O	S	0	0
			5	4	1		


 \bullet Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $\mathrm{C_2H_6O_2}).$





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	В	1	Total 4	C 2	O 2	0	0

 \bullet Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: $\mathrm{C_2H_3O_2}).$



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	Е	1	Total 4	C 2	O 2	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total O 3 3	0	0
6	В	4	Total O 4 4	0	0
6	С	4	Total O 4 4	0	0
6	D	2	Total O 2 2	0	0
6	Е	7	Total O 7 7	0	0
6	F	6	Total O 6 6	0	0
6	G	3	Total O 3 3	0	0
6	Н	6	Total O 6 6	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetyltransferase (GNAT) family protein






• Molecule 1: Acetyltransferase (GNAT) family protein



• Molecule 1: Acetyltransferase (GNAT) family protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	92.96Å 92.96Å 218.51Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.48 - 2.50	Depositor
Resolution (A)	46.48 - 2.50	EDS
% Data completeness	99.3 (46.48-2.50)	Depositor
(in resolution range)	$100.0 \ (46.48 - 2.50)$	EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.93 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
P. P.	0.189 , 0.224	Depositor
R, R_{free}	0.193 , 0.195	DCC
R_{free} test set	3894 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.33 \; , 19.5$	EDS
L-test for twinning ²	$< L > = 0.43, < L^2> = 0.25$	Xtriage
	0.050 for -h,-k,l	
Estimated twinning fraction	0.457 for h,-h-k,-l	Xtriage
	0.054 for -k,-h,-l	
Reported twinning fraction	0.500 for k,h,-l	Depositor
Outliers	0 of 81309 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10683	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, COA, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.12	0/1311	0.26	0/1773
1	В	0.11	0/1300	0.25	0/1759
1	С	0.11	0/1300	0.26	0/1759
1	D	0.10	0/1300	0.25	0/1759
1	Е	0.11	0/1300	0.27	0/1759
1	F	0.11	0/1309	0.25	0/1771
1	G	0.12	0/1308	0.25	0/1769
1	Н	0.11	0/1300	0.26	0/1759
All	All	0.11	0/10428	0.25	0/14108

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1277	0	1203	9	0
1	В	1266	0	1191	11	0
1	С	1266	0	1191	11	0
1	D	1266	0	1191	9	0
1	Е	1266	0	1191	16	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1275	0	1196	11	0
1	G	1274	0	1199	15	0
1	Н	1266	0	1191	15	0
2	A	48	0	32	1	0
2	В	48	0	32	2	0
2	С	48	0	32	0	0
2	D	48	0	32	0	0
2	Ε	48	0	32	1	0
2	F	48	0	32	2	0
2	G	48	0	32	0	0
2	Н	48	0	32	0	0
3	A	20	0	0	0	0
3	В	15	0	0	0	0
3	С	5	0	0	0	0
3	D	15	0	0	0	0
3	Ε	10	0	0	0	0
3	F	15	0	0	0	0
3	G	10	0	0	0	0
3	Н	10	0	0	0	0
4	В	4	0	6	0	0
5	Ε	4	0	3	0	0
6	A	3	0	0	0	0
6	В	4	0	0	0	0
6	С	4	0	0	0	0
6	D	2	0	0	0	0
6	Е	7	0	0	0	0
6	F	6	0	0	0	0
6	G	3	0	0	0	0
6	Н	6	0	0	0	0
All	All	10683	0	9818	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:F:6:LEU:HG	1:F:41:LEU:HD21	1.77	0.67
1:G:6:LEU:HG	1:G:41:LEU:HD21	1.79	0.64
1:E:6:LEU:HG	1:E:41:LEU:HD21	1.85	0.59
1:E:129:ASN:HB2	1:E:149:LYS:HE2	1.88	0.56



Continued from previous page...

Atom-1	1100111 1		$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$	
1:E:55:ASN:HB3	1:E:61:ILE:HD13	1.86	0.56	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	151/154 (98%)	148 (98%)	3 (2%)	0	100	100
1	В	150/154 (97%)	147 (98%)	3 (2%)	0	100	100
1	С	150/154 (97%)	144 (96%)	6 (4%)	0	100	100
1	D	150/154 (97%)	145 (97%)	5 (3%)	0	100	100
1	E	150/154 (97%)	145 (97%)	5 (3%)	0	100	100
1	F	151/154 (98%)	143 (95%)	8 (5%)	0	100	100
1	G	151/154 (98%)	146 (97%)	5 (3%)	0	100	100
1	Н	150/154 (97%)	145 (97%)	5 (3%)	0	100	100
All	All	1203/1232 (98%)	1163 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	$134/132\ (102\%)$	131 (98%)	3 (2%)	47	73	
1	В	$133/132\ (101\%)$	132 (99%)	1 (1%)	79	91	
1	C	133/132 (101%)	132 (99%)	1 (1%)	79	91	
1	D	$133/132\ (101\%)$	131 (98%)	2 (2%)	60	82	
1	E	133/132 (101%)	130 (98%)	3 (2%)	45	72	
1	F	$134/132\ (102\%)$	132 (98%)	2 (2%)	60	82	
1	G	$134/132\ (102\%)$	129 (96%)	5 (4%)	29	55	
1	Н	133/132 (101%)	128 (96%)	5 (4%)	28	53	
All	All	1067/1056 (101%)	1045 (98%)	22 (2%)	48	74	

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	27	GLU
1	Н	8	LYS
1	G	61	ILE
1	Н	30	ARG
1	D	68	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	Н	69	ASN
1	G	148	GLN
1	F	93	HIS
1	Е	148	GLN
1	G	114	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / L - 1	T	Cl :-	D	T ! 1-	Вс	ond leng	ths	Bond angles			
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	SO4	A	203	-	4,4,4	0.66	0	6,6,6	0.10	0	
4	EDO	В	202	-	3,3,3	0.25	0	2,2,2	0.33	0	
2	COA	A	201	-	43,50,50	0.62	0	56,75,75	0.93	2 (3%)	
3	SO4	С	202	-	4,4,4	0.66	0	6,6,6	0.10	0	
5	ACT	Е	202	-	3,3,3	1.09	0	3,3,3	1.13	0	
3	SO4	Н	202	_	4,4,4	0.67	0	6,6,6	0.10	0	
3	SO4	В	203	_	4,4,4	0.67	0	6,6,6	0.10	0	
2	COA	D	201	-	43,50,50	0.64	0	56,75,75	1.08	3 (5%)	
3	SO4	F	202	-	4,4,4	0.67	0	6,6,6	0.09	0	
3	SO4	A	202	-	4,4,4	0.67	0	6,6,6	0.08	0	
3	SO4	Е	203	-	4,4,4	0.66	0	6,6,6	0.11	0	
3	SO4	F	203	-	4,4,4	0.67	0	6,6,6	0.07	0	
3	SO4	D	204	_	4,4,4	0.67	0	6,6,6	0.08	0	
3	SO4	F	204	_	4,4,4	0.67	0	6,6,6	0.15	0	
3	SO4	D	202	_	4,4,4	0.67	0	6,6,6	0.09	0	
2	COA	В	201	_	43,50,50	0.63	0	56,75,75	0.90	2 (3%)	
2	COA	Е	201	-	43,50,50	0.63	0	56,75,75	1.02	3 (5%)	
2	COA	G	201	-	43,50,50	0.63	0	56,75,75	0.93	2 (3%)	
3	SO4	G	202	-	4,4,4	0.67	0	6,6,6	0.07	0	
3	SO4	G	203	-	4,4,4	0.67	0	6,6,6	0.06	0	
3	SO4	В	205	-	4,4,4	0.66	0	6,6,6	0.12	0	
3	SO4	A	205	-	4,4,4	0.66	0	6,6,6	0.08	0	
2	COA	F	201	-	43,50,50	0.63	0	56,75,75	0.96	2 (3%)	
3	SO4	Н	203	-	4,4,4	0.67	0	6,6,6	0.09	0	
3	SO4	D	203	-	4,4,4	0.66	0	6,6,6	0.08	0	
3	SO4	A	204	-	4,4,4	0.67	0	6,6,6	0.07	0	



Mol Type	Tuno	Chain	Res	Link	Во	ond leng	ths	Bond angles		
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	В	204	-	4,4,4	0.67	0	6,6,6	0.09	0
2	COA	С	201	-	43,50,50	0.63	0	56,75,75	0.95	2 (3%)
3	SO4	Е	204	-	4,4,4	0.67	0	6,6,6	0.07	0
2	COA	Н	201	-	43,50,50	0.64	0	56,75,75	0.96	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	В	201	-	-	13/44/64/64	0/3/3/3
2	COA	Е	201	-	-	11/44/64/64	0/3/3/3
2	COA	G	201	-	-	2/44/64/64	0/3/3/3
4	EDO	В	202	-	-	0/1/1/1	-
2	COA	A	201	-	-	12/44/64/64	0/3/3/3
2	COA	С	201	-	-	9/44/64/64	0/3/3/3
2	COA	D	201	-	-	4/44/64/64	0/3/3/3
2	COA	Н	201	-	-	7/44/64/64	0/3/3/3
2	COA	F	201	-	-	13/44/64/64	0/3/3/3

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	Е	201	COA	P3B-O3B-C3B	-5.27	109.36	123.43
2	F	201	COA	P3B-O3B-C3B	-5.01	110.04	123.43
2	Н	201	COA	P3B-O3B-C3B	-4.96	110.18	123.43
2	A	201	COA	P3B-O3B-C3B	-4.91	110.33	123.43
2	С	201	COA	P3B-O3B-C3B	-4.87	110.43	123.43

There are no chirality outliers.

5 of 71 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	COA	OAP-CAP-CBP-CCP
2	A	201	COA	C9P-CAP-CBP-CCP
2	A	201	COA	OAP-CAP-CBP-CDP
2	A	201	COA	C9P-CAP-CBP-CDP



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	201	COA	C9P-CAP-CBP-CEP


There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	COA	1	0
2	В	201	COA	2	0
2	Е	201	COA	1	0
2	F	201	COA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#	#RSR	Z>2	$OWAB(A^2)$	Q < 0.9
1	A	149/154 (96%)	-1.75	0	100	100	23, 42, 61, 104	1 (0%)
1	В	149/154 (96%)	-1.77	0	100	100	34, 42, 63, 140	0
1	С	149/154 (96%)	-1.75	0	100	100	34, 47, 65, 88	0
1	D	149/154 (96%)	-1.72	0	100	100	39, 49, 71, 94	0
1	E	149/154 (96%)	-1.72	0	100	100	31, 47, 64, 116	0
1	F	149/154 (96%)	-1.75	0	100	100	25, 46, 59, 72	1 (0%)
1	G	149/154 (96%)	-1.73	0	100	100	35, 48, 70, 80	0
1	Н	149/154 (96%)	-1.73	0	100	100	40, 49, 69, 107	0
All	All	1192/1232 (96%)	-1.74	0	100	100	23, 46, 67, 140	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	COA	D	201	48/48	0.99	0.03	56,70,99,105	0
2	COA	Н	201	48/48	0.99	0.04	56,72,86,89	0
3	SO4	A	202	5/5	0.99	0.03	57,61,80,88	0
3	SO4	A	205	5/5	0.99	0.06	53,58,64,72	0
3	SO4	В	204	5/5	0.99	0.03	46,55,69,71	0
3	SO4	С	202	5/5	0.99	0.04	59,60,66,67	0
3	SO4	D	202	5/5	0.99	0.04	60,61,73,79	0
3	SO4	D	203	5/5	0.99	0.03	55,60,75,78	0
3	SO4	D	204	5/5	0.99	0.04	53,58,64,70	0
3	SO4	Е	204	5/5	0.99	0.04	65,65,86,92	0
3	SO4	F	204	5/5	0.99	0.03	59,61,73,79	0
3	SO4	G	202	5/5	0.99	0.06	66,69,77,81	0
3	SO4	G	203	5/5	0.99	0.03	69,75,80,90	0
3	SO4	Н	202	5/5	0.99	0.03	68,69,82,89	0
3	SO4	Н	203	5/5	0.99	0.04	65,66,69,88	0
4	EDO	В	202	4/4	0.99	0.04	39,42,42,43	0
5	ACT	Е	202	4/4	0.99	0.05	36,41,43,46	0
2	COA	В	201	48/48	1.00	0.03	40,49,76,85	0
2	COA	С	201	48/48	1.00	0.03	48,58,85,88	0
3	SO4	Е	203	5/5	1.00	0.03	52,60,67,69	0
3	SO4	A	203	5/5	1.00	0.04	46,51,57,64	5
3	SO4	F	202	5/5	1.00	0.03	46,48,67,71	0
3	SO4	F	203	5/5	1.00	0.04	57,64,79,90	0
3	SO4	A	204	5/5	1.00	0.03	59,60,69,72	0
2	COA	A	201	48/48	1.00	0.03	35,44,76,79	0
3	SO4	В	203	5/5	1.00	0.03	48,48,60,71	0
2	COA	E	201	48/48	1.00	0.03	40,56,71,79	0
3	SO4	В	205	5/5	1.00	0.02	53,54,59,62	0
2	COA	F	201	48/48	1.00	0.03	42,58,76,80	0
2	COA	G	201	48/48	1.00	0.03	52,59,75,78	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around COA D 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around COA H 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around COA C 201: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)














6.5 Other polymers (i)

There are no such residues in this entry.

