



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:46 AM UTC

PDB ID : 8Y4K / pdb_00008y4k
Title : Metal Beta-Lactamase VIM-2 in Complex with MBL inhibitor B17
Authors : Li, G.-B.; Wang, S.-Y.
Deposited on : 2024-01-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	 10% 84% 16%
1	B	231	 10% 73% 24% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metal-Beta-Lactamase VIM-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	231	Total 1733	C 1094	N 299	O 339	S 1	0	0	0
1	B	231	Total 1733	C 1094	N 299	O 339	S 1	0	0	0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	2	Total 2	Zn 2	0	0

- Molecule 3 is 2-[[[(3 {R})-6-(azetidino-3-yloxy)-1,1-bis(oxidanylo)-3 {H}-2,1-benzoxaborol-1-ium-3-yl)methyl]prop-2-enoic acid (CCD ID: A1LXG) (formula: C₁₄H₁₇BNO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	B	C	N	O	0	0
			22	1	14	1	6		
3	B	1	Total	B	C	N	O	0	0
			22	1	14	1	6		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			7	4 3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total O	0	0
			3 3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metal-Beta-Lactamase VIM-2



- Molecule 1: Metal-Beta-Lactamase VIM-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.31Å 62.95Å 68.35Å 90.00° 100.80° 90.00°	Depositor
Resolution (Å)	67.14 – 2.50 67.14 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (67.14-2.50) 99.8 (67.14-2.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.21-5207	Depositor
R, R_{free}	0.199 , 0.278 0.211 , 0.286	Depositor DCC
R_{free} test set	1523 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å ²)	37.6	Xtrriage
Anisotropy	0.460	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3524	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1LXG, ZN, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/1774	0.86	1/2429 (0.0%)
1	B	0.77	0/1774	0.84	2/2429 (0.1%)
All	All	0.76	0/3548	0.85	3/4858 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	209	GLY	N-CA-C	-9.06	100.97	111.85
1	A	210	ASN	N-CA-C	7.04	120.38	107.99
1	B	208	ALA	N-CA-C	5.24	121.96	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1733	0	1682	12	0
1	B	1733	0	1682	24	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	22	0	0	0	0
3	B	22	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	7	0	10	1	0
5	A	3	0	0	0	0
All	All	3524	0	3374	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:GLU:OE2	1:B:102:GLN:NE2	2.19	0.76
1:B:117:ASP:OD1	1:B:148:ASN:ND2	2.19	0.73
4:A:304:PEG:O1	3:B:303:A1LXG:N19	2.28	0.66
1:B:116:HIS:NE2	1:B:213:ASP:OD2	2.22	0.66
1:B:210:ASN:HB2	3:B:303:A1LXG:O09	1.97	0.64
1:A:49:ILE:HD13	1:A:237:ILE:HD11	1.83	0.60
1:A:98:GLU:OE2	1:A:102:GLN:NE2	2.29	0.60
1:B:178:ALA:HB1	1:B:219:TRP:CD1	2.38	0.58
1:A:60:GLN:HB3	1:A:87:TRP:CE3	2.41	0.56
1:B:141:ARG:O	1:B:144:GLU:HB3	2.07	0.55
1:B:84:ASP:OD1	1:B:112:SER:HA	2.08	0.54
1:B:60:GLN:HB3	1:B:87:TRP:CE3	2.43	0.53
1:A:34:PRO:HB2	1:A:39:ILE:HD11	1.91	0.52
1:B:56:HIS:NE2	1:B:71:GLY:HA3	2.25	0.52
1:B:80:LEU:HD11	1:B:103:ILE:HG21	1.94	0.50
1:A:228:GLN:O	1:A:231:PRO:HD3	2.13	0.47
1:B:85:THR:HG23	1:B:119:ARG:O	2.14	0.47
1:A:82:LEU:O	1:A:110:ALA:HA	2.14	0.47
1:B:96:LEU:HD11	1:B:126:LEU:HD23	1.96	0.47
1:B:34:PRO:HG2	1:B:57:ILE:HD13	1.96	0.47
1:A:201:TYR:CD2	1:A:205:ARG:HD3	2.49	0.46
1:B:60:GLN:HB3	1:B:87:TRP:CZ3	2.52	0.44
1:A:258:ALA:O	1:A:259:HIS:C	2.61	0.44
1:B:225:ARG:O	1:B:229:HIS:HB2	2.18	0.43
1:B:144:GLU:HA	1:B:150:ILE:HD11	1.99	0.43
1:B:179:HIS:O	1:B:214:ALA:HA	2.19	0.42
1:B:135:ALA:HB2	1:B:151:PRO:HG2	2.01	0.42
1:B:208:ALA:HB2	1:B:259:HIS:CE1	2.54	0.42
1:A:42:GLY:HA2	1:A:91:ASN:OD1	2.20	0.42
1:B:105:LEU:HB3	1:B:106:PRO:HD2	2.01	0.42
1:A:172:LEU:HD22	1:A:186:VAL:HG22	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:ASP:CG	1:B:79:GLU:HG3	2.45	0.41
1:B:86:ALA:O	1:B:91:ASN:HB3	2.21	0.41
1:B:73:ILE:HG12	1:B:82:LEU:HD12	2.03	0.41
1:A:134:TYR:HA	1:A:153:HIS:O	2.21	0.41
1:B:121:GLY:HA2	1:B:149:GLU:OE2	2.22	0.40
1:A:201:TYR:HE2	1:A:208:ALA:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/231 (99%)	210 (92%)	17 (7%)	2 (1%)	14	27
1	B	229/231 (99%)	202 (88%)	16 (7%)	11 (5%)	2	2
All	All	458/462 (99%)	412 (90%)	33 (7%)	13 (3%)	4	6

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	113	THR
1	B	145	VAL
1	B	208	ALA
1	A	178	ALA
1	B	127	ARG
1	B	84	ASP
1	B	152	THR
1	B	144	GLU
1	B	130	GLY
1	A	84	ASP
1	B	87	TRP
1	B	178	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	49	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/182 (100%)	169 (93%)	13 (7%)	13	29
1	B	182/182 (100%)	170 (93%)	12 (7%)	15	32
All	All	364/364 (100%)	339 (93%)	25 (7%)	14	30

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ILE
1	A	51	ASP
1	A	54	TRP
1	A	72	LEU
1	A	123	VAL
1	A	159	SER
1	A	183	ASN
1	A	190	SER
1	A	216	LEU
1	A	224	GLU
1	A	242	LEU
1	A	253	THR
1	A	260	THR
1	B	48	GLN
1	B	51	ASP
1	B	114	HIS
1	B	123	VAL
1	B	145	VAL
1	B	146	GLU
1	B	153	HIS
1	B	160	SER
1	B	170	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	183	ASN
1	B	232	GLU
1	B	248	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	ASN
1	A	229	HIS
1	B	261	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	A	304	-	6,6,6	0.55	0	5,5,5	0.60	0
3	A1LXG	A	303	2	21,24,24	4.20	5 (23%)	21,36,36	2.46	6 (28%)
3	A1LXG	B	303	2	21,24,24	4.51	9 (42%)	21,36,36	3.14	9 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	304	-	-	2/4/4/4	-
3	A1LXG	A	303	2	-	3/12/33/33	0/3/3/3
3	A1LXG	B	303	2	-	4/12/33/33	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	303	A1LXG	C12-C22	-12.01	1.32	1.39
3	A	303	A1LXG	B02-O01	10.30	1.67	1.48
3	B	303	A1LXG	B02-O03	10.29	1.67	1.48
3	A	303	A1LXG	C12-C22	-10.24	1.33	1.39
3	B	303	A1LXG	B02-O01	9.45	1.66	1.48
3	A	303	A1LXG	B02-O03	9.39	1.66	1.48
3	A	303	A1LXG	C08-C07	6.91	1.54	1.49
3	B	303	A1LXG	C06-C07	5.65	1.57	1.50
3	B	303	A1LXG	B02-O04	-3.79	1.35	1.45
3	B	303	A1LXG	C08-C07	3.44	1.51	1.49
3	B	303	A1LXG	C13-C12	2.97	1.43	1.39
3	B	303	A1LXG	C18-C17	-2.23	1.49	1.53
3	B	303	A1LXG	C20-C17	-2.16	1.49	1.53
3	A	303	A1LXG	C18-C17	-2.01	1.50	1.53

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	A1LXG	C11-C07-C08	-10.06	112.01	120.65
3	A	303	A1LXG	O09-C08-C07	6.51	122.15	114.89
3	A	303	A1LXG	O10-C08-C07	-5.48	116.58	121.74
3	B	303	A1LXG	C06-C07-C08	5.03	122.13	116.16
3	B	303	A1LXG	C15-O16-C17	-4.08	111.79	119.18
3	B	303	A1LXG	O09-C08-C07	4.05	119.40	114.89
3	B	303	A1LXG	C13-C12-C22	-3.79	119.64	121.92
3	A	303	A1LXG	C13-C12-C22	-3.37	119.90	121.92
3	A	303	A1LXG	C11-C07-C08	3.28	123.47	120.65
3	B	303	A1LXG	O10-C08-C07	-3.25	118.68	121.74
3	A	303	A1LXG	O01-B02-O04	2.92	116.98	110.09
3	A	303	A1LXG	C06-C05-C12	-2.18	105.55	114.49
3	B	303	A1LXG	C21-C22-C12	2.18	119.73	118.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	A1LXG	O04-C05-C06	-2.16	107.07	111.55
3	B	303	A1LXG	O03-B02-O04	-2.12	105.07	110.09

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	303	A1LXG	C18-C17-O16-C15
3	B	303	A1LXG	O04-C05-C06-C07
4	A	304	PEG	O2-C3-C4-O4
3	B	303	A1LXG	C14-C15-O16-C17
3	B	303	A1LXG	C21-C15-O16-C17
3	B	303	A1LXG	C12-C05-C06-C07
3	A	303	A1LXG	C20-C17-O16-C15
4	A	304	PEG	C4-C3-O2-C2
3	A	303	A1LXG	O04-C05-C06-C07

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	304	PEG	1	0
3	B	303	A1LXG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/231 (100%)	0.00	1 (0%) 88 86	20, 34, 49, 64	0
1	B	231/231 (100%)	0.77	22 (9%) 14 12	25, 48, 76, 88	0
All	All	462/462 (100%)	0.39	23 (4%) 34 30	20, 39, 72, 88	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	113	THR	4.4
1	B	262	ALA	3.7
1	A	262	ALA	3.6
1	B	151	PRO	3.3
1	B	125	VAL	3.2
1	B	103	ILE	3.0
1	B	131	VAL	2.9
1	B	107	VAL	2.7
1	B	81	LEU	2.6
1	B	105	LEU	2.6
1	B	153	HIS	2.5
1	B	96	LEU	2.5
1	B	126	LEU	2.5
1	B	94	ALA	2.3
1	B	134	TYR	2.3
1	B	132	ALA	2.3
1	B	133	THR	2.3
1	B	108	THR	2.2
1	B	102	GLN	2.1
1	B	80	LEU	2.1
1	B	122	GLY	2.1
1	B	111	VAL	2.0
1	B	123	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PEG	A	304	7/7	0.85	0.20	48,57,87,125	0
3	A1LXG	A	303	22/22	0.86	0.14	34,51,64,75	0
3	A1LXG	B	303	22/22	0.90	0.11	15,31,40,50	0
2	ZN	B	302	1/1	0.99	0.02	27,27,27,27	0
2	ZN	A	301	1/1	0.99	0.02	37,37,37,37	0
2	ZN	A	302	1/1	0.99	0.02	35,35,35,35	0
2	ZN	B	301	1/1	0.99	0.03	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1LXG A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1LXG B 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.