



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 16, 2024 – 06:14 PM JST

PDB ID : 8YBO
Title : Crystal structure of nanobody SEB-Nb11 bound to staphylococcal enterotoxin B (SEB)
Authors : Ding, Y.; Zong, X.; Liu, R.; Liu, P.
Deposited on : 2024-02-15
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 11% 86% 5% 10%
1	C	240	 9% 84% 5% 10%
2	B	128	 18% 94% 6%
2	D	128	 17% 94% 5%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enterotoxin type B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	217	1818	1167	289	352	10	0	0	0
1	C	216	1810	1163	287	350	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP P01552
C	2	MET	-	initiating methionine	UNP P01552

- Molecule 2 is a protein called nanobody SEB-Nb11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	128	983	621	166	192	4	0	0	0
2	D	128	983	621	166	192	4	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total 120	O 120	0	0
3	B	65	Total 65	O 65	0	0
3	C	147	Total 147	O 147	0	0
3	D	75	Total 75	O 75	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enterotoxin type B




- Molecule 1: Enterotoxin type B



- Molecule 2: nanobody SEB-Nb11



- Molecule 2: nanobody SEB-Nb11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.15Å 82.96Å 83.55Å 90.00° 97.44° 90.00°	Depositor
Resolution (Å)	82.84 – 1.59 82.84 – 1.59	Depositor EDS
% Data completeness (in resolution range)	98.1 (82.84-1.59) 98.2 (82.84-1.59)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.59Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.261 , 0.284 0.268 , 0.293	Depositor DCC
R_{free} test set	5494 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtrriage
Anisotropy	0.161	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 25.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6001	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9671e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1860	0.60	0/2503
1	C	0.42	0/1852	0.59	0/2492
2	B	0.35	0/1007	0.58	0/1368
2	D	0.35	0/1007	0.59	0/1368
All	All	0.39	0/5726	0.59	0/7731

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	2
2	B	0	1
2	D	0	1
All	All	0	5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	ARG	Sidechain
2	B	19	ARG	Sidechain
1	C	132	ARG	Sidechain
1	C	137	ARG	Sidechain
2	D	19	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1818	0	1745	5	0
1	C	1810	0	1739	10	0
2	B	983	0	957	4	0
2	D	983	0	957	4	0
3	A	120	0	0	0	0
3	B	65	0	0	1	0
3	C	147	0	0	5	0
3	D	75	0	0	0	0
All	All	6001	0	5398	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:67:ARG:HD2	3:C:371:HOH:O	1.83	0.77
1:C:67:ARG:CD	3:C:371:HOH:O	2.33	0.77
1:C:97:PHE:CG	3:C:371:HOH:O	2.38	0.76
1:C:97:PHE:CD2	3:C:371:HOH:O	2.49	0.62
2:B:11:LEU:HD11	2:B:127:SER:HB3	1.84	0.60
1:A:9:LYS:HB3	1:A:10:PRO:CD	2.32	0.59
1:A:97:PHE:O	1:A:98:SER:HB2	2.07	0.54
1:C:9:LYS:HB3	1:C:10:PRO:HD2	1.89	0.54
1:C:97:PHE:O	1:C:98:SER:HB2	2.07	0.54
2:D:22:CYS:HG	2:D:96:CYS:HG	1.56	0.53
2:B:22:CYS:HG	2:B:96:CYS:CB	2.23	0.51
2:B:22:CYS:SG	2:B:96:CYS:SG	3.02	0.51
1:A:9:LYS:HB3	1:A:10:PRO:HD2	1.91	0.51
1:C:132:ARG:HD2	1:C:228:LYS:HA	1.95	0.48
1:C:67:ARG:HD3	3:C:371:HOH:O	2.09	0.48
2:B:89:GLU:HG2	3:B:232:HOH:O	2.15	0.47
1:A:128:LEU:HD11	1:A:228:LYS:HZ1	1.80	0.46
2:D:19:ARG:HG2	2:D:82:GLN:OE1	2.16	0.46
1:A:41:LYS:CG	1:A:81:ASP:HA	2.45	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:11:LEU:HD11	2:D:127:SER:HB3	1.98	0.45
2:D:22:CYS:SG	2:D:96:CYS:SG	3.09	0.45
1:C:41:LYS:CG	1:C:81:ASP:HA	2.46	0.45
1:C:89:ALA:H	1:C:160:GLN:HE21	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/240 (88%)	205 (97%)	6 (3%)	0	100	100
1	C	210/240 (88%)	205 (98%)	5 (2%)	0	100	100
2	B	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
2	D	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
All	All	673/736 (91%)	660 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/226 (90%)	201 (98%)	3 (2%)	60	39
1	C	203/226 (90%)	199 (98%)	4 (2%)	50	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	106/106 (100%)	104 (98%)	2 (2%)	52	29
2	D	106/106 (100%)	104 (98%)	2 (2%)	52	29
All	All	619/664 (93%)	608 (98%)	11 (2%)	54	31

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	31	ASP
1	A	97	PHE
2	B	77	ASN
2	B	107	THR
1	C	31	ASP
1	C	97	PHE
1	C	129	ASP
1	C	209	LYS
2	D	77	ASN
2	D	107	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	160	GLN
2	D	13	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/240 (90%)	0.74	26 (11%) 10 9	9, 17, 41, 59	0
1	C	216/240 (90%)	0.66	22 (10%) 13 12	9, 17, 40, 67	0
2	B	128/128 (100%)	1.01	23 (17%) 4 3	13, 21, 44, 65	0
2	D	128/128 (100%)	0.84	22 (17%) 5 4	12, 19, 39, 54	0
All	All	689/736 (93%)	0.78	93 (13%) 8 7	9, 19, 41, 67	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	41	PRO	5.7
1	C	63	TYR	5.6
1	A	3	GLU	5.6
1	A	62	ASN	5.2
1	A	81	ASP	5.2
1	C	96	TYR	5.0
1	A	10	PRO	4.9
1	A	98	SER	4.9
2	B	118	TRP	4.8
1	C	98	SER	4.8
1	A	96	TYR	4.5
2	B	117	TYR	4.5
1	C	97	PHE	4.4
2	D	2	VAL	4.4
2	D	118	TRP	4.4
2	D	40	ALA	4.4
2	B	2	VAL	4.4
1	C	3	GLU	4.2
2	B	128	SER	4.2
1	A	11	ASP	4.1
1	C	10	PRO	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	1	GLN	4.1
1	A	130	LYS	4.0
2	B	127	SER	3.8
2	D	117	TYR	3.7
1	C	130	LYS	3.7
2	D	42	GLY	3.7
1	C	195	GLU	3.7
1	C	55	ILE	3.7
1	A	129	ASP	3.7
2	B	1	GLN	3.6
1	A	97	PHE	3.6
1	A	128	LEU	3.5
2	B	5	VAL	3.5
1	A	239	LYS	3.5
1	A	63	TYR	3.4
1	A	132	ARG	3.2
2	B	23	ALA	3.2
1	A	55	ILE	3.2
2	D	45	ARG	3.1
1	C	129	ASP	3.1
2	D	128	SER	3.1
1	A	9	LYS	3.0
1	C	145	LEU	2.9
2	B	42	GLY	2.9
2	B	3	GLN	2.9
1	C	4	SER	2.9
1	C	194	ASN	2.9
1	C	9	LYS	2.8
1	A	4	SER	2.8
2	B	54	GLY	2.8
2	B	112	LEU	2.8
2	B	56	SER	2.7
1	C	209	LYS	2.7
1	A	194	ASN	2.7
2	D	111	LEU	2.7
1	C	11	ASP	2.7
1	C	196	ASN	2.7
1	A	82	LYS	2.7
1	C	239	LYS	2.6
1	A	131	TYR	2.6
1	A	228	LYS	2.5
1	A	6	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	LEU	2.5
1	C	131	TYR	2.4
1	A	209	LYS	2.4
2	D	127	SER	2.4
2	B	88	THR	2.4
2	B	41	PRO	2.4
2	D	75	ALA	2.4
2	B	65	LYS	2.3
2	B	75	ALA	2.3
1	C	82	LYS	2.3
2	D	112	LEU	2.3
2	B	126	VAL	2.3
2	D	3	GLN	2.3
2	D	54	GLY	2.3
2	D	65	LYS	2.3
1	C	81	ASP	2.2
2	D	100	LYS	2.2
2	B	44	GLY	2.2
1	A	195	GLU	2.2
2	D	107	THR	2.1
2	D	89	GLU	2.1
1	A	196	ASN	2.1
1	C	128	LEU	2.1
2	B	100	LYS	2.1
2	D	19	ARG	2.1
2	D	26	GLY	2.1
2	B	63	SER	2.1
2	B	45	ARG	2.1
2	B	76	LYS	2.1
2	D	43	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.