



wwPDB EM Validation Summary Report ⓘ

Jul 16, 2025 – 12:05 PM JST

PDB ID : 8YGN / pdb_00008ygn
EMDB ID : EMD-39257
Title : The Dimer Structure of DSR2-SPR with NAD
Authors : Gao, X.; Zhu, H.; Cui, S.
Deposited on : 2024-02-26
Resolution : 4.27 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : **FAILED**
MolProbity : 4-5-2 with Phenix2.0rc1
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	C	264	
1	D	264	
2	A	1005	
2	B	1005	
2	E	1005	
2	F	1005	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 43648 atoms, of which 20406 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SPR.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	D	151	Total	C	N	O	S	0	0
			1198	758	189	247	4		
1	C	151	Total	C	N	O	S	0	0
			1198	758	189	247	4		

- Molecule 2 is a protein called SIR2-like domain-containing protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	A	983	Total	C	H	N	O	S	0	0
			16194	5292	8011	1322	1538	31		
2	B	983	Total	C	H	N	O	S	0	0
			16194	5292	8011	1322	1538	31		
2	E	277	Total	C	H	N	O	S	0	0
			4470	1455	2210	367	430	8		
2	F	272	Total	C	H	N	O	S	0	0
			4394	1432	2174	360	421	7		


There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	171	ALA	HIS	engineered mutation	UNP D4G637
B	171	ALA	HIS	engineered mutation	UNP D4G637
E	171	ALA	HIS	engineered mutation	UNP D4G637
F	171	ALA	HIS	engineered mutation	UNP D4G637

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SPR

Chain D: 



- Molecule 1: SPR

Chain C: 



- Molecule 2: SIR2-like domain-containing protein

Chain A: 

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	14482	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	66	Depositor
Minimum defocus (nm)	1800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.19	0/1218	0.54	0/1645
1	D	0.18	0/1218	0.52	0/1645
2	A	0.19	0/8374	0.47	7/11280 (0.1%)
2	B	0.20	0/8374	0.42	1/11280 (0.0%)
2	E	0.12	0/2314	0.30	0/3136
2	F	0.13	0/2274	0.31	0/3082
All	All	0.18	0/23772	0.43	8/32068 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	859	VAL	CA-C-N	8.93	138.59	121.54
2	A	859	VAL	C-N-CA	8.93	138.59	121.54
2	A	858	SER	CA-C-N	8.27	136.86	121.97
2	A	858	SER	C-N-CA	8.27	136.86	121.97
2	A	299	ASN	N-CA-C	-7.54	103.34	112.54

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	297	GLN	Peptide
2	B	297	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1198	0	1159	165	0
1	D	1198	0	1159	152	0
2	A	8183	8011	8010	128	0
2	B	8183	8011	8010	122	0
2	E	2260	2210	2209	26	0
2	F	2220	2174	2173	35	0
All	All	23242	20406	22720	584	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 584 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:196:PRO:HG3	1:D:230:GLU:HA	1.47	0.95
1:D:50:LEU:HD12	1:D:51:TYR:H	1.31	0.95
1:D:189:SER:HB3	1:D:238:SER:HB3	1.49	0.93
1:D:11:VAL:HG13	1:D:175:TYR:HE1	1.35	0.91
1:D:45:ILE:HG23	1:D:47:ASN:H	1.37	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	147/264 (56%)	109 (74%)	30 (20%)	8 (5%)	1	16
1	D	147/264 (56%)	105 (71%)	35 (24%)	7 (5%)	2	17
2	A	981/1005 (98%)	931 (95%)	46 (5%)	4 (0%)	30	67
2	B	981/1005 (98%)	935 (95%)	45 (5%)	1 (0%)	48	83
2	E	275/1005 (27%)	270 (98%)	5 (2%)	0	100	100
2	F	270/1005 (27%)	263 (97%)	7 (3%)	0	100	100
All	All	2801/4548 (62%)	2613 (93%)	168 (6%)	20 (1%)	21	56

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	4	VAL
1	D	26	GLU
1	D	49	PRO
1	D	50	LEU
1	C	4	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	130/225 (58%)	130 (100%)	0	100	100
1	D	130/225 (58%)	130 (100%)	0	100	100
2	A	901/922 (98%)	899 (100%)	2 (0%)	92	93
2	B	901/922 (98%)	900 (100%)	1 (0%)	92	94
2	E	249/922 (27%)	249 (100%)	0	100	100
2	F	244/922 (26%)	244 (100%)	0	100	100
All	All	2555/4138 (62%)	2552 (100%)	3 (0%)	92	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	295	GLU
2	A	298	GLU
2	B	861	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	263	ASN
2	B	863	ASN
2	B	548	ASN
2	B	978	HIS
2	A	332	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.