



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 28, 2024 – 10:09 AM JST

PDB ID : 8YJ2
Title : N17.1.2 recognition of NRAS neoantigens
Authors : Wu, D.C.; Mariuzza, R.A.
Deposited on : 2024-02-29
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 5% 89% 10% •
2	B	100	 3% 89% 11% •
3	C	10	 80% 20%
4	E	247	 2% 89% 9% ••
5	D	207	 4% 84% 13% •

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called human leukocyte antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2228	1384	406	428	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP F6IQR9

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	824	524	138	158	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	expression tag	UNP P61769

- Molecule 3 is a protein called ILE-LEU-ASP-THR-ALA-GLY-ARG-GLU-GLU-TYR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	10	82	50	13	19	0	0	0

- Molecule 4 is a protein called tcr beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	245	1968	1246	339	375	8	0	0	0

- Molecule 5 is a protein called tcr alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	D	202	1578	995	259	315	9	0	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	110	Total 110	O 110	0	0
6	B	48	Total 48	O 48	0	0
6	C	9	Total 9	O 9	0	0
6	E	163	Total 163	O 163	0	0
6	D	96	Total 96	O 96	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: human leukocyte antigen




- Molecule 2: Beta-2-microglobulin




- Molecule 3: ILE-LEU-ASP-THR-ALA-GLY-ARG-GLU-GLU-TYR




- Molecule 4: tcr beta



- Molecule 5: tcr alpha





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	71.52Å 134.16Å 222.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.78 – 2.26 42.78 – 2.26	Depositor EDS
% Data completeness (in resolution range)	98.5 (42.78-2.26) 98.4 (42.78-2.26)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.27Å)	Xtrriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.189 , 0.229 0.192 , 0.228	Depositor DCC
R_{free} test set	2522 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtrriage
Anisotropy	0.449	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7106	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.48	0/2288	0.64	1/3102 (0.0%)
2	B	0.42	0/847	0.60	0/1150
3	C	0.44	0/82	0.52	0/108
4	E	0.54	2/2022 (0.1%)	0.63	2/2755 (0.1%)
5	D	0.48	0/1612	0.65	0/2184
All	All	0.49	2/6851 (0.0%)	0.63	3/9299 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	103	GLU	CD-OE1	-7.56	1.17	1.25
4	E	103	GLU	CD-OE2	-6.98	1.18	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ARG	NE-CZ-NH2	-9.01	115.79	120.30
4	E	112	ARG	NE-CZ-NH2	-5.28	117.66	120.30
4	E	207	ARG	NE-CZ-NH1	-5.04	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2079	17	1
2	B	824	0	769	10	0
3	C	82	0	77	1	0
4	E	1968	0	1901	21	0
5	D	1578	0	1521	31	1
6	A	110	0	0	4	0
6	B	48	0	0	3	0
6	C	9	0	0	0	0
6	D	96	0	0	7	0
6	E	163	0	0	5	2
All	All	7106	0	6347	74	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:102:LYS:HD2	4:E:102:LYS:H	1.14	1.13
4:E:244:ARG:NH1	6:E:301:HOH:O	1.81	1.13
5:D:148:GLN:NE2	6:D:301:HOH:O	1.86	1.01
1:A:176:LYS:NZ	6:A:301:HOH:O	1.93	1.01
1:A:231:VAL:O	1:A:243:LYS:NZ	1.93	1.00
5:D:65:VAL:CG1	5:D:74:VAL:HG23	2.04	0.86
2:B:20:SER:HA	2:B:71:THR:HG22	1.57	0.85
5:D:99:LYS:NZ	6:D:305:HOH:O	2.10	0.84
4:E:102:LYS:HD2	4:E:102:LYS:N	1.92	0.82
5:D:33:TYR:CE2	5:D:50:ARG:NH1	2.47	0.81
5:D:133:SER:O	6:D:302:HOH:O	1.99	0.81
5:D:34:ILE:HG21	5:D:74:VAL:HG11	1.68	0.76
4:E:35:TYR:OH	6:E:303:HOH:O	2.07	0.73
5:D:202:SER:O	6:D:304:HOH:O	2.07	0.71
2:B:84:HIS:O	6:B:101:HOH:O	2.09	0.70
1:A:82:ARG:NH1	6:A:303:HOH:O	2.25	0.69
5:D:33:TYR:HE2	5:D:50:ARG:NH1	1.90	0.69
5:D:65:VAL:HG11	5:D:74:VAL:HG23	1.74	0.69
1:A:186:LYS:NZ	6:A:305:HOH:O	2.26	0.68
5:D:193:ILE:O	6:D:306:HOH:O	2.15	0.64
4:E:101:PRO:HA	5:D:50:ARG:HH21	1.63	0.64
5:D:65:VAL:CG1	5:D:74:VAL:CG2	2.77	0.61
1:A:225:THR:HA	1:A:228:THR:HG21	1.85	0.59
4:E:102:LYS:HD3	4:E:103:GLU:HG3	1.87	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:119:LEU:HD13	4:E:219:LEU:HD22	1.87	0.57
4:E:100:LEU:HB3	4:E:101:PRO:HD2	1.87	0.56
5:D:65:VAL:HG13	5:D:74:VAL:HG23	1.85	0.56
1:A:225:THR:HA	1:A:228:THR:CG2	2.36	0.55
2:B:1:ILE:HA	6:B:107:HOH:O	2.08	0.54
2:B:24:ASN:HB3	2:B:65:LEU:HD11	1.91	0.53
5:D:74:VAL:HG22	5:D:74:VAL:O	2.08	0.53
5:D:65:VAL:HG11	5:D:74:VAL:CG2	2.38	0.51
5:D:148:GLN:NE2	5:D:149:SER:H	2.08	0.51
4:E:26:ASN:ND2	6:E:308:HOH:O	2.43	0.51
5:D:65:VAL:HG13	5:D:74:VAL:CG2	2.39	0.51
1:A:226:GLN:O	1:A:226:GLN:OE1	2.29	0.50
4:E:221:GLU:O	4:E:231:LYS:NZ	2.38	0.50
5:D:107:THR:OG1	6:D:307:HOH:O	2.19	0.49
1:A:214:THR:HB	1:A:262:GLN:HB2	1.95	0.49
4:E:56:ASP:OD1	6:E:304:HOH:O	2.20	0.49
5:D:34:ILE:HG21	5:D:74:VAL:CG1	2.41	0.49
5:D:155:TYR:O	5:D:176:ALA:HA	2.13	0.48
1:A:50:PRO:O	6:A:302:HOH:O	2.20	0.47
4:E:207:ARG:NH1	6:E:312:HOH:O	2.47	0.47
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.50	0.47
1:A:19:GLU:H	1:A:19:GLU:HG2	1.46	0.46
4:E:173:CYS:HB2	5:D:162:LEU:HB3	1.96	0.46
1:A:120:GLY:HA3	2:B:1:ILE:HD11	1.98	0.46
5:D:73:SER:OG	6:D:303:HOH:O	2.04	0.46
2:B:20:SER:CA	2:B:71:THR:HG22	2.37	0.46
5:D:24:LEU:CD2	5:D:73:SER:HB2	2.47	0.45
1:A:218:GLN:HA	1:A:222:GLU:O	2.17	0.45
1:A:111:ARG:HD3	1:A:113:TYR:OH	2.17	0.44
4:E:14:VAL:HA	4:E:116:LEU:O	2.17	0.44
4:E:102:LYS:N	4:E:102:LYS:CD	2.71	0.44
3:C:3:ASP:OD1	3:C:4:THR:N	2.51	0.43
5:D:38:LYS:HB2	5:D:48:LEU:HD11	2.00	0.43
2:B:81:ARG:NH2	6:B:109:HOH:O	2.50	0.43
4:E:101:PRO:CA	5:D:50:ARG:HH21	2.31	0.43
4:E:23:CYS:HB2	4:E:34:TRP:CZ2	2.54	0.43
4:E:101:PRO:HA	5:D:50:ARG:NH2	2.31	0.43
5:D:34:ILE:HG12	5:D:74:VAL:HG11	2.00	0.43
1:A:49:ALA:O	1:A:52:ILE:HG22	2.19	0.42
4:E:62:GLU:H	4:E:62:GLU:CD	2.22	0.42
1:A:197:HIS:C	1:A:251:SER:HB2	2.39	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:86:THR:HG23	4:E:112:ARG:HH22	1.84	0.42
5:D:137:PHE:O	5:D:173:SER:HA	2.19	0.42
5:D:126:ASP:OD1	5:D:127:SER:N	2.53	0.42
1:A:5:MET:O	1:A:100:GLY:HA3	2.20	0.41
2:B:25:CYS:HB2	2:B:39:LEU:HD21	2.01	0.41
5:D:197:ASP:OD1	5:D:197:ASP:N	2.37	0.41
4:E:182:GLN:O	4:E:188:SER:HB2	2.21	0.40
5:D:161:VAL:HG22	5:D:172:ASN:OD1	2.22	0.40
2:B:1:ILE:O	2:B:1:ILE:HG13	2.20	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:E:372:HOH:O	6:E:428:HOH:O[3_654]	2.05	0.15
1:A:108:ARG:NH1	5:D:129:SER:O[7_554]	2.07	0.13
6:E:430:HOH:O	6:E:445:HOH:O[3_654]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/276 (99%)	264 (97%)	9 (3%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
3	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
4	E	243/247 (98%)	237 (98%)	6 (2%)	0	100	100
5	D	200/207 (97%)	192 (96%)	8 (4%)	0	100	100
All	All	822/840 (98%)	797 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/233 (99%)	225 (98%)	5 (2%)	47	56
2	B	92/95 (97%)	91 (99%)	1 (1%)	70	79
3	C	8/8 (100%)	8 (100%)	0	100	100
4	E	220/222 (99%)	216 (98%)	4 (2%)	54	64
5	D	178/184 (97%)	176 (99%)	2 (1%)	70	79
All	All	728/742 (98%)	716 (98%)	12 (2%)	58	69

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	75	ARG
1	A	196	ASP
1	A	251	SER
1	A	268	LYS
2	B	70	PHE
4	E	71	LYS
4	E	102	LYS
4	E	112	ARG
4	E	195	ARG
5	D	129	SER
5	D	141	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	HIS
5	D	148	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/276 (99%)	0.12	14 (5%) 34 34	20, 35, 65, 137	0
2	B	100/100 (100%)	0.06	3 (3%) 52 53	23, 37, 55, 63	0
3	C	10/10 (100%)	-0.44	0 100 100	24, 27, 31, 34	0
4	E	245/247 (99%)	-0.33	4 (1%) 70 71	20, 28, 47, 62	0
5	D	202/207 (97%)	0.21	8 (3%) 43 43	24, 36, 60, 75	0
All	All	832/840 (99%)	-0.01	29 (3%) 47 48	20, 33, 59, 137	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	D	130	SER	5.5
1	A	17	ARG	4.8
1	A	16	GLY	4.3
5	D	131	ASP	4.2
1	A	227	ASP	3.5
1	A	225	THR	3.5
4	E	2	ALA	3.1
1	A	105	PRO	3.1
1	A	262	GLN	3.0
1	A	18	GLY	3.0
5	D	129	SER	3.0
1	A	194	ILE	3.0
2	B	98	ASP	2.9
1	A	232	GLU	2.7
2	B	0	MET	2.7
5	D	115	GLN	2.4
1	A	178	THR	2.3
5	D	181	SER	2.2
4	E	101	PRO	2.2
4	E	102	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
5	D	196	GLU	2.1
5	D	190	ASN	2.1
4	E	246	ASP	2.1
1	A	104	GLY	2.1
1	A	15	PRO	2.1
1	A	177	GLU	2.1
5	D	147	SER	2.1
2	B	1	ILE	2.1
1	A	110	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.