



wwPDB EM Validation Summary Report i

Jul 15, 2025 – 02:02 PM JST

PDB ID : 8YJP / pdb_00008yjp
EMDB ID : EMD-39345
Title : Cryo-EM structure of human apo GPR156
Authors : Ma, X.Y.; Chen, L.N.; Liao, M.H.; Zhang, L.Y.; Xi, K.; Guo, J.M.
Deposited on : 2024-03-02
Resolution : 3.09 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:


EMDB validation analysis	: FAILED
Mogul	: 1.8.5 (274361), CSD as541be (2020)
MolProbity	: 4-5-2 with Phenix2.0rc1
buster-report	: 1.1.7 (2018)
Percentile statistics	: 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	: FAILED
Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.44

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain			
1	A	346	63%	23%	14%	
1	B	346	62%	24%	14%	

2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 4946 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable G-protein coupled receptor 156.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	298	Total	C	N	O	S	0	0
			2241	1474	363	387	17		


Mol	Chain	Residues	Total	C	N	O	S	AltConf	Trace
1	B	298	Total	C	N	O	S	0	0
			2241	1474	363	387	17		

- Molecule 2 is (21R,24R,27S)-24,27,28-trihydroxy-18,24-dioxo-19,23,25-trioxa-24lambd 5 -p phosphaoctacosan-21-yl (9Z)-octadec-9-enoate (CCD ID: MW9) (formula: C₄₂H₈₁O₁₀P).



Mol	Chain	Residues	Total	C	O	P	AltConf
2	A	1	53	42	10	1	0
2	A	1	53	42	10	1	0
2	B	1	53	42	10	1	0
2	B	1	53	42	10	1	0

- Molecule 3 is CHOLESTEROL (CCD ID: CLR) (formula: C₂₇H₄₆O).




Mol	Chain	Residues	Atoms	AltConf
3	A	1	Total C O 28 27 1	0
3	A	1	Total C O 28 27 1	0
3	A	1	Total C O 28 27 1	0
3	A	1	Total C O 28 27 1	0
3	A	1	Total C O 28 27 1	0
3	B	1	Total C O 28 27 1	0
3	B	1	Total C O 28 27 1	0
3	B	1	Total C O 28 27 1	0
3	B	1	Total C O 28 27 1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable G-protein coupled receptor 156



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	164823	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	52	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CLR, MW9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/2288	0.37	0/3123
1	B	0.26	0/2288	0.44	2/3123 (0.1%)
All	All	0.24	0/4576	0.40	2/6246 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	251	SER	CA-C-N	-6.73	113.44	120.38
1	B	251	SER	C-N-CA	-6.73	113.44	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2241	0	2299	74	0
1	B	2241	0	2299	73	0
2	A	106	0	0	1	0
2	B	106	0	0	1	0
3	A	140	0	230	34	0
3	B	112	0	184	17	0
All	All	4946	0	5012	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:THR:HG21	3:A:404:CLR:H213	1.46	0.98
1:B:120:ILE:CD1	1:B:216:CYS:SG	2.52	0.97
3:A:405:CLR:H271	1:B:237:LEU:HD21	1.48	0.95
1:A:301:THR:CG2	3:A:404:CLR:H213	1.97	0.94
1:A:297:PHE:HE2	3:A:404:CLR:H241	1.45	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	296/346 (86%)	281 (95%)	15 (5%)	0	100 100
1	B	296/346 (86%)	283 (96%)	13 (4%)	0	100 100
All	All	592/692 (86%)	564 (95%)	28 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/311 (79%)	245 (100%)	0	100	100
1	B	245/311 (79%)	245 (100%)	0	100	100
All	All	490/622 (79%)	490 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265	ASN
1	B	193	GLN
1	B	265	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CLR	A	406	-	31,31,31	0.85	1 (3%)	48,48,48	1.06	5 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CLR	A	405	-	31,31,31	0.79	1 (3%)	48,48,48	1.89	12 (25%)
3	CLR	B	401	-	31,31,31	0.67	1 (3%)	48,48,48	1.81	11 (22%)
2	MW9	B	404	-	52,52,52	1.15	4 (7%)	55,58,58	1.20	3 (5%)
3	CLR	B	406	-	31,31,31	0.67	1 (3%)	48,48,48	1.81	11 (22%)
3	CLR	B	402	-	31,31,31	0.83	0	48,48,48	0.99	2 (4%)
3	CLR	A	404	-	31,31,31	0.83	0	48,48,48	1.04	4 (8%)
3	CLR	B	405	-	31,31,31	0.67	1 (3%)	48,48,48	1.81	11 (22%)
2	MW9	A	402	-	52,52,52	1.15	4 (7%)	55,58,58	1.21	3 (5%)
2	MW9	A	401	-	52,52,52	1.16	4 (7%)	55,58,58	1.19	3 (5%)
2	MW9	B	403	-	52,52,52	1.16	4 (7%)	55,58,58	1.20	3 (5%)
3	CLR	A	403	-	31,31,31	0.86	1 (3%)	48,48,48	1.31	6 (12%)
3	CLR	A	407	-	31,31,31	0.67	1 (3%)	48,48,48	1.81	11 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CLR	A	406	-	-	3/10/68/68	0/4/4/4
3	CLR	A	405	-	-	6/10/68/68	0/4/4/4
3	CLR	B	401	-	-	0/10/68/68	0/4/4/4
2	MW9	B	404	-	-	30/57/57/57	-
3	CLR	B	406	-	-	0/10/68/68	0/4/4/4
3	CLR	B	402	-	-	3/10/68/68	0/4/4/4
3	CLR	A	404	-	-	4/10/68/68	0/4/4/4
3	CLR	B	405	-	-	0/10/68/68	0/4/4/4
2	MW9	A	402	-	-	26/57/57/57	-
2	MW9	A	401	-	-	40/57/57/57	-
2	MW9	B	403	-	-	32/57/57/57	-
3	CLR	A	403	-	-	5/10/68/68	0/4/4/4
3	CLR	A	407	-	-	0/10/68/68	0/4/4/4

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	MW9	C9-C8	-3.19	1.33	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	404	MW9	C9-C8	-3.18	1.33	1.51
2	B	403	MW9	C9-C8	-3.14	1.34	1.51
2	A	401	MW9	C9-C8	-3.13	1.34	1.51
2	B	403	MW9	O1-C17	3.06	1.42	1.33

The worst 5 of 85 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	405	CLR	C13-C17-C20	-5.88	110.28	119.49
3	A	407	CLR	C13-C17-C20	-5.87	110.30	119.49
3	B	406	CLR	C13-C17-C20	-5.85	110.32	119.49
3	B	401	CLR	C13-C17-C20	-5.84	110.35	119.49
3	A	405	CLR	C4-C5-C10	5.01	123.08	116.42

There are no chirality outliers.

5 of 149 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	MW9	O5-C21-C22-C23
2	A	401	MW9	C21-C22-C23-O6
2	A	401	MW9	C25-C24-O8-C19
2	A	401	MW9	C20-O2-P-O4
2	A	401	MW9	C21-O5-P-O3


There are no ring outliers.


11 monomers are involved in 53 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	406	CLR	3	0
3	A	405	CLR	10	0
3	B	401	CLR	6	0
2	B	404	MW9	1	0
3	B	406	CLR	4	0
3	B	402	CLR	1	0
3	A	404	CLR	12	0
3	B	405	CLR	6	0
2	A	402	MW9	1	0
3	A	403	CLR	6	0
3	A	407	CLR	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.