

wwPDB X-ray Structure Validation Summary Report (i)

Nov 14, 2023 – 01:09 AM JST

PDB ID : 5YQT

Title : Crystal Structure of the L74F/M78V/I80V/L114F mutant of LEH complexed

with cyclopentene oxide

Authors: Kong, X.D.; Sun, Z.T.; Wu, L.; Reetz, M.T.; Zhou, J.H.; Xu, J.H.

Deposited on : 2017-11-07

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.30 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	155	80%	%	• 7%
1	В	155		5%	6%
1	С	155	78%	6	• 7%
1	D	155	82%	12%	6%
1	Е	155	77% 14%		10%
1	F	155	77%		9%





Mol	Chain	Length	Quality of chain		
1	G	155	70%	19%	11%
1	Н	155	71%	21%	8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3ZS	С	201[B]	-	-	X	-
2	3ZS	D	201	_	-	X	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Limonene-1,2-epoxide hydrolase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	С	144	Total	С	N	О	S	0	1	0
1		144	1140	732	182	223	3	0	1	U
1	D	145	Total	С	N	О	S	0	1	0
1	ע	149	1144	734	183	224	3	0	1	U
1	A	144	Total	С	N	О	S	0	2	0
1	Λ	144	1145	737	182	223	3	0	_	U
1	В	145	Total	С	N	О	S	0	2	0
1	Ъ	140	1148	738	183	224	3	0	2	U
1	E	140	Total	С	N	О	S	0	1	0
1	Ľ	140	1106	709	175	219	3	0	1	U
1	F	141	Total	С	N	О	S	0	0	0
1	I.	141	1106	707	176	220	3	0		U
1	G	138	Total	С	N	О	S	0	0	0
1	G	130	1083	692	173	215	3	0		U
1	Н	149	Total	С	N	О	S	0	0	0
1	11	142	1113	712	177	221	3		U	U

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	-5	MET	-	expression tag	UNP Q9ZAG3
С	-4	HIS	-	expression tag	UNP Q9ZAG3
С	-3	HIS	-	expression tag	UNP Q9ZAG3
С	-2	HIS	-	expression tag	UNP Q9ZAG3
С	-1	HIS	-	expression tag	UNP Q9ZAG3
С	0	HIS	-	expression tag	UNP Q9ZAG3
С	1	HIS	_	expression tag	UNP Q9ZAG3
С	2	THR	-	expression tag	UNP Q9ZAG3
С	3	SER	-	expression tag	UNP Q9ZAG3
С	4	LEU	-	expression tag	UNP Q9ZAG3
С	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
С	78	VAL	MET	engineered mutation	UNP Q9ZAG3
С	80	VAL	ILE	engineered mutation	UNP Q9ZAG3



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
С	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
D	-5	MET	-	expression tag	UNP Q9ZAG3
D	-4	HIS	-	expression tag	UNP Q9ZAG3
D	-3	HIS	_	expression tag	UNP Q9ZAG3
D	-2	HIS	-	expression tag	UNP Q9ZAG3
D	-1	HIS	-	expression tag	UNP Q9ZAG3
D	0	HIS	-	expression tag	UNP Q9ZAG3
D	1	HIS	-	expression tag	UNP Q9ZAG3
D	2	THR	-	expression tag	UNP Q9ZAG3
D	3	SER	-	expression tag	UNP Q9ZAG3
D	4	LEU	-	expression tag	UNP Q9ZAG3
D	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
D	78	VAL	MET	engineered mutation	UNP Q9ZAG3
D	80	VAL	ILE	engineered mutation	UNP Q9ZAG3
D	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
A	-5	MET	-	expression tag	UNP Q9ZAG3
A	-4	HIS	-	expression tag	UNP Q9ZAG3
A	-3	HIS	-	expression tag	UNP Q9ZAG3
A	-2	HIS	-	expression tag	UNP Q9ZAG3
A	-1	HIS	-	expression tag	UNP Q9ZAG3
A	0	HIS	-	expression tag	UNP Q9ZAG3
A	1	HIS	-	expression tag	UNP Q9ZAG3
A	2	THR	-	expression tag	UNP Q9ZAG3
A	3	SER	-	expression tag	UNP Q9ZAG3
A	4	LEU	-	expression tag	UNP Q9ZAG3
A	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
A	78	VAL	MET	engineered mutation	UNP Q9ZAG3
A	80	VAL	ILE	engineered mutation	UNP Q9ZAG3
A	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
В	-5	MET	-	expression tag	UNP Q9ZAG3
В	-4	HIS	-	expression tag	UNP Q9ZAG3
В	-3	HIS	_	expression tag	UNP Q9ZAG3
В	-2	HIS	-	expression tag	UNP Q9ZAG3
В	-1	HIS	_	expression tag	UNP Q9ZAG3
В	0	HIS	-	expression tag	UNP Q9ZAG3
В	1	HIS	-	expression tag	UNP Q9ZAG3
В	2	THR	-	expression tag	UNP Q9ZAG3
В	3	SER	-	expression tag	UNP Q9ZAG3
В	4	LEU	-	expression tag	UNP Q9ZAG3
В	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
В	78	VAL	MET	engineered mutation	UNP Q9ZAG3
В	80	VAL	ILE	engineered mutation	UNP Q9ZAG3



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
Е	-5	MET	-	expression tag	UNP Q9ZAG3
Е	-4	HIS	-	expression tag	UNP Q9ZAG3
Е	-3	HIS	-	expression tag	UNP Q9ZAG3
Е	-2	HIS	-	expression tag	UNP Q9ZAG3
Е	-1	HIS	-	expression tag	UNP Q9ZAG3
Е	0	HIS	-	expression tag	UNP Q9ZAG3
Е	1	HIS	-	expression tag	UNP Q9ZAG3
Е	2	THR	-	expression tag	UNP Q9ZAG3
Е	3	SER	-	expression tag	UNP Q9ZAG3
Е	4	LEU	-	expression tag	UNP Q9ZAG3
Е	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
Е	78	VAL	MET	engineered mutation	UNP Q9ZAG3
Е	80	VAL	ILE	engineered mutation	UNP Q9ZAG3
Е	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
F	-5	MET	-	expression tag	UNP Q9ZAG3
F	-4	HIS	-	expression tag	UNP Q9ZAG3
F	-3	HIS	-	expression tag	UNP Q9ZAG3
F	-2	HIS	-	expression tag	UNP Q9ZAG3
F	-1	HIS	-	expression tag	UNP Q9ZAG3
F	0	HIS	-	expression tag	UNP Q9ZAG3
F	1	HIS	-	expression tag	UNP Q9ZAG3
F	2	THR	-	expression tag	UNP Q9ZAG3
F	3	SER	-	expression tag	UNP Q9ZAG3
F	4	LEU	-	expression tag	UNP Q9ZAG3
F	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
F	78	VAL	MET	engineered mutation	UNP Q9ZAG3
F	80	VAL	ILE	engineered mutation	UNP Q9ZAG3
F	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
G	-5	MET	-	expression tag	UNP Q9ZAG3
G	-4	HIS	-	expression tag	UNP Q9ZAG3
G	-3	HIS	-	expression tag	UNP Q9ZAG3
G	-2	HIS	-	expression tag	UNP Q9ZAG3
G	-1	HIS	-	expression tag	UNP Q9ZAG3
G	0	HIS	-	expression tag	UNP Q9ZAG3
G	1	HIS	-	expression tag	UNP Q9ZAG3
G	2	THR	-	expression tag	UNP Q9ZAG3
G	3	SER	-	expression tag	UNP Q9ZAG3
G	4	LEU	-	expression tag	UNP Q9ZAG3
G	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
G	78	VAL	MET	engineered mutation	UNP Q9ZAG3
G	80	VAL	ILE	engineered mutation	UNP Q9ZAG3



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	114	PHE	LEU	engineered mutation	UNP Q9ZAG3
Н	-5	MET	-	expression tag	UNP Q9ZAG3
Н	-4	HIS	-	expression tag	UNP Q9ZAG3
Н	-3	HIS	-	expression tag	UNP Q9ZAG3
Н	-2	HIS	-	expression tag	UNP Q9ZAG3
Н	-1	HIS	-	expression tag	UNP Q9ZAG3
Н	0	HIS	-	expression tag	UNP Q9ZAG3
Н	1	HIS	-	expression tag	UNP Q9ZAG3
Н	2	THR	-	expression tag	UNP Q9ZAG3
Н	3	SER	-	expression tag	UNP Q9ZAG3
Н	4	LEU	-	expression tag	UNP Q9ZAG3
Н	74	PHE	LEU	engineered mutation	UNP Q9ZAG3
Н	78	VAL	MET	engineered mutation	UNP Q9ZAG3
Н	80	VAL	ILE	engineered mutation	UNP Q9ZAG3
Н	114	PHE	LEU	engineered mutation	UNP Q9ZAG3

 $\bullet \ \, \text{Molecule 2 is (1R,5S)-6-oxabicyclo[3.1.0]} \\ \text{hexane (three-letter code: 3ZS) (formula: C_5H_8O)}. \\$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total C O 12 10 2	0	1
2	D	1	Total C O 6 5 1	0	0
2	A	1	Total C O 6 5 1	0	0
2	В	1	Total C O 6 5 1	0	0



Mo	l Chain	Residues	Atoms	ZeroOcc	AltConf
2	Н	1	Total C O 6 5 1	0	0

• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	55	Total O 55 55	0	0
3	D	71	Total O 71 71	0	0
3	A	70	Total O 70 70	0	0
3	В	76	Total O 76 76	0	0
3	Ε	36	Total O 36 36	0	0
3	F	42	Total O 42 42	0	0
3	G	30	Total O 30 30	0	0
3	Н	30	Total O 30 30	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Limonene-1,2-epoxide hydrolase



• Molecule 1: Limonene-1,2-epoxide hydrolase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	104.25Å 60.68Å 118.76Å	Denogitor
a, b, c, α , β , γ	90.00° 90.53° 90.00°	Depositor
Resolution (Å)	30.16 - 2.30	Depositor
Resolution (A)	30.15 - 2.30	EDS
% Data completeness	94.8 (30.16-2.30)	Depositor
(in resolution range)	92.9 (30.15-2.30)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.27 (at 2.31Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
D D.	0.224 , 0.283	Depositor
R, R_{free}	0.226 , 0.284	DCC
R_{free} test set	3117 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32 , 27.7	EDS
L-test for twinning ²	$< L > = 0.46, < L^2> = 0.29$	Xtriage
Estimated twinning fraction	0.268 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9431	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 31.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0382e-03.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 3ZS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.50	0/1178	0.62	0/1602
1	В	0.50	0/1181	0.62	0/1606
1	С	0.56	1/1170 (0.1%)	0.64	0/1591
1	D	0.51	0/1174	0.63	0/1596
1	Е	0.37	0/1135	0.58	0/1543
1	F	0.40	0/1131	0.55	0/1538
1	G	0.42	0/1108	0.58	0/1506
1	Н	0.36	0/1139	0.54	0/1550
All	All	0.46	1/9216 (0.0%)	0.60	0/12532

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\textup{\AA})$	$\operatorname{Ideal}(\text{\AA})$
1	С	57	PRO	N-CD	-6.78	1.38	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1145	0	1110	17	0
1	В	1148	0	1111	14	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1140	0	1099	18	0
1	D	1144	0	1102	20	0
1	Ε	1106	0	1057	15	0
1	F	1106	0	1059	18	0
1	G	1083	0	1035	21	0
1	Η	1113	0	1066	22	0
2	A	6	0	8	1	0
2	В	6	0	8	0	0
2	С	12	0	16	5	0
2	D	6	0	8	4	0
2	Η	6	0	8	0	0
3	A	70	0	0	1	0
3	В	76	0	0	0	0
3	С	55	0	0	1	0
3	D	71	0	0	1	0
3	Ε	36	0	0	0	0
3	F	42	0	0	1	0
3	G	30	0	0	0	0
3	Н	30	0	0	1	0
All	All	9431	0	8687	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:42:LYS:NZ	3:A:301:HOH:O	1.61	1.20
1:D:108:THR:HG23	1:D:110:LYS:H	1.30	0.94
1:F:38:ASN:HD21	1:F:80:VAL:H	1.15	0.94
1:G:5:ILE:HG21	1:G:100:VAL:HG21	1.59	0.83
1:G:39:ASP:HB3	1:G:42:LYS:HZ2	1.43	0.81

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	144/155 (93%)	141 (98%)	3 (2%)	0	100	100
1	В	145/155 (94%)	142 (98%)	3 (2%)	0	100	100
1	С	143/155 (92%)	141 (99%)	2 (1%)	0	100	100
1	D	144/155 (93%)	142 (99%)	2 (1%)	0	100	100
1	E	139/155 (90%)	137 (99%)	2 (1%)	0	100	100
1	F	139/155 (90%)	137 (99%)	2 (1%)	0	100	100
1	G	136/155 (88%)	126 (93%)	10 (7%)	0	100	100
1	Н	140/155 (90%)	137 (98%)	3 (2%)	0	100	100
All	All	1130/1240 (91%)	1103 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	122/130 (94%)	119 (98%)	3 (2%)	47	65
1	В	122/130 (94%)	121 (99%)	1 (1%)	81	91
1	С	121/130 (93%)	119 (98%)	2 (2%)	60	76
1	D	121/130 (93%)	121 (100%)	0	100	100
1	E	117/130 (90%)	117 (100%)	0	100	100
1	F	117/130 (90%)	117 (100%)	0	100	100



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	G	114/130 (88%)	113 (99%)	1 (1%)	78	89	
1	Н	118/130 (91%)	116 (98%)	2 (2%)	60	76	
All	All	952/1040 (92%)	943 (99%)	9 (1%)	81	89	

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	15	SER
1	Н	111	SER
1	A	127[A]	ILE
1	A	127[B]	ILE
1	В	13	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	55	ASN
1	D	38	ASN
1	F	38	ASN
1	F	54	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	В	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3ZS	A	201	-	7,7,7	1.37	2 (28%)	6,10,10	2.37	3 (50%)
2	3ZS	Н	201	-	7,7,7	1.49	2 (28%)	6,10,10	2.68	3 (50%)
2	3ZS	D	201	-	7,7,7	1.62	2 (28%)	6,10,10	2.85	3 (50%)
2	3ZS	С	201[A]	-	7,7,7	1.46	2 (28%)	6,10,10	2.61	3 (50%)
2	3ZS	В	201	-	7,7,7	1.44	2 (28%)	6,10,10	2.62	3 (50%)
2	3ZS	С	201[B]	-	7,7,7	1.51	2 (28%)	6,10,10	2.65	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3ZS	A	201	-	=	-	0/2/2/2
2	3ZS	Н	201	-	-	-	0/2/2/2
2	3ZS	D	201	-	=	-	0/2/2/2
2	3ZS	С	201[A]	-	-	-	0/2/2/2
2	3ZS	В	201	-	=	-	0/2/2/2
2	3ZS	С	201[B]	-	-	-	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(\mathring{A})$	Ideal(A)
2	D	201	3ZS	O06-C05	-2.96	1.39	1.45
2	С	201[B]	3ZS	O06-C05	-2.59	1.40	1.45
2	D	201	3ZS	O06-C04	-2.58	1.40	1.45
2	A	201	3ZS	O06-C05	-2.50	1.40	1.45
2	С	201[B]	3ZS	O06-C04	-2.37	1.40	1.45

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
2	D	201	3ZS	C05-O06-C04	5.30	64.51	60.69



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
2	С	201[B]	3ZS	C05-O06-C04	5.00	64.29	60.69
2	Н	201	3ZS	C05-O06-C04	4.95	64.26	60.69
2	С	201[A]	3ZS	C05-O06-C04	4.88	64.21	60.69
2	В	201	3ZS	C05-O06-C04	4.77	64.13	60.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	3ZS	1	0
2	D	201	3ZS	4	0
2	С	201[A]	3ZS	1	0
2	С	201[B]	3ZS	4	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	144/155~(92%)	-0.35	1 (0%) 87 91	28, 35, 51, 70	0
1	В	145/155 (93%)	-0.41	0 100 100	27, 36, 52, 73	0
1	С	144/155 (92%)	-0.36	1 (0%) 87 91	26, 33, 54, 62	0
1	D	145/155 (93%)	-0.31	2 (1%) 75 80	25, 33, 53, 71	1 (0%)
1	Е	140/155 (90%)	-0.22	1 (0%) 87 91	32, 49, 67, 86	0
1	F	141/155 (90%)	-0.20	1 (0%) 87 91	33, 48, 68, 89	0
1	G	138/155 (89%)	-0.02	1 (0%) 87 91	33, 61, 84, 100	2 (1%)
1	Н	142/155 (91%)	-0.13	0 100 100	37, 55, 73, 94	0
All	All	1139/1240 (91%)	-0.25	7 (0%) 89 92	25, 42, 71, 100	3 (0%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	147	LEU	4.7
1	С	147	LEU	3.8
1	Е	145	LEU	3.2
1	G	74	PHE	2.8
1	A	74[A]	PHE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	3ZS	В	201	6/6	0.87	0.46	37,42,47,57	6
2	3ZS	A	201	6/6	0.88	0.29	35,39,46,47	6
2	3ZS	Н	201	6/6	0.91	0.49	57,65,66,70	6
2	3ZS	С	201[A]	6/6	0.93	0.29	30,33,37,37	6
2	3ZS	С	201[B]	6/6	0.93	0.29	30,30,34,35	6
2	3ZS	D	201	6/6	0.93	0.38	24,32,39,40	6

6.5 Other polymers (i)

There are no such residues in this entry.

