



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:19 PM UTC

PDB ID : 8ZAD / pdb\_00008zad  
Title : Crystal structure of RuABA3 from Rutstroemia sp. NJR-2017a WRK4  
Authors : Li, S.Y.; Li, H.; Yang, Y.; Huang, J.-W.; Chen, C.-C.; Guo, R.-T.  
Deposited on : 2024-04-25  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aba 3 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	2806	1792	478	516	20	0	1	0

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	HIS	-	expression tag	UNP A0A2S7P8J3
A	39	HIS	-	expression tag	UNP A0A2S7P8J3
A	40	HIS	-	expression tag	UNP A0A2S7P8J3
A	41	HIS	-	expression tag	UNP A0A2S7P8J3
A	42	HIS	-	expression tag	UNP A0A2S7P8J3
A	43	HIS	-	expression tag	UNP A0A2S7P8J3
A	44	HIS	-	expression tag	UNP A0A2S7P8J3
A	45	HIS	-	expression tag	UNP A0A2S7P8J3
A	46	HIS	-	expression tag	UNP A0A2S7P8J3
A	47	HIS	-	expression tag	UNP A0A2S7P8J3
A	48	GLU	-	expression tag	UNP A0A2S7P8J3
A	49	ASN	-	expression tag	UNP A0A2S7P8J3
A	50	LEU	-	expression tag	UNP A0A2S7P8J3
A	51	TYR	-	expression tag	UNP A0A2S7P8J3
A	52	PHE	-	expression tag	UNP A0A2S7P8J3
A	53	GLN	-	expression tag	UNP A0A2S7P8J3
A	54	GLY	-	expression tag	UNP A0A2S7P8J3
A	55	ALA	-	expression tag	UNP A0A2S7P8J3
A	56	GLY	-	expression tag	UNP A0A2S7P8J3
A	57	ALA	-	expression tag	UNP A0A2S7P8J3
A	58	GLY	-	expression tag	UNP A0A2S7P8J3
A	59	ALA	-	expression tag	UNP A0A2S7P8J3
A	60	GLY	-	expression tag	UNP A0A2S7P8J3
A	61	ALA	-	expression tag	UNP A0A2S7P8J3
A	62	GLY	-	expression tag	UNP A0A2S7P8J3
A	63	ALA	-	expression tag	UNP A0A2S7P8J3
A	64	GLY	-	expression tag	UNP A0A2S7P8J3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	ALA	-	expression tag	UNP A0A2S7P8J3
A	66	MET	-	expression tag	UNP A0A2S7P8J3

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	375	Total	O	0	0
			375	375		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.32Å 141.01Å 90.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.39 – 2.10 36.39 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.39-2.10) 99.9 (36.39-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.77 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.164 , 0.212 0.174 , 0.220	Depositor DCC
$R_{free}$ test set	1308 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.5	Xtrriage
Anisotropy	0.474	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 53.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3182	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.05	0/2874	1.34	2/3889 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	403	PHE	CB-CA-C	6.43	121.37	109.54
1	A	425	GLU	CB-CG-CD	5.89	122.62	112.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2806	0	2707	24	0
2	A	1	0	0	0	0
3	A	375	0	0	5	0
All	All	3182	0	2707	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291[A]:ASN:HD22	1:A:291[A]:ASN:N	1.59	1.01
1:A:182:ASP:OD1	1:A:396:GLY:HA3	1.89	0.72
1:A:251:LYS:O	1:A:255:GLU:HG2	1.92	0.69
1:A:291[A]:ASN:N	1:A:291[A]:ASN:ND2	2.30	0.68
1:A:425:GLU:HG3	3:A:951:HOH:O	1.96	0.64
1:A:256:GLY:HA2	1:A:402:GLN:O	1.98	0.63
1:A:312:MET:HE2	1:A:314:ARG:HG2	1.83	0.60
1:A:97:GLU:OE1	1:A:313:ARG:HD3	2.02	0.60
1:A:284:LEU:HD13	1:A:434:LEU:HD21	1.88	0.55
1:A:292:PRO:O	1:A:295:VAL:HG22	2.08	0.54
1:A:290:ARG:C	1:A:291[A]:ASN:HD22	2.13	0.52
1:A:425:GLU:HG3	3:A:838:HOH:O	2.12	0.49
1:A:425:GLU:CD	1:A:425:GLU:H	2.20	0.48
1:A:96:TRP:CH2	1:A:100:ARG:HD3	2.48	0.48
1:A:71:PRO:HD3	1:A:96:TRP:CD2	2.50	0.47
1:A:385:VAL:CG2	3:A:786:HOH:O	2.64	0.46
1:A:291[A]:ASN:ND2	1:A:291[A]:ASN:H	2.14	0.45
1:A:251:LYS:NZ	3:A:624:HOH:O	2.50	0.45
1:A:128:ARG:NH2	3:A:621:HOH:O	2.49	0.45
1:A:71:PRO:HD3	1:A:96:TRP:CE3	2.54	0.42
1:A:425:GLU:CD	1:A:425:GLU:N	2.77	0.42
1:A:316:ARG:HA	1:A:319:GLU:OE2	2.20	0.42
1:A:205:ASP:C	1:A:205:ASP:OD1	2.64	0.41
1:A:140:ILE:HD12	1:A:145:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	341/405 (84%)	333 (98%)	8 (2%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	291/338 (86%)	274 (94%)	17 (6%)	18 16

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	LEU
1	A	128	ARG
1	A	134	VAL
1	A	255	GLU
1	A	284	LEU
1	A	290	ARG
1	A	291[A]	ASN
1	A	291[B]	ASN
1	A	314	ARG
1	A	316	ARG
1	A	326	LYS
1	A	376	GLU
1	A	393	GLU
1	A	398	GLU
1	A	425	GLU
1	A	430	VAL
1	A	435	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	159	HIS
1	A	221	ASN
1	A	402	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	344/405 (84%)	-0.14	12 (3%) 47 49	12, 20, 48, 73	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	397	ALA	3.7
1	A	396	GLY	3.6
1	A	399	VAL	3.5
1	A	326	LYS	3.2
1	A	291[A]	ASN	3.0
1	A	398	GLU	3.0
1	A	137	SER	2.4
1	A	290	ARG	2.3
1	A	66	MET	2.3
1	A	436	THR	2.2
1	A	254	ALA	2.1
1	A	402	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	500	1/1	1.00	0.03	26,26,26,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.