



# wwPDB NMR Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:29 AM UTC

PDB ID : 1ZGW / pdb\_00001zgw  
Title : NMR structure of E. Coli Ada protein in complex with DNA  
Authors : He, C.; Hus, J.C.; Sun, L.J.; Zhou, P.; Norman, D.P.; Doetsch, V.; Wei, H.;  
Gross, J.D.; Lane, W.S.; Wagner, G.; Verdine, G.L.  
Deposited on : 2005-04-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	18	6% (red) 89% (orange) 6% (red)
2	C	18	22% (yellow) 72% (orange) 6% (red)
3	A	139	16% (green) 55% (yellow) 22% (orange) 6% (cyan)

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:8-A:37, A:39-A:139 (131)	1.04	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 6 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 8, 10, 15, 17, 18, 19
2	2, 4, 5, 11
3	12, 20
4	3, 14
5	7, 9
6	6, 16
Single-model clusters	13

### 3 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3364 atoms, of which 1517 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(\*GP\*CP\*AP\*AP\*AP\*TP\*TP\*AP\*AP\*AP\*GP\*C P\*GP\*CP\*AP\*AP\*GP\*A)-3'.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	B	18	573	177	202	78	99	17	0

- Molecule 2 is a DNA chain called 5'-D(\*TP\*CP\*TP\*TP\*GP\*CP\*GP\*CP\*TP\*TP\*TP\*AP \*AP\*TP\*TP\*TP\*GP\*C)-3'.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	C	18	570	176	209	55	113	17	0

- Molecule 3 is a protein called Ada polyprotein.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
3	A	139	2220	692	1106	219	194	9	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	SMC	CYS	modified residue	UNP P06134

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms
			Total
4	A	1	Zn 1

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(\*GP\*CP\*AP\*AP\*AP\*TP\*TP\*AP\*AP\*AP\*GP\*CP\*GP\*CP\*AP\*AP\*GP\*A)-3'

Chain B: 






- Molecule 2: 5'-D(\*TP\*CP\*TP\*TP\*GP\*CP\*GP\*CP\*TP\*TP\*TP\*AP\*AP\*TP\*TP\*TP\*GP\*C)-3'

Chain C: 



- Molecule 3: Ada polyprotein

Chain A: 



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.


- Molecule 1: 5'-D(\*GP\*CP\*AP\*AP\*AP\*TP\*TP\*AP\*AP\*AP\*GP\*CP\*GP\*CP\*AP\*AP\*GP\*A)-3'

Chain B:  6% 89% 6%




- Molecule 2: 5'-D(\*TP\*CP\*TP\*TP\*GP\*CP\*GP\*CP\*TP\*TP\*TP\*AP\*AP\*TP\*TP\*TP\*GP\*C)-3'

Chain C:  28% 67% 6%



- Molecule 3: Ada polyprotein

Chain A:  17% 57% 17% 6%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 20 were deposited, based on the following criterion: *The submitted conformer models have no restraint violations and lowest energies*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851
Procheck nmr	structure solution	3.5.4

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.54±0.00	18±0/419 ( 4.3± 0.0%)	1.80±0.00	18±0/645 ( 2.8± 0.0%)
2	C	1.55±0.00	18±0/401 ( 4.5± 0.0%)	1.86±0.00	18±0/617 ( 2.9± 0.0%)
3	A	1.40±0.01	0±1/1080 ( 0.0± 0.1%)	1.70±0.01	12±1/1459 ( 0.8± 0.1%)
All	All	1.46	727/38000 ( 1.9%)	1.76	955/54420 ( 1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	1.0±0.0
2	C	0.0±0.0	1.0±0.0
All	All	0	40

5 of 42 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
3	A	77	ALA	N-CA	6.86	1.50	1.46	13	2
3	A	39	ARG	C-N	-6.38	1.27	1.33	13	1
1	B	191	DC	C3'-C2'	-6.23	1.40	1.52	19	20
2	C	233	DA	C3'-C2'	-6.20	1.40	1.52	16	20
1	B	190	DG	C3'-C2'	-6.09	1.40	1.52	12	20

5 of 65 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )	Models	
								Worst	Total
3	A	115	HIS	CA-CB-CG	-10.46	103.34	113.80	5	20
1	B	204	DA	C3'-C2'-C1'	9.52	115.88	101.60	15	20
2	C	235	DT	C3'-C2'-C1'	9.37	115.66	101.60	6	20
1	B	199	DA	C3'-C2'-C1'	9.16	115.34	101.60	7	20
2	C	227	DG	C3'-C2'-C1'	9.14	115.31	101.60	15	20

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	200	DG	Sidechain	20
2	C	227	DG	Sidechain	20

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	371	202	202	28 $\pm$ 4
2	C	361	209	209	37 $\pm$ 6
3	A	1055	1034	1034	230 $\pm$ 16
All	All	35760	28900	28900	5250

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 81.

5 of 1054 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash( $\text{\AA}$ )	Distance( $\text{\AA}$ )	Models	
				Worst	Total
2:C:235:DT:OP1	3:A:46:HIS:O	1.18	1.62	13	5
3:A:97:GLU:OE2	3:A:101:THR:OG1	1.12	1.65	1	3
2:C:224:DT:OP2	3:A:129:LYS:HB3	1.06	1.50	11	19
2:C:234:DT:O3'	3:A:46:HIS:O	1.05	1.75	18	17
3:A:102:LEU:HD23	3:A:116:LEU:HD23	1.02	1.23	1	3

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	130/139 (94%)	113±2 (87±1%)	13±2 (10±1%)	4±1 (3±1%)	5	38
All	All	2600/2780 (94%)	2266 (87%)	256 (10%)	78 (3%)	5	38

5 of 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	98	THR	20
3	A	128	PRO	20
3	A	70	LYS	16
3	A	77	ALA	9
3	A	9	ASP	4

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	108/114 (95%)	81±4 (75±4%)	27±4 (25±4%)	2	24
All	All	2160/2280 (95%)	1613 (75%)	547 (25%)	2	24

5 of 64 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	28	VAL	20
3	A	52	VAL	20
3	A	101	THR	20
3	A	131	TRP	20
3	A	135	TRP	20

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
3	SMC	A	38	3,4	5,6,7	0.65±0.10	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	SMC	A	38	3,4	3,6,8	2.01±0.05	1±0 (33±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SMC	A	38	3,4	-	0±0,3,5,7	-

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	A	38	SMC	CA-CB-SG	3.43	108.50	114.04	13	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided