



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:50 PM UTC

PDB ID : 9ZM4 / pdb\_00009zm4  
Title : Crystal structure of a Heat shock protein (DnaJ) from Brucella melitensis  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2025-12-09  
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	 81% 8% 11%
1	B	172	 76% 8% 15%
1	C	172	 81% 7% 12%
1	D	172	 84% 11% 5%
1	E	172	 83% 8% 10%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	172	 77% 13% 10%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	153	1163	735	210	217	1	0	0	0
1	B	146	1047	658	186	202	1	0	0	0
1	C	152	1171	739	213	218	1	0	0	0
1	D	163	1255	792	229	233	1	0	0	0
1	E	155	1197	756	217	223	1	0	0	0
1	F	155	1181	745	214	221	1	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	140	MET	-	initiating methionine	UNP Q2YMC9
A	141	ALA	-	expression tag	UNP Q2YMC9
A	142	HIS	-	expression tag	UNP Q2YMC9
A	143	HIS	-	expression tag	UNP Q2YMC9
A	144	HIS	-	expression tag	UNP Q2YMC9
A	145	HIS	-	expression tag	UNP Q2YMC9
A	146	HIS	-	expression tag	UNP Q2YMC9
A	147	HIS	-	expression tag	UNP Q2YMC9
B	140	MET	-	initiating methionine	UNP Q2YMC9
B	141	ALA	-	expression tag	UNP Q2YMC9
B	142	HIS	-	expression tag	UNP Q2YMC9
B	143	HIS	-	expression tag	UNP Q2YMC9
B	144	HIS	-	expression tag	UNP Q2YMC9
B	145	HIS	-	expression tag	UNP Q2YMC9
B	146	HIS	-	expression tag	UNP Q2YMC9
B	147	HIS	-	expression tag	UNP Q2YMC9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	140	MET	-	initiating methionine	UNP Q2YMC9
C	141	ALA	-	expression tag	UNP Q2YMC9
C	142	HIS	-	expression tag	UNP Q2YMC9
C	143	HIS	-	expression tag	UNP Q2YMC9
C	144	HIS	-	expression tag	UNP Q2YMC9
C	145	HIS	-	expression tag	UNP Q2YMC9
C	146	HIS	-	expression tag	UNP Q2YMC9
C	147	HIS	-	expression tag	UNP Q2YMC9
D	140	MET	-	initiating methionine	UNP Q2YMC9
D	141	ALA	-	expression tag	UNP Q2YMC9
D	142	HIS	-	expression tag	UNP Q2YMC9
D	143	HIS	-	expression tag	UNP Q2YMC9
D	144	HIS	-	expression tag	UNP Q2YMC9
D	145	HIS	-	expression tag	UNP Q2YMC9
D	146	HIS	-	expression tag	UNP Q2YMC9
D	147	HIS	-	expression tag	UNP Q2YMC9
E	140	MET	-	initiating methionine	UNP Q2YMC9
E	141	ALA	-	expression tag	UNP Q2YMC9
E	142	HIS	-	expression tag	UNP Q2YMC9
E	143	HIS	-	expression tag	UNP Q2YMC9
E	144	HIS	-	expression tag	UNP Q2YMC9
E	145	HIS	-	expression tag	UNP Q2YMC9
E	146	HIS	-	expression tag	UNP Q2YMC9
E	147	HIS	-	expression tag	UNP Q2YMC9
F	140	MET	-	initiating methionine	UNP Q2YMC9
F	141	ALA	-	expression tag	UNP Q2YMC9
F	142	HIS	-	expression tag	UNP Q2YMC9
F	143	HIS	-	expression tag	UNP Q2YMC9
F	144	HIS	-	expression tag	UNP Q2YMC9
F	145	HIS	-	expression tag	UNP Q2YMC9
F	146	HIS	-	expression tag	UNP Q2YMC9
F	147	HIS	-	expression tag	UNP Q2YMC9


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain A: 





- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain B: 




- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain C: 





- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain D: 





- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain E:  83% 8% 10%



- Molecule 1: Heat shock protein DnaJ, N-terminal:Chaperone DnaJ, C-terminal:Heat shock protein DnaJ

Chain F:  77% 13% 10%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.48Å 94.29Å 150.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.20 – 3.09 46.20 – 3.09	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.20-3.09) 99.9 (46.20-3.09)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 3.06Å)	Xtrriage
Refinement program	PHENIX (2.0_5896: ???)	Depositor
R, $R_{free}$	0.241 , 0.293 0.240 , 0.289	Depositor DCC
$R_{free}$ test set	955 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	96.1	Xtrriage
Anisotropy	0.756	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 90.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7015	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	133.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/1182	0.39	0/1591
1	B	0.17	0/1059	0.36	0/1432
1	C	0.17	0/1188	0.34	0/1595
1	D	0.18	0/1276	0.37	0/1716
1	E	0.21	0/1215	0.39	0/1630
1	F	0.20	0/1199	0.39	0/1614
All	All	0.20	0/7119	0.37	0/9578

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1163	0	1177	10	0
1	B	1047	0	983	10	0
1	C	1171	0	1199	8	0
1	D	1255	0	1284	14	0
1	E	1197	0	1237	10	0
1	F	1181	0	1199	18	0
2	A	1	0	0	0	0
All	All	7015	0	7079	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:307:LEU:HD13	1:F:239:ASP:HA	1.60	0.84
1:A:234:LEU:HD13	1:A:249:VAL:HG22	1.63	0.79
1:E:239:ASP:HA	1:F:307:LEU:HD13	1.68	0.75
1:A:294:MET:HE1	1:B:267:ASP:HB3	1.71	0.73
1:C:267:ASP:HB3	1:D:294:MET:HE1	1.75	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/172 (87%)	145 (97%)	4 (3%)	0	100	100
1	B	140/172 (81%)	135 (96%)	4 (3%)	1 (1%)	18	49
1	C	148/172 (86%)	145 (98%)	3 (2%)	0	100	100
1	D	159/172 (92%)	156 (98%)	3 (2%)	0	100	100
1	E	151/172 (88%)	148 (98%)	3 (2%)	0	100	100
1	F	151/172 (88%)	146 (97%)	5 (3%)	0	100	100
All	All	898/1032 (87%)	875 (97%)	22 (2%)	1 (0%)	48	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	297	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/143 (86%)	123 (100%)	0	100	100
1	B	99/143 (69%)	99 (100%)	0	100	100
1	C	125/143 (87%)	125 (100%)	0	100	100
1	D	135/143 (94%)	135 (100%)	0	100	100
1	E	129/143 (90%)	129 (100%)	0	100	100
1	F	125/143 (87%)	125 (100%)	0	100	100
All	All	736/858 (86%)	736 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	GLN
1	E	163	GLN
1	E	192	GLN
1	F	179	HIS
1	F	199	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/172 (88%)	-0.55	0 <a href="#">100</a> <a href="#">100</a>	74, 104, 175, 215	0
1	B	146/172 (84%)	-0.53	1 (0%) <a href="#">84</a> <a href="#">68</a>	115, 163, 190, 215	0
1	C	152/172 (88%)	-0.53	0 <a href="#">100</a> <a href="#">100</a>	104, 139, 187, 211	0
1	D	163/172 (94%)	-0.57	0 <a href="#">100</a> <a href="#">100</a>	93, 131, 178, 201	0
1	E	155/172 (90%)	-0.50	0 <a href="#">100</a> <a href="#">100</a>	88, 119, 163, 182	0
1	F	155/172 (90%)	-0.49	0 <a href="#">100</a> <a href="#">100</a>	77, 127, 179, 196	0
All	All	924/1032 (89%)	-0.53	1 (0%) <a href="#">92</a> <a href="#">86</a>	74, 132, 184, 215	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	238	ILE	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	401	1/1	0.13	0.20	217,217,217,217	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.