

wwPDB X-ray Structure Validation Summary Report (i)

Apr 9, 2025 – 06:16 PM JST

PDB ID	:	8 ZN6 / pdb_00008zn6
Title	:	Crystal Structure of Designed Clock Protein KaiC
Authors	:	Furuike, Y.; Akiyama, S.
Deposited on	:	2024-05-26
Resolution	:	2.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	164625	$1067 \ (2.96-2.92)$
Clashscore	180529	1122 (2.96-2.92)
Ramachandran outliers	177936	1075 (2.96-2.92)
Sidechain outliers	177891	1075 (2.96-2.92)
RSRZ outliers	164620	1067 (2.96-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	515	2% 85%	6% • 8%
-		010	% •	0,0 - 0,0
1	F	515	80%	10% • 9%
1	J	515	82%	8% 10%
1	K	515	2% 82%	7% 11%
2	В	515	% 82%	10% 8%
2	С	515	3% 82%	9% • 9%

Continued on next page...



Conti	nueu jron	i previous	page	
Mol	Chain	Length	Quality of chain	
2	D	515	2% 81%	7% • 10%
2	Е	515	% 8 3%	10% 8%
2	G	515	2% 83%	7% • 10%
2	Н	515	% 8 2%	9% • 8%
2	Ι	515	% 	8% • 10%
2	L	515	% 8 2%	9% • 9%

Continued from previous page



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 41205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	F	460	Total	С	Ν	0	Р	\mathbf{S}	0	0	0
1	I.	409	3437	2172	597	657	1	10	0	0	0
1	Δ	475	Total	С	Ν	0	Р	S	0	0	0
1	Л	475	3477	2205	603	658	1	10			0
1	K	460	Total	С	Ν	0	Р	S	0	0	0
1	К	400	3241	2036	566	629	1	9			0
1	1 T	466	Total	С	Ν	0	Р	S	0	0	0
I J	400	3316	2090	578	637	1	10	0	0	0	

• Molecule 1 is a protein called KaiC.

• Molecule 2 is a protein called KaiC.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	л	464	Total	С	Ν	0	\mathbf{S}	0	0	0
	D	404	3304	2095	574	626	9	0	0	0
0	F	476	Total	С	Ν	0	S	0	0	0
	Ľ	470	3446	2180	601	655	10	0		0
0	В	479	Total	С	Ν	0	S	0	0	0
	D	412	3396	2154	586	646	10		0	0
0	C	471	Total	С	Ν	0	S	0	0	0
	U	471	3350	2115	582	641	12			
0	С	466	Total	С	Ν	0	S	0	0	0
	G	400	3376	2144	588	634	10			
2	Ц	473	Total	С	Ν	0	S	0	0	0
	11	475	3321	2110	584	618	9	0	0	0
0	т	471	Total	С	Ν	0	S	0	0	0
		471	3427	2172	597	647	11	0	U	0
2	т	466	Total	С	Ν	0	S	0	0	0
	1	400	3296	2091	574	621	10	U		0

• Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
0	F	1	Total	С	Ν	Ο	Р	0	0
3	F	1	31	10	6	12	3	0	0
9	Б	1	Total	С	Ν	Ο	Р	0	0
່ <u>ບ</u>	Г	1	31	10	6	12	3	0	0
3	Λ	1	Total	С	Ν	Ο	Р	0	0
່ <u>ບ</u>	Л	1	31	10	6	12	3	0	0
3	Λ	1	Total	С	Ν	Ο	Р	0	0
5	Л	1	31	10	6	12	3	0	0
3	л	1	Total	С	Ν	Ο	Р	0	0
5	D	I	31	10	6	12	3	0	0
3	л	1	Total	С	Ν	Ο	Р	0	0
5	D	I	31	10	6	12	3	0	0
3	F	1	Total	С	Ν	Ο	Р	0	0
0	Ľ	1	31	10	6	12	3		
3	E	1	Total	С	Ν	Ο	Р	0	0
5	Ľ	I	31	10	6	12	3	0	0
3	В	1	Total	С	Ν	Ο	Р	0	0
5	D	1	31	10	6	12	3	0	0
2	В	1	Total	С	Ν	Ο	Р	0	0
5	D	1	31	10	6	12	3	0	0
2	С	1	Total	С	Ν	Ο	Р	0	0
5	U	1	31	10	6	12	3	0	0
3	C	1	Total	С	Ν	Ο	Р	0	0
5	U	1	31	10	6	12	3	U	0
3	G	1	Total	$\overline{\mathbf{C}}$	Ν	Ο	Р	0	0
	G	L	31	10	6	12	3	0	U
3	G	1	Total	$\overline{\mathbf{C}}$	Ν	Ο	Р	0	0
0	G	L	31	10	6	12	3	U	0

Continued on next page...



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	Ц	1	Total	С	Ν	Ο	Р	0	0
0	5 11		31	10	6	12	3	0	0
9	ц	1	Total	С	Ν	Ο	Р	0	0
0	п	1	31	10	6	12	3	0	0
2	K	1	Total	С	Ν	Ο	Р	0	0
5	Γ	1	31	10	6	12	3	0	0
3	K	1	Total	С	Ν	Ο	Р	0	0
5	Γ	1	31	10	6	12	3	0	0
3	т	1	Total	С	Ν	Ο	Р	0	0
5	L	T	31	10	6	12	3		
3	T	1	Total	С	Ν	Ο	Р	0	0
5	Ľ	T	31	10	6	12	3		0
3	T	1	Total	С	Ν	Ο	Р	0	0
5	1	L	31	10	6	12	3	0	0
3	T	1	Total	С	Ν	Ο	Р	0	0
5	I	I	31	10	6	12	3	0	0
3	T	1	Total	С	Ν	Ο	Р	0	0
5	J J	1	31	10	6	12	3	0	0
3	Т	1	Total	С	Ν	Ο	Р	0	0
5	J	1	31	10	6	12	3	0	0

Continued from previous page...

• Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	F	2	Total Mg 2 2	0	0
4	А	2	Total Mg 2 2	0	0
4	D	1	Total Mg 1 1	0	0
4	Е	2	Total Mg 2 2	0	0
4	В	1	Total Mg 1 1	0	0
4	С	1	Total Mg 1 1	0	0
4	G	2	Total Mg 2 2	0	0
4	Н	2	Total Mg 2 2	0	0
4	K	2	Total Mg 2 2	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	2	Total Mg 2 2	0	0
4	Ι	2	Total Mg 2 2	0	0
4	J	1	Total Mg 1 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	F	5	Total O 5 5	0	0
5	А	5	Total O 5 5	0	0
5	D	5	Total O 5 5	0	0
5	Ε	6	Total O 6 6	0	0
5	В	3	Total O 3 3	0	0
5	С	4	Total O 4 4	0	0
5	G	4	Total O 4 4	0	0
5	Н	8	Total O 8 8	0	0
5	К	2	Total O 2 2	0	0
5	L	5	Total O 5 5	0	0
5	Ι	4	Total O 4 4	0	0
5	J	3	Total O 3 3	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: KaiC









GLY GLN GLU VAL VAL VAL GLY ASN PHE ASP MET THR ASP PRO PRO ASN ASN ASN GLU GLU GLU GLU ALA ALA T496 VAL ASP CLU CLV CLV CLV SER SER ARG CLV VAL CLV CLV GLN GLN GLV GLV • Molecule 2: KaiC 3% Chain C: 82% 9% 9% MET THR ASP PRO PRO PRO ASN ASN GLU GLU GLU GLU GLU PRO GLV GLV GLV VAL VAL VAL VAL VAL SNN • Molecule 2: KaiC Chain G: 83% 7% • 10% MET THR ASP PPRO PPRO PPRO ASN ASN ASN ASN HIS GLU GLU GLU THR GLY GLN GLU VAL VAL VAL VAL VAL CLY GLY GLY • Molecule 2: KaiC Chain H: 82% 9% 8% MET THR ASP PRO ASN ASN ASN GLU GLU HIS GLU GLU LLEU GLY ALA MET ARG







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	90.76Å 110.27Å 166.89Å	Depositor
a, b, c, α , β , γ	77.97° 87.25° 82.41°	Depositor
Bosolution (Å)	48.74 - 2.94	Depositor
	48.74 - 2.94	EDS
% Data completeness	99.5 (48.74-2.94)	Depositor
(in resolution range)	99.5(48.74-2.94)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.49 (at 2.96 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
R R.	0.239 , 0.292	Depositor
II, II, <i>free</i>	0.237 , 0.288	DCC
R_{free} test set	6615 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	79.2	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , 75.3	EDS
L-test for $twinning^2$	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	41205	wwPDB-VP
Average B, all atoms $(Å^2)$	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, SEP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond	angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.27	0/3527	0.44	0/4784
1	F	0.26	0/3484	0.43	0/4727
1	J	0.27	0/3363	0.44	0/4570
1	Κ	0.27	0/3287	0.43	0/4467
2	В	0.26	0/3454	0.45	0/4697
2	С	0.27	0/3407	0.44	0/4633
2	D	0.27	0/3362	0.44	0/4571
2	Е	0.26	0/3506	0.44	0/4767
2	G	0.26	0/3433	0.44	0/4664
2	Н	0.26	0/3379	0.44	0/4604
2	Ι	0.27	0/3351	0.44	0/4555
2	L	0.27	0/3486	0.44	0/4727
All	All	0.27	0/41039	0.44	0/55766

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	С	0	1
2	D	0	4
All	All	0	5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
2	С	213	ARG	Sidechain
2	D	164	ARG	Sidechain
2	D	213	ARG	Sidechain
2	D	486	ARG	Sidechain
2	D	494	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3477	0	3234	13	0
1	F	3437	0	3225	23	0
1	J	3316	0	2972	14	0
1	K	3241	0	2847	15	0
2	В	3396	0	3162	18	0
2	С	3350	0	3030	21	0
2	D	3304	0	2996	20	0
2	Е	3446	0	3184	19	0
2	G	3376	0	3154	10	0
2	Н	3321	0	3003	24	0
2	Ι	3296	0	3006	19	0
2	L	3427	0	3217	19	0
3	А	62	0	26	1	0
3	В	62	0	26	2	0
3	С	62	0	26	1	0
3	D	62	0	26	2	0
3	Е	62	0	26	0	0
3	F	62	0	26	0	0
3	G	62	0	26	0	0
3	Н	62	0	26	0	0
3	Ι	62	0	26	1	0
3	J	62	0	26	2	0
3	Κ	62	0	26	2	0
3	L	62	0	26	1	0
4	A	2	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	Е	2	0	0	0	0

Continued on next page...



	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	2	0	0	0	0
4	G	2	0	0	0	0
4	Н	2	0	0	0	0
4	Ι	2	0	0	0	0
4	J	1	0	0	0	0
4	K	2	0	0	0	0
4	L	2	0	0	0	0
5	А	5	0	0	0	0
5	В	3	0	0	0	0
5	С	4	0	0	0	0
5	D	5	0	0	0	0
5	Е	6	0	0	0	0
5	F	5	0	0	0	0
5	G	4	0	0	0	0
5	Н	8	0	0	0	0
5	Ι	4	0	0	0	0
5	J	3	0	0	0	0
5	Κ	2	0	0	0	0
5	L	5	0	0	0	0
All	All	41205	0	37342	199	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:213:ARG:HG2	2:C:213:ARG:HH21	1.29	0.97
2:B:398:THR:HG21	2:B:431:ILE:HG23	1.65	0.79
2:C:398:THR:HG21	2:C:431:ILE:HG23	1.66	0.77
2:B:183:ILE:HD11	2:C:188:PRO:HA	1.66	0.76
2:D:213:ARG:HH11	2:D:213:ARG:HB2	1.57	0.68

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	466/515~(90%)	444 (95%)	21 (4%)	1 (0%)	44	66
1	F	464/515~(90%)	437 (94%)	23~(5%)	4 (1%)	14	35
1	J	459/515~(89%)	436 (95%)	22 (5%)	1 (0%)	44	66
1	Κ	451/515 (88%)	434 (96%)	15 (3%)	2(0%)	30	55
2	В	468/515~(91%)	441 (94%)	23~(5%)	4 (1%)	14	35
2	С	465/515~(90%)	429 (92%)	33 (7%)	3 (1%)	22	46
2	D	454/515~(88%)	426 (94%)	28 (6%)	0	100	100
2	Ε	472/515~(92%)	440 (93%)	27 (6%)	5 (1%)	12	30
2	G	458/515~(89%)	438 (96%)	17 (4%)	3 (1%)	19	41
2	Н	467/515~(91%)	435 (93%)	28 (6%)	4 (1%)	14	35
2	Ι	460/515~(89%)	436 (95%)	21 (5%)	3 (1%)	19	41
2	L	465/515~(90%)	433 (93%)	31 (7%)	1 (0%)	44	66
All	All	5549/6180 (90%)	5229 (94%)	289 (5%)	31 (1%)	22	46

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	Ε	153	ASP
2	В	155	ALA
2	С	108	PRO
1	F	155	ALA
1	А	194	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	А	332/442~(75%)	316~(95%)	16 (5%)	21	44
1	F	334/442~(76%)	315 (94%)	19 (6%)	17	37
1	J	300/442~(68%)	280 (93%)	20 (7%)	13	31
1	К	287/442~(65%)	270 (94%)	17 (6%)	16	36
2	В	325/443~(73%)	305 (94%)	20 (6%)	15	34
2	С	309/443~(70%)	287 (93%)	22 (7%)	12	29
2	D	304/443~(69%)	282 (93%)	22 (7%)	12	29
2	Е	326/443~(74%)	310~(95%)	16 (5%)	21	43
2	G	323/443 (73%)	299~(93%)	24 (7%)	11	27
2	Н	296/443~(67%)	283~(96%)	13 (4%)	24	48
2	Ι	300/443~(68%)	284 (95%)	16 (5%)	19	40
2	L	330/443 (74%)	308 (93%)	22 (7%)	13	31
All	All	3766/5312 (71%)	3539 (94%)	227 (6%)	16	35

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

5 of 227 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
2	G	146	THR
1	J	404	GLU
2	Н	215	ARG
1	J	354	LEU
2	Ι	268	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such side chains are listed below:

Mol	Chain	Res	Type
2	G	412	ASN
1	J	412	ASN
2	Н	207	ASN
2	Ι	207	ASN
2	Н	31	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Turne	Chain	Dec	Tink	Bond lengths			Bond angles		
MOI	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
1	SEP	F	429	1	8,9,10	0.62	0	8,12,14	0.63	0
1	SEP	J	429	1	8,9,10	0.59	0	8,12,14	0.67	0
1	SEP	А	429	1	8,9,10	0.61	0	8,12,14	0.63	0
1	SEP	K	429	1	8,9,10	0.60	0	8,12,14	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	F	429	1	-	5/5/8/10	-
1	SEP	J	429	1	-	4/5/8/10	-
1	SEP	А	429	1	-	3/5/8/10	-
1	SEP	K	429	1	-	1/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	F	429	SEP	N-CA-CB-OG
1	F	429	SEP	CB-OG-P-O1P
1	F	429	SEP	CB-OG-P-O2P
1	F	429	SEP	CB-OG-P-O3P
1	А	429	SEP	CB-OG-P-O2P

There are no ring outliers.



No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 44 ligands modelled in this entry, 20 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dec	Tiple	Bo	ond leng	ths	B	ond ang	gles
IVIOI	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	E	601	4	29,33,33	1.20	5 (17%)	31,52,52	1.11	2 (6%)
3	ANP	D	601	-	29,33,33	1.19	5 (17%)	31,52,52	1.11	2 (6%)
3	ANP	А	602	4	29,33,33	1.19	<mark>5 (17%)</mark>	31,52,52	1.09	2 (6%)
3	ANP	Ι	602	4	29,33,33	1.20	<mark>5 (17%)</mark>	31,52,52	1.13	2 (6%)
3	ANP	G	601	4	29,33,33	1.19	<mark>5 (17%)</mark>	31,52,52	1.16	3 (9%)
3	ANP	Н	601	4	29,33,33	1.19	<mark>5 (17%)</mark>	31,52,52	1.11	3 (9%)
3	ANP	F	602	4	29,33,33	1.20	<mark>5 (17%)</mark>	31,52,52	1.19	2 (6%)
3	ANP	J	601	-	29,33,33	1.20	5 (17%)	31,52,52	1.09	2 (6%)
3	ANP	Н	602	4	29,33,33	1.21	5 (17%)	31,52,52	1.14	2 (6%)
3	ANP	L	602	4	29,33,33	1.19	5 (17%)	31,52,52	1.19	3 (9%)
3	ANP	J	602	4	29,33,33	1.22	<mark>5 (17%)</mark>	31,52,52	1.13	3 (9%)
3	ANP	В	601	-	29,33,33	1.19	<mark>5 (17%)</mark>	31,52,52	1.09	2 (6%)
3	ANP	K	601	4	29,33,33	1.20	<mark>5 (17%)</mark>	31,52,52	1.15	3 (9%)
3	ANP	F	601	4	29,33,33	1.22	<mark>5 (17%)</mark>	31,52,52	1.09	2 (6%)
3	ANP	K	602	4	29,33,33	1.22	5 (17%)	31,52,52	1.13	3 (9%)
3	ANP	G	602	4	29,33,33	1.20	<mark>5 (17%)</mark>	31,52,52	1.13	3 (9%)
3	ANP	Е	602	4	29,33,33	1.21	<mark>5 (17%)</mark>	31,52,52	1.12	2 (6%)
3	ANP	L	601	4	29,33,33	1.20	<mark>5 (17%)</mark>	31,52,52	1.10	2 (6%)
3	ANP	А	601	4	29,33,33	1.19	5 (17%)	31,52,52	1.15	2(6%)
3	ANP	В	602	4	29,33,33	1.19	5 (17%)	31,52,52	1.15	2 (6%)



Mal	Aol Type Chain	Dec	Tinle	Bo	Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	ANP	С	602	4	29,33,33	1.19	5 (17%)	31,52,52	1.08	2 (6%)
3	ANP	Ι	601	4	29,33,33	1.20	5 (17%)	31,52,52	1.11	2 (6%)
3	ANP	D	602	4	29,33,33	1.18	5 (17%)	31,52,52	1.14	2 (6%)
3	ANP	С	601	-	29,33,33	1.20	5 (17%)	31,52,52	1.14	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	Е	601	4	-	8/14/38/38	0/3/3/3
3	ANP	D	601	-	-	5/14/38/38	0/3/3/3
3	ANP	А	602	4	-	5/14/38/38	0/3/3/3
3	ANP	Ι	602	4	-	3/14/38/38	0/3/3/3
3	ANP	G	601	4	-	9/14/38/38	0/3/3/3
3	ANP	Н	601	4	-	7/14/38/38	0/3/3/3
3	ANP	F	602	4	-	3/14/38/38	0/3/3/3
3	ANP	J	601	-	-	7/14/38/38	0/3/3/3
3	ANP	Н	602	4	-	6/14/38/38	0/3/3/3
3	ANP	L	602	4	-	3/14/38/38	0/3/3/3
3	ANP	J	602	4	-	6/14/38/38	0/3/3/3
3	ANP	В	601	-	-	5/14/38/38	0/3/3/3
3	ANP	K	601	4	-	4/14/38/38	0/3/3/3
3	ANP	F	601	4	-	4/14/38/38	0/3/3/3
3	ANP	K	602	4	-	6/14/38/38	0/3/3/3
3	ANP	G	602	4	-	6/14/38/38	0/3/3/3
3	ANP	Е	602	4	-	3/14/38/38	0/3/3/3
3	ANP	L	601	4	-	8/14/38/38	0/3/3/3
3	ANP	А	601	4	-	3/14/38/38	0/3/3/3
3	ANP	В	602	4	-	6/14/38/38	0/3/3/3
3	ANP	С	602	4	-	1/14/38/38	0/3/3/3
3	ANP	Ι	601	4	-	5/14/38/38	0/3/3/3
3	ANP	D	602	4	-	5/14/38/38	0/3/3/3
3	ANP	С	601	-	-	2/14/38/38	0/3/3/3



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	601	ANP	PG-01G	3.30	1.51	1.46
3	L	601	ANP	PG-01G	3.22	1.51	1.46
3	J	601	ANP	PG-01G	3.21	1.51	1.46
3	J	602	ANP	PG-01G	3.21	1.51	1.46
3	G	601	ANP	PG-01G	3.18	1.51	1.46

The worst 5 of 120 bond length outliers are listed below:

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	В	602	ANP	O2B-PB-O1B	4.28	118.90	109.92
3	J	601	ANP	O2B-PB-O1B	4.23	118.79	109.92
3	G	601	ANP	O2B-PB-O1B	4.23	118.78	109.92
3	L	602	ANP	O2B-PB-O1B	4.22	118.77	109.92
3	Ι	602	ANP	O2B-PB-O1B	4.21	118.75	109.92

There are no chirality outliers.

5 of 120 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
3	F	601	ANP	PB-N3B-PG-O1G
3	F	601	ANP	PG-N3B-PB-O1B
3	F	601	ANP	PA-O3A-PB-O1B
3	F	601	ANP	PA-O3A-PB-O2B
3	F	602	ANP	PB-N3B-PG-O1G

There are no ring outliers.

11 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Ι	602	ANP	1	0
3	J	601	ANP	1	0
3	J	602	ANP	1	0
3	В	601	ANP	1	0
3	Κ	601	ANP	1	0
3	Κ	602	ANP	1	0
3	L	601	ANP	1	0
3	А	601	ANP	1	0
3	В	602	ANP	1	0
3	С	602	ANP	1	0
3	D	602	ANP	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















8ZN6































Torsions



Rings





Torsions

Rings























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ :	>2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	А	474/515~(92%)	0.31	12 (2%) 58	55	51, 74, 104, 127	0
1	F	468/515~(90%)	0.18	6 (1%) 74	72	52, 74, 108, 125	0
1	J	465/515~(90%)	0.26	9 (1%) 66	64	57, 77, 115, 140	0
1	Κ	459/515~(89%)	0.26	9 (1%) 64	62	52, 79, 107, 135	0
2	В	472/515~(91%)	0.21	5 (1%) 77	75	49, 74, 106, 144	0
2	С	471/515~(91%)	0.32	13 (2%) 55	51	57, 82, 111, 134	0
2	D	464/515~(90%)	0.28	8 (1%) 69	66	57, 81, 107, 135	0
2	Ε	476/515~(92%)	0.17	7 (1%) 71	69	51, 70, 102, 128	0
2	G	466/515~(90%)	0.20	10 (2%) 63	61	55, 76, 99, 124	0
2	Н	473/515~(91%)	0.26	6 (1%) 74	72	55, 80, 104, 132	0
2	Ι	466/515~(90%)	0.29	7 (1%) 71	69	57, 78, 112, 132	0
2	L	471/515~(91%)	0.19	5 (1%) 77	75	55, 75, 100, 125	0
All	All	5625/6180~(91%)	0.24	97 (1%) 69	66	49, 77, 108, 144	0

The worst 5 of 97 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	423	ILE	3.9
2	D	153	ASP	3.8
1	А	154	ALA	3.5
2	С	148	VAL	3.3
2	Ι	255	ARG	3.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q < 0.9
1	SEP	F	429	10/11	0.76	0.21	90,101,120,121	0
1	SEP	А	429	10/11	0.79	0.19	95,106,121,121	0
1	SEP	J	429	10/11	0.79	0.17	90,101,117,119	0
1	SEP	Κ	429	10/11	0.85	0.14	83,98,115,117	0

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	MG	Е	604	1/1	0.62	0.15	90,90,90,90	0
4	MG	G	604	1/1	0.74	0.17	84,84,84,84	0
4	MG	K	604	1/1	0.74	0.09	82,82,82,82	0
4	MG	В	603	1/1	0.77	0.20	71,71,71,71	0
4	MG	А	604	1/1	0.85	0.14	69,69,69,69	0
4	MG	Ι	604	1/1	0.85	0.12	80,80,80,80	0
4	MG	L	604	1/1	0.87	0.10	73,73,73,73	0
4	MG	F	604	1/1	0.88	0.10	76,76,76,76	0
3	ANP	J	601	31/31	0.90	0.10	76,81,109,109	0
3	ANP	K	601	31/31	0.90	0.10	73,76,96,96	0
3	ANP	Ι	601	31/31	0.90	0.10	81,84,94,96	0
3	ANP	D	601	31/31	0.91	0.10	64,65,91,95	0
4	MG	K	603	1/1	0.91	0.12	58,58,58,58	0
3	ANP	L	601	31/31	0.91	0.10	64,66,89,92	0
3	ANP	Е	601	31/31	0.91	0.09	71,74,92,94	0
3	ANP	Н	601	31/31	0.91	0.09	69,76,89,92	0
3	ANP	G	601	31/31	0.92	0.09	$61,\!64,\!94,\!97$	0
4	MG	Н	604	1/1	0.92	0.06	74,74,74,74	0
3	ANP	F	601	31/31	0.92	0.09	71,72,91,95	0
3	ANP	A	602	31/31	0.92	0.10	62,66,84,86	0
3	ANP	В	601	31/31	0.92	0.09	68,71,92,94	0
3	ANP	С	601	31/31	0.92	0.09	71,77,90,91	0

Continued on next page...



8ZN6	
------	--

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
3	ANP	G	602	31/31	0.93	0.08	67, 76, 81, 82	0
3	ANP	Ι	602	31/31	0.94	0.07	61,63,74,75	0
3	ANP	F	602	31/31	0.94	0.08	58,61,66,68	0
3	ANP	K	602	31/31	0.94	0.07	62,66,70,71	0
3	ANP	С	602	31/31	0.95	0.07	$65,\!69,\!78,\!79$	0
3	ANP	Н	602	31/31	0.95	0.07	70,73,77,78	0
3	ANP	В	602	31/31	0.95	0.07	55,58,60,61	0
3	ANP	D	602	31/31	0.95	0.07	65,66,74,76	0
3	ANP	J	602	31/31	0.95	0.07	$63,\!67,\!72,\!73$	0
3	ANP	L	602	31/31	0.96	0.06	56,58,64,65	0
3	ANP	Е	602	31/31	0.96	0.07	63,67,70,71	0
3	ANP	А	601	31/31	0.96	0.06	54,61,62,64	0
4	MG	Ι	603	1/1	0.97	0.04	58, 58, 58, 58	0
4	MG	D	603	1/1	0.97	0.06	$57,\!57,\!57,\!57$	0
4	MG	G	603	1/1	0.98	0.05	$54,\!54,\!54,\!54$	0
4	MG	Н	603	1/1	0.99	0.03	76, 76, 76, 76, 76	0
4	MG	J	603	1/1	0.99	0.02	$65,\!65,\!65,\!65$	0
4	MG	L	603	1/1	1.00	0.03	$55,\!55,\!55,\!55$	0
4	MG	А	603	1/1	1.00	0.03	59, 59, 59, 59, 59	0
4	MG	С	603	1/1	1.00	0.03	71,71,71,71	0
4	MG	Е	603	1/1	1.00	0.04	$47,\!47,\!47,\!47$	0
4	MG	F	603	1/1	1.00	0.01	49,49,49,49	0

Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















































6.5 Other polymers (i)

There are no such residues in this entry.

