



wwPDB EM Validation Summary Report ⓘ

Jul 16, 2025 – 02:38 AM JST

PDB ID : 8ZR4 / pdb_00008zr4
EMDB ID : EMD-60391
Title : Cryo-EM structure of the N2-4N2C402 complex at a resolution of 1.9 angstrom
Authors : Chu, B.X.; Wang, X.; Yang, Q.; Wu, X.D.; Zhang, Z.L.
Deposited on : 2024-06-03
Resolution : 1.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	135	
1	F	135	
1	I	135	
1	L	135	
2	B	113	
2	G	113	
2	J	113	
2	M	113	
3	D	393	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	393	 87%11%
3	K	393	 87%11%
3	N	393	 88%10%
4	O	5	 40%60%
4	P	5	 40%60%
4	Q	5	 40%40%20%
4	R	5	 40%40%20%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 39296 atoms, of which 18770 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4N2C402_Fab_H.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	126	Total	C	H	N	O	S	0	0
			1929	616	936	179	194	4		
1	F	125	Total	C	H	N	O	S	0	0
			1913	611	927	178	193	4		
1	I	126	Total	C	H	N	O	S	0	0
			1929	616	936	179	194	4		
1	L	126	Total	C	H	N	O	S	0	0
			1929	616	936	179	194	4		

- Molecule 2 is a protein called 4N2C402_Fab_L.


Mol	Chain	Residues	Atoms						AltConf	Trace
2	B	111	Total	C	H	N	O	S	0	0
			1682	547	820	142	169	4		
2	G	110	Total	C	H	N	O	S	0	0
			1663	541	809	141	168	4		
2	J	110	Total	C	H	N	O	S	0	0
			1663	541	809	141	168	4		
2	M	110	Total	C	H	N	O	S	0	0
			1663	541	809	141	168	4		

- Molecule 3 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	D	386	Total	C	H	N	O	S	3	0
			5866	1860	2869	534	581	22		
3	H	386	Total	C	H	N	O	S	3	0
			5866	1860	2869	534	581	22		
3	K	386	Total	C	H	N	O	S	3	0
			5866	1860	2869	534	581	22		
3	N	386	Total	C	H	N	O	S	3	0
			5866	1860	2869	534	581	22		


- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran

ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					AltConf	Trace
4	O	5	Total	C	H	N	O	0	0
			113	34	52	2	25		
4	P	5	Total	C	H	N	O	0	0
			113	34	52	2	25		
4	Q	5	Total	C	H	N	O	0	0
			113	34	52	2	25		
4	R	5	Total	C	H	N	O	0	0
			113	34	52	2	25		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
5	D	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	D	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	H	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	H	1	Total	C	H	N	O	0
			27	8	13	1	5	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
5	K	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	K	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	N	1	Total	C	H	N	O	0
			27	8	13	1	5	
5	N	1	Total	C	H	N	O	0
			27	8	13	1	5	

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
6	D	1	Total	Ca	0
			1	1	
6	H	1	Total	Ca	0
			1	1	
6	K	1	Total	Ca	0
			1	1	
6	N	1	Total	Ca	0
			1	1	

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		AltConf
7	A	14	Total	O	0
			14	14	
7	B	22	Total	O	0
			22	22	
7	D	161	Total	O	0
			161	161	
7	F	13	Total	O	0
			13	13	
7	G	23	Total	O	0
			23	23	
7	H	164	Total	O	0
			164	164	
7	I	14	Total	O	0
			14	14	
7	J	22	Total	O	0
			22	22	
7	K	158	Total	O	0
			158	158	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		AltConf
7	L	15	Total 15	O 15	0
7	M	22	Total 22	O 22	0
7	N	161	Total 161	O 161	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4N2C402_Fab_H

Chain A: 



- Molecule 1: 4N2C402_Fab_H

Chain F: 




- Molecule 1: 4N2C402_Fab_H

Chain I: 




- Molecule 1: 4N2C402_Fab_H

Chain L: 



- Molecule 2: 4N2C402_Fab_L

Chain B: 




- Molecule 2: 4N2C402_Fab_L

Chain G: 



- Molecule 2: 4N2C402_Fab_L

Chain J: 86% 12% .



- Molecule 2: 4N2C402_Fab_L

Chain M: 86% 12% .



- Molecule 3: Neuraminidase

Chain D: 87% 11% .



- Molecule 3: Neuraminidase

Chain H: 87% 11% .




- Molecule 3: Neuraminidase

Chain K: 87% 11% .




- Molecule 3: Neuraminidase

Chain N: 88% 10% .




- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	663111	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/1013	0.26	0/1370
1	F	0.13	0/1006	0.27	0/1360
1	I	0.13	0/1013	0.27	0/1370
1	L	0.13	0/1013	0.27	0/1370
2	B	0.11	0/885	0.24	0/1206
2	G	0.11	0/877	0.24	0/1195
2	J	0.11	0/877	0.24	0/1195
2	M	0.11	0/877	0.24	0/1195
3	D	0.16	0/3069	0.37	0/4161
3	H	0.16	0/3069	0.37	0/4161
3	K	0.16	0/3069	0.38	0/4161
3	N	0.16	0/3069	0.37	0/4161
All	All	0.14	0/19837	0.33	0/26905

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	993	936	932	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	986	927	923	8	0
1	I	993	936	932	12	0
1	L	993	936	932	8	0
2	B	862	820	819	10	0
2	G	854	809	808	7	0
2	J	854	809	808	9	0
2	M	854	809	808	9	0
3	D	2997	2869	2878	29	0
3	H	2997	2869	2878	30	0
3	K	2997	2869	2878	31	0
3	N	2997	2869	2878	25	0
4	O	61	52	52	3	0
4	P	61	52	52	2	0
4	Q	61	52	52	3	0
4	R	61	52	52	3	0
5	D	28	26	26	1	0
5	H	28	26	26	3	0
5	K	28	26	26	3	0
5	N	28	26	26	2	0
6	D	1	0	0	0	0
6	H	1	0	0	0	0
6	K	1	0	0	0	0
6	N	1	0	0	0	0
7	A	14	0	0	1	0
7	B	22	0	0	1	0
7	D	161	0	0	3	0
7	F	13	0	0	0	0
7	G	23	0	0	1	0
7	H	164	0	0	4	0
7	I	14	0	0	1	0
7	J	22	0	0	0	0
7	K	158	0	0	2	0
7	L	15	0	0	0	0
7	M	22	0	0	0	0
7	N	161	0	0	3	0
All	All	20526	18770	18786	172	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 172 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:13:VAL:HG21	2:J:83:VAL:HG11	1.64	0.79
2:G:13:VAL:HG21	2:G:83:VAL:HG11	1.65	0.77
2:M:13:VAL:HG21	2:M:83:VAL:HG11	1.64	0.77
2:B:13:VAL:HG21	2:B:83:VAL:HG11	1.67	0.76
3:D:141:SER:OG	3:D:167:ASP:OD2	2.11	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/135 (87%)	115 (98%)	3 (2%)	0	100	100
1	F	117/135 (87%)	115 (98%)	2 (2%)	0	100	100
1	I	118/135 (87%)	116 (98%)	2 (2%)	0	100	100
1	L	118/135 (87%)	116 (98%)	2 (2%)	0	100	100
2	B	109/113 (96%)	108 (99%)	1 (1%)	0	100	100
2	G	108/113 (96%)	107 (99%)	1 (1%)	0	100	100
2	J	108/113 (96%)	107 (99%)	1 (1%)	0	100	100
2	M	108/113 (96%)	107 (99%)	1 (1%)	0	100	100
3	D	387/393 (98%)	374 (97%)	11 (3%)	2 (0%)	25	17
3	H	387/393 (98%)	372 (96%)	13 (3%)	2 (0%)	25	17
3	K	387/393 (98%)	370 (96%)	15 (4%)	2 (0%)	25	17
3	N	387/393 (98%)	373 (96%)	12 (3%)	2 (0%)	25	17
All	All	2452/2564 (96%)	2380 (97%)	64 (3%)	8 (0%)	45	29

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	146[A]	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	146[B]	ILE
3	H	146[A]	ILE
3	H	146[B]	ILE
3	K	146[A]	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/113 (94%)	103 (97%)	3 (3%)	38	33
1	F	105/113 (93%)	103 (98%)	2 (2%)	52	49
1	I	106/113 (94%)	105 (99%)	1 (1%)	75	77
1	L	106/113 (94%)	105 (99%)	1 (1%)	75	77
2	B	96/98 (98%)	96 (100%)	0	100	100
2	G	95/98 (97%)	95 (100%)	0	100	100
2	J	95/98 (97%)	95 (100%)	0	100	100
2	M	95/98 (97%)	95 (100%)	0	100	100
3	D	337/341 (99%)	335 (99%)	2 (1%)	84	86
3	H	337/341 (99%)	336 (100%)	1 (0%)	91	92
3	K	337/341 (99%)	336 (100%)	1 (0%)	91	92
3	N	337/341 (99%)	336 (100%)	1 (0%)	91	92
All	All	2152/2208 (98%)	2140 (99%)	12 (1%)	82	86

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	77	THR
1	I	132	THR
3	N	77	THR
3	K	77	THR
3	D	77	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	85	HIS
3	K	10	ASN
1	L	85	HIS
3	K	266	ASN
3	H	124	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	O	1	4	14,14,15	0.51	0	17,19,21	0.41	0
4	NAG	O	2	4	14,14,15	0.31	0	17,19,21	0.38	0
4	BMA	O	3	4	11,11,12	0.41	0	15,15,17	0.64	0
4	MAN	O	4	4	11,11,12	0.60	0	15,15,17	0.98	1 (6%)
4	MAN	O	5	4	11,11,12	0.63	0	15,15,17	0.90	1 (6%)
4	NAG	P	1	4	14,14,15	0.46	0	17,19,21	0.55	0
4	NAG	P	2	4	14,14,15	0.42	0	17,19,21	0.43	0
4	BMA	P	3	4	11,11,12	0.42	0	15,15,17	0.65	0
4	MAN	P	4	4	11,11,12	0.61	0	15,15,17	0.97	1 (6%)
4	MAN	P	5	4	11,11,12	0.64	0	15,15,17	0.91	1 (6%)
4	NAG	Q	1	4	14,14,15	0.85	1 (7%)	17,19,21	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	Q	2	4	14,14,15	0.43	0	17,19,21	0.40	0
4	BMA	Q	3	4	11,11,12	0.43	0	15,15,17	0.64	0
4	MAN	Q	4	4	11,11,12	0.59	0	15,15,17	0.99	1 (6%)
4	MAN	Q	5	4	11,11,12	0.64	0	15,15,17	0.90	1 (6%)
4	NAG	R	1	4	14,14,15	0.84	1 (7%)	17,19,21	0.61	0
4	NAG	R	2	4	14,14,15	0.44	0	17,19,21	0.40	0
4	BMA	R	3	4	11,11,12	0.42	0	15,15,17	0.63	0
4	MAN	R	4	4	11,11,12	0.58	0	15,15,17	0.99	1 (6%)
4	MAN	R	5	4	11,11,12	0.64	0	15,15,17	0.90	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	O	1	4	-	2/6/23/26	0/1/1/1
4	NAG	O	2	4	-	0/6/23/26	0/1/1/1
4	BMA	O	3	4	-	0/2/19/22	0/1/1/1
4	MAN	O	4	4	-	0/2/19/22	0/1/1/1
4	MAN	O	5	4	-	0/2/19/22	0/1/1/1
4	NAG	P	1	4	-	1/6/23/26	0/1/1/1
4	NAG	P	2	4	-	0/6/23/26	0/1/1/1
4	BMA	P	3	4	-	0/2/19/22	0/1/1/1
4	MAN	P	4	4	-	0/2/19/22	0/1/1/1
4	MAN	P	5	4	-	0/2/19/22	0/1/1/1
4	NAG	Q	1	4	-	0/6/23/26	0/1/1/1
4	NAG	Q	2	4	-	0/6/23/26	0/1/1/1
4	BMA	Q	3	4	-	0/2/19/22	0/1/1/1
4	MAN	Q	4	4	-	0/2/19/22	0/1/1/1
4	MAN	Q	5	4	-	0/2/19/22	0/1/1/1
4	NAG	R	1	4	-	0/6/23/26	0/1/1/1
4	NAG	R	2	4	-	0/6/23/26	0/1/1/1
4	BMA	R	3	4	-	0/2/19/22	0/1/1/1
4	MAN	R	4	4	-	0/2/19/22	0/1/1/1
4	MAN	R	5	4	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	Q	1	NAG	O5-C1	-3.08	1.38	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	R	1	NAG	O5-C1	-3.04	1.38	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	Q	4	MAN	C1-O5-C5	2.32	115.33	112.19
4	R	4	MAN	C1-O5-C5	2.30	115.30	112.19
4	O	4	MAN	C1-O5-C5	2.27	115.26	112.19
4	P	4	MAN	C1-O5-C5	2.18	115.15	112.19
4	P	5	MAN	O2-C2-C3	-2.18	105.77	110.14

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	P	1	NAG	C1-C2-N2-C7
4	O	1	NAG	C4-C5-C6-O6
4	O	1	NAG	O5-C5-C6-O6


There are no ring outliers.


4 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	P	1	NAG	2	0
4	O	1	NAG	3	0
4	R	1	NAG	3	0
4	Q	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	H	401	-	14,14,15	0.19	0	17,19,21	0.54	0
5	NAG	K	402	-	14,14,15	0.29	0	17,19,21	0.51	0
5	NAG	N	401	-	14,14,15	0.19	0	17,19,21	0.55	0
5	NAG	N	402	-	14,14,15	0.27	0	17,19,21	0.51	0
5	NAG	D	401	-	14,14,15	0.19	0	17,19,21	0.56	0
5	NAG	D	402	-	14,14,15	0.27	0	17,19,21	0.54	0
5	NAG	H	402	-	14,14,15	0.31	0	17,19,21	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	K	401	-	14,14,15	0.19	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	H	401	-	-	2/6/23/26	0/1/1/1
5	NAG	K	402	-	-	0/6/23/26	0/1/1/1
5	NAG	N	401	-	-	2/6/23/26	0/1/1/1
5	NAG	N	402	-	-	0/6/23/26	0/1/1/1
5	NAG	D	401	-	-	4/6/23/26	0/1/1/1
5	NAG	D	402	-	-	2/6/23/26	0/1/1/1
5	NAG	H	402	-	-	0/6/23/26	0/1/1/1
5	NAG	K	401	-	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	401	NAG	C8-C7-N2-C2
5	D	401	NAG	O7-C7-N2-C2
5	D	402	NAG	C8-C7-N2-C2
5	D	402	NAG	O7-C7-N2-C2
5	H	401	NAG	C4-C5-C6-O6



There are no ring outliers.



6 monomers are involved in 9 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	401	NAG	2	0
5	K	402	NAG	1	0
5	N	401	NAG	2	0
5	D	401	NAG	1	0
5	H	402	NAG	1	0
5	K	401	NAG	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.